

MATH1081 notes

Nira (z5417727)

September 17th, 2022

Contents

1	Topic 1	2
1.1	Introduction	2
1.2	Sets and subsets	3
1.3	Power Sets and Stability	4
1.4	Set Operations	5
1.5	The Inclusion-Exclusion Principle	6
1.6	Sets Proofs	7
1.7	Laws of Set Algebra	8
1.8	Generalised Set Operations	9
1.9	Russel's Paradox	10
1.10	Cartesian Product	11
1.11	Functions	12
1.12	Image and Inverse Image	13
1.13	Injective, Surjective, Bijective	14
1.14	Composition of Functions	15
1.15	Identity and Inverse Functions	16

1 Topic 1

1.1 Introduction

1. addition, multiplication, division and subtraction
2. Mainly dealing with finite sets

1.2 Sets and subsets

A set is a well defined collection of distinct objects

Example: $S = \{1, a, 3\}, A = \{\Pi, 1\}$.

1. $e \notin A$; it is not in A
2. For example, if A is a set of all integers; $\{\text{all even integers}\} = \{n \in \mathbb{R} | n \text{ is even}\}$.
3. We can remove superfluous items (elements that occur more than one).
 $A = \{1, 2, 3, 3\}$ where 3 can be removed.

Example:

$A = \{1, 2, 3\}, B = \{2, 3, 1\}, C = \{1, 2, 3, 3\}, D = \{1, 3\}$.

Here, D is a proper subset of A, B, C; A, B, C are supersets of D.

\subseteq : Subset (proper subset), \supseteq : Superset.

1. To prove if a set is a proper subset; do the following:

For example, if $D \in A$, then check if $e \in D$

If $e \in D$, then $e \in A$. Thus, it would be a proper subset (here, e is just an element).

2. To prove that two sets are equal;

For example, if $A = B$, prove:

- i) $A \subseteq B$; if an element is in A, then the element is in B.
- ii) $B \subseteq A$; if an element is in B, then the element is in A.

1.3 Power Sets and Stability

Subsets of $A = \{1, 2, 3\}$:

1. Could throw everything out to get empty set Φ ,
2. One element each: $\{1\}, \{2\}, \{3\}$,
3. Two elements: $\{1, 2\}, \{2, 3\}, \{1, 3\}$,
4. Set itself: A .

The set containing 1, 2, 3, 4 is called the powerset of A.

Given $A = \{1, 2, 3\}, B = \{1, 2, 3, 3\}, C = \{1, 3\}, D = \{1, 3\}$, where $A = B, C \subseteq A, B$ and $D \not\subseteq A, B, C$.

1. size of A = 3, B = 3, C = 2, D = 2.

[Exercise with A = 0, 1, 0, 1, B done in word].

1.4 Set Operations

Boolean Operators ("not" operation in programming):

1. Complement:

Let there be a set A in U (A : all of the people in the video, U : universal set of everyone in the world, A^c = complement of A).

$$A^c = \{x \in U | x \notin A\}.$$

2. Intersecting ("and" operation in programming):

If there is A, B , intersecting,

$$A \cap B = \{x \in A | x \in B\}.$$

3. Union ("or" operation in programming): If there is A, B , A or B is:

$$A \cup B = \{x \in U | x \in A \text{ or } x \in B\}.$$

4. Difference: If there is A, B , intersecting,

$$A - B = \{x \in A | x \notin B\}.$$

[examples in word doc]

1.5 The Inclusion-Exclusion Principle

[example in Word]

$$|A \cup B| = |A| + |B| - |A \cap B|.$$

For three elements,

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|.$$

[example in word]

1.6 Sets Proofs

[proof question in word]

Hints for proofs:

1. To prove that $S \subseteq T$, we can assume that $x \in S$ and show that $x \in T$.
2. To prove that $S = T$, we can show that $S \subseteq T$ and $T \subseteq S$.

Scaffold:

Proof: Suppose that (proof) we see that/ it follows ... (conclusion) (end with shaded box to indicate end of proof)

Note that the "Suppose that" part of the proof is usually whatever the if statement mentions.

For example, if the question is "Prove that if $A \cap B = A$, then $A \cup B = B$ ", then the proof starts like this:

<u>Proof</u> : Suppose that $A \cap B = A$.
--

For questions like "is this statement true", there are two ways to approach the question:

1. If the statement is true (if you think it is true), then prove it.
2. If the statement is false, then give a counter-example that proves it false.

[examples in word]

1.7 Laws of Set Algebra

Laws of Set Algebra

1. $A \cap B = B \cap A$: Commutative Law.
2. $A \cap (B \cap C) = (A \cap B) \cap C$: Associative Law.
3. $A \cap (B \cap C) = (A \cap B) \cup (A \cap C)$: Distributive Law.
4. $A \cap (A \cup B) = A$: Absorption Law.
5. $A \cap U = U \cap A = A$: Identity Law.
6. $A \cap A = A$: Idempotent Law.
7. $(A^c)^c = A$: Double Complement Law.
8. $A \cap \emptyset = \emptyset \cap A = \emptyset$: Domination Law.
9. $A \cap A^c = \emptyset$: Intersection with Complement Law.
10. $(A \cup B)^c = A^c \cap B^c$: De Moirve's Law.

The intersection can be swapped with the union to form another law (like, $A \cup B = B \cup A$ swapped as $A \cap B = B \cap A$). Similarly, U should be swapped with \emptyset and vice versa.

[examples in word]

1.8 Generalised Set Operations

Unions and Intersections; A saga:

$$1. \cup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n,$$

$$2. \cap_{i=1}^n A_i = A_1 \cap A_2 \cap \cdots \cap A_n.$$

Example:

$$\begin{aligned} A_k &= k, k+1; \\ \cup_{i=1}^3 A_k &= A_1(\{1, 2\}) \cup A_2(\{2, 3\}) \cup A_3(\{3, 4\}), \\ &= \{1, 2, 3, 4\}. \end{aligned}$$

[example in word]

1.9 Russel's Paradox

A set may contain another set as one of its elements.

This raises the possibility that a set may contain itself as an element.

Problem: Try to let S be the set of all sets that are not elements of themselves, i.e., $S = \{A \mid A \text{ is a set and } A \notin A\}$.

Is S an element of itself?

i) If $S \in S$, then the definition of S implies that $S \notin S$, a contradiction.

ii) If $S \notin S$, then the definition of S implies that $S \in S$, also a contradiction.

Hence neither $S \in S$ nor $S \notin S$. This is Russell's paradox.

1.10 Cartesian Product

[example in word]

The Cartesian product of two sets A and B, denoted by $A \times B$, is the set of all ordered pairs from A to B:

$$A \times B = \{(a, b) | a \in A \text{ and } b \in B\}$$

If $|A| = m$ and $|B| = n$, then we have $|A \times B| = mn$.

Sets with more than 2 elements:

Example: $A = \{a, b\}, B = \{1, 2, 3\}$.

Cartesian Product $(A \times B) = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$

(all of the ordered pairs – combinations)

[example in word]

When X and Y are small finite sets, we can use an arrow diagram to represent a subset S of $X \times Y$: we list the elements of X and the elements of Y , and then we draw an arrow from x to y for each pair $(x, y) \in S$.

1.11 Functions

Example: Take 2 sets X and Y , for which we have to find a function.

$$X = \{\text{all MATH 1081 students}\}, Y = \{0, 1, \dots, 84, 85, \dots, 100\}.$$

X : number of students; Y : marks from 0 – 100.

Take function $f : X \rightarrow Y$; where X is the domain and Y is the co domain.

Ie, $f(x)$ = X 's mark (Y).

Function $f : X \rightarrow Y$ satisfies $\{(x, f(x)) | x \in X\} \subseteq X \times Y$ so that, for each $x \in X$;

1. $f(x)$ exists
2. $f(x)$ is unique

[example in word]

Note: be vary of the one-to-one function property lol

Floor function and ceiling functions:

1. Floor function (rounds down; smallest integer):

$$\lfloor x \rfloor = \max \{z \in \mathbb{Z} | z \leq x\}.$$

2. Ceiling function (rounds up; largest integer):

$$\lceil x \rceil = \min \{z \in \mathbb{Z} | z \geq x\}.$$

[example in word] Domain/codomain: $\lfloor x \rfloor / \lceil x \rceil : \mathbb{R} \rightarrow \mathbb{Z}$.

Range($\lceil x \rceil$) = \mathbb{Z} .

[example in word]

1.12 Image and Inverse Image

- The image of a set $A \subseteq X$ under a function $f : X \rightarrow Y$ is $f(A) = \{y \in Y \mid y = f(x) \text{ for some } x \in A\} = \{f(x) \mid x \in A\}$.

- The inverse image of a set $B \subseteq Y$ under a function $f : X \rightarrow Y$ is $f^{-1}(B) = \{x \in X \mid f(x) \in B\}$.

(image is just function values in the domain and inverse image is function values in range).

note: this is just function and inverse functions.
--

[example in word]

1.13 Injective, Surjective, Bijective

Formal Definitions:

Recall that if f is a function from X to Y , then for every $x \in X$, there is exactly one $y \in Y$ such that $f(x) = y$.

1. We say that a function $f : X \rightarrow Y$ is injective or one-to-one if, for every $y \in Y$, there is at most one $x \in X$ such that $f(x) = y$.

Example: for all $x_1, x_2 \in X$, if $f(x_1) = f(x_2)$ then $x_1 = x_2$.

2. We say that a function $f : X \rightarrow Y$ is surjective or onto if, for every $y \in Y$, there is at least one $x \in X$ such that $f(x) = y$. the range of f is the same as the codomain of f ($\text{range}(f) = Y$).

3. We say that a function $f : X \rightarrow Y$ is bijective if f is both injective and surjective (one-to-one and onto).

for every $y \in Y$, there is exactly one $x \in X$ such that $f(x) = y$.

[example in word]

1.14 Composition of Functions

For functions $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, the composite of f and g is the function $g \circ f : X \rightarrow Z$ defined by $(g \circ f)(x) = g(f(x))$ for all $x \in X$.

The composite function $g \circ f$ exists whenever the range of f is a subset of the domain of g .

In general, $g \circ f$ and $f \circ g$ are not the same composite functions. Associativity of composition (assuming they exist): $h \circ (g \circ f) = (h \circ g) \circ f$.

Example: Take sets $X = \{ \text{all MATH1081 students} \}$, $Y = \{0, 1, \dots, 100\}$, $Z = \{F, P, CR, D, HD\}$.

Maps: $f : X \rightarrow Y$; $g : Y \rightarrow Z$.

A) $g \circ f : X \rightarrow Z$.
 $(f \circ g)(y) = f(g(y))$.
[examples in word]

1.15 Identity and Inverse Functions