

02

Reduction

Metal hydrides reduction (LiAlH_4 ; NaBH_4 ; DiBAL-H ; B_2H_6):

Certain complex metal and boron hydrides are important reagents for reduction.

(i) LiAlH_4 (LAH) Lithium aluminium hydride [LiAlH_4 /Ether or THF] :


LAH is most common and versatile reagent. It is sensitive to protic solvent and therefore used in ether.

Reagent	LiAlH_4	LiAlH_4	LiAlH_4 /Excess	LiAlH_4 /Excess	LiAlH_4	LiAlH_4	LiAlH_4	LiAlH_4	LiAlH_4
Reactant	Aldehyde	Ketone	Acid	Acid anhydride	Acid chloride	Ester	Cyanide	Amide	Nitro
Product	1° alcohol	2° alcohol	1° alcohol	1° alcohol	1° alcohol	1° alcohol	1° amine	1° amine	1° amine

NOTE :

(i) Alkene, alkyne, benzene rings are not reduced by LiAlH_4 .

(ii) double bond can be reduced by LiAlH_4 / THF in few cases like :



[Cinnamaldehyde]

Mechanism of LiAlH_4








Illustration 1



NOTE : It can reduce 1° and 2° alkyl halide into Alkane but not 3°






(ii) Sodium borohydride [NaBH₄ / C₂H₅OH or Ether] :

It is more specific than LAH as a reducing agent. It reduces ketones and aldehydes to the corresponding alcohols without affecting other functional groups, reduces acid chlorides to 1° alcohols. It is effective even in protic solvent like alcohol.

Reagent	NaBH ₄	NaBH ₄	NaBH ₄
Reactant	Aldehyde	Ketone	Acid chloride
Product	1° alcohol	2° alcohol	1° alcohol



NOTE :


NaBH₄ can also reduce imine group ($\text{C}=\text{NH}$).

(iii) Diisobutyl Aluminium Hydride [DIBAL-H / Inert solvent] :



Diisobutyl aluminium hydride is parallel to LAH (Lithium aluminium hydride) as a reducing agent but it is more selective.

Reduction of acyl halides, esters and nitriles :

(a) Acyl chlorides can be reduced to aldehydes by treating them with lithium-tri-tert-butoxyaluminium hydride, LiAlH[OC(CH₃)₃]₃ or with DIBAL-H at -78°C.




(b) Both esters and nitriles can be reduced to aldehydes by DIBAL-H. Reduction must be carried out at low temperatures. Hydrolysis of the intermediates gives the aldehyde.



Reagent	DiBAL-H/(-78°C)		
Reactant	Ester	Cyanide	Acyl chlorides
Product	Aldehyde	Aldehyde	Aldehyde

Mechanism of DIBAL H



Reduction

NOTE :

By DIBAL at ordinary temperature esters are reduced to alcohols but at low temperature esters are reduced to aldehyde.




Illustration 2





Illustration 3

LAH reduce RCN to amine but DIBAL is found to be reduce it to aldehyde.



(iv) B₂H₆ or BH₃/THF

B₂H₆/THF gives BH₃ which is a electrophile(Lewis acid) so B₂H₆ does not reduce electrophilic functional groups.

NOTE : -

Acyl chloride and esters are relatively electron poor (Cl and OR are very electronegative). Boron will not reduce acyl chloride and reduces esters slowly.

Reactant	Remark	Product
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{X}$	Not usually Reduced	
R - NO ₂	Not usually Reduced	
R - X	Not usually Reduced	
$\text{>C}=\text{N}-\text{OH}$	Not usually Reduced	
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$	Reduced slowly	-CH ₂ OH
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-$	Reduced slowly	$\begin{array}{c} \text{OH} \\ \\ -\text{CH}- \end{array}$
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OR}$	Reduced slowly	R - CH ₂ OH + ROH
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Reduced	-CH ₂ OH
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2$	Reduced	R-CH ₂ -NH ₂



NOTE :

1. B_2H_6 in THF not able to reduce cyclic ester.

Catalytic Hydrogenation:


From alkenes and alkynes (Sabatier and Senderens reaction)

Alkenes and alkynes on catalytic hydrogenation gives alkanes.



Catalyst :

- (a) Pd/Pt at ordinary temp. and pressure
- (b) Ni, 200–300° C (sabatier)
- (c) Raney Nickel at room temp.
- (d) Raney nickel is obtained by boiling Ni/Al with NaOH. Al dissolved & Ni obtained in finally divided state.
- (e) Methane cannot be prepared by this method (From unsaturated hydrocarbon).





Higher electrophilicity more reactive for hydrogenation.

They do not reduce $-COOH$.


Alkenes and alkynes on catalytic hydrogenation gives alkanes

Illustration 4




This method is called Sabatier–senderens reaction or hydrogenation of alkenes and alkynes

Selective reduction by H_2 -Pd/C



Greater the crowding at double bond carbon more difficult the hydrogenation

Rate of Hydrogenation




Mendius reduction (complete reduction)

NOTE : -


Alkyl cyanides (nitriles) can be reduced to 1° amine and isocyanides are reduced to 2° amine .

Reagent $-H_2 + Na/ROH$.



Reduction


Illustration 5




By partial reduction of Alkynes -

By catalytic Hydrogenation of Alkynes in presence of poisoned catalyst (A Syn Addition of Hydrogen : Synthesis of cis-Alkenes : This is performed by)


(i) **Lindlar's catalyst (Pd/CaCO₃)** : Metallic palladium deposited on calcium carbonate conditioned with lead acetate and quinoline.




(ii) **Rosenmund catalyst (Pd/BaSO₄)** : Metallic palladium deposited on Barium sulfate conditioned with lead acetate and quinoline.



(iii) **P-2 catalyst (Ni₂B nickel boride)**



Mechanism of hydrogenation :




Steps : The reactant alkyne molecules and hydrogen molecules get adsorbed at the surface of metal catalyst. It is chemical adsorption (chemisorption).

In this state, the reactants lie very close to each other and so the hydrogen atoms start forming bond with carbon. Two hydrogen atoms are added to two triply bonded carbon atoms from the same side of π bond and a **cis or syn addition product** is formed. The product alkene now escapes away from the surface of the catalyst. Quinoline occupies the metal surface inhibiting further reduction to alkanes **Quinoline** therefore is called catalyst poison and such palladium is called deactivated catalyst or poisoned catalyst.

NOTE : Rosenmund's reduction :

Quinoline or sulphur act as a poisoned catalyst, controls the further reduction of aldehyde to alcohols.



This reaction controls further reduction of aldehyde to alcohols.

In case if only Pd is used, formation of alcohol takes place.



Formaldehyde cannot be prepared by this method because HCOCl is highly unstable at room temperature it decompose into $\text{CO} + \text{HCl}$.




Illustration 6





Illustration 7



Wilkinson's catalyst :


Reagent - $[(\text{Ph}_3\text{P})_3\text{RhCl}] + \text{H}_2$

It gives Syn addition.

Reduce only alkene or alkyne.

METAL ACID REDUCTION:

From alkyl Halides (By reduction) :



Catalyst :



(i) $\text{Zn} + \text{HCl}$

(ii) $\text{Zn} + \text{CH}_3\text{COOH}$



(iii) $\text{Zn}-\text{Cu}$ couple in $\text{C}_2\text{H}_5\text{OH}$

(iv) $\text{Al} + \text{Hg} + \text{ethanol}$

Mechanism :



Product



Key points:

(a) Alkyl halides can also be reduced to alkane by H_2/Pd or LiAlH_4 or H_2/Ni .


(b) Reduction is due to the electron transfer from the metal to the substrate ($\text{R}-\text{X}$)

(c) If any alkyl halide is asked, the H-atom of any carbon atom of given alkane is removed by halogen atom.

The Wolf Kishner reduction:


When a ketone or an aldehyde is heated in a basic solution of hydrazine, the carbonyl group is converted to a methylene group this process is called Deoxygenation because an oxygen is removed from the reactant. The reaction is known as the Wolf-kishner Reduction.

Reagent : $\text{N}_2\text{H}_4/\text{OH}/\Delta$



Reduction

Function : Aldehyde/Ketone to Alkane




NOTE :

It reduce aldehydes and ketones to alkanes, but does not reduce alkene and alkyne.


This is a base catalysed reaction, so simultaneously dehydrohalogenation of RX also take place.

Mechanism :



Reduction by transfer hydrogenation:

Reagent $\text{N}_2\text{H}_4/\text{H}_2\text{O}_2$ Or N_2H_2



Mechanism:





Illustration 8





Illustration 9




Clemmensen's Reduction ; Mozingo Reduction:

The reduction of carbonyl groups of aldehydes and ketones to methylene groups with amalgamated zinc and concentrated hydrochloric acid is known as Clemmensen reduction.

Reagent : Zn-Hg/Conc. HCl

Function : Aldehyde/Ketone to Alkane




In this reaction $-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-$ is converted into $-\text{CH}_2-$


NOTE :-

If other acid sensitive group is also attached then H^+ will react with that group and If we want retain acid-sensitive group in the product then we will not use clemensen Reduction.


Mechanism




Mozingo Reaction :



Birch Reduction:




Mechanism




Reduction


Birch reduction in conjugated double bond




Mechanism:




Birch Reduction In Benzene Ring



Mechanism:




CASE-I $\Rightarrow$ Presence of electron releasing group ($-\text{R}$, $-\text{OR}$, $-\text{NH}_2$ etc.)




G = E.W.G

Mechanism:



CASE-II $\Rightarrow$ Presence of electron withdrawing group ($-\text{NO}_2$, $-\text{COOH}$, $-\text{CHO}$, $-\text{CN}$ etc.)



G = E.D.G

Mechanism:




Illustration 10






Illustration 11




Rate of reaction with Na/liq NH₃



(iii) > (i) > (ii)

Illustration 12

Identify the reagent for following synthesis



Solution:

H₂/Lindlar's catalyst






Illustration 13

Identify the product in the following reaction :




Solution:



Reduction

Bouveault Blank Reduction:

Reduces all which are reduced by LAH except -COOH



Mechanism:









Illustration 14




Stephen's Reduction (SnCl₂/HCl)

Partial reduction




MPV Reduction:








In this reaction, carbonyl is reduced into corresponding alcohol by aluminium isopropoxide taken in isopropanol.

Reduction with (Red P + HI):









Mechanism:











All organic compound containing O convert into Alkane Containing N Convert into amine




No. of moles of (Red P + HI) required for Reduction of alcohol, aldehyde, ketone, carboxylic acid in presence of Red P and HI




In the above reaction I_2 is formed which acts as reducing agent and may reduce alkane and form alkyl halide. So red P is added in the reaction to remove I_2 formed in the reaction.



(i) Alcohol




No. of moles of (Red P + HI) required = 2




No. of moles of (Red P + HI) required = 4

(ii) Ketone



No. of moles of (Red P + HI) required = 4

(iii) Carboxylic Acid



No. of moles of (Red P + HI) required = 6

Illustration 15




Illustration 16




Illustration 17




Illustration 18



Illustration 19



Solution:






Illustration 20



Solution:






Illustration 21



Solution:








Illustration 22



Solution:




(ii) No reaction

Illustration 23




Reduction

Solution:




(ii) No reaction

Illustration 24




Solution:



(ii) No reaction

Illustration 25



Solution:




Illustration 26













Illustration 27



Chart-1

Group	Product	LAH in Ether	NaBH ₄ in H ₂ O	B ₂ H ₆ in THF	H ₂ /Catalyst Δ
-CHO	-CH ₂ OH	+	+	+	+
>C = O	>CH - OH	+	+	+	+
-CO ₂ H	-CH ₂ OH	+	-	+	-
-CO ₂ R	-CH ₂ OH	+	-	+	+
-COCl	-CH ₂ OH	+	+	-	+
-CONH ₂	-CH ₂ NH ₂	+	-	+	+
(RCO) ₂ O	RCH ₂ OH	+	-	+	+
-CN	-CH ₂ NH ₂	+	-	+	+
>C = NOH	-CH ₂ NH ₂	+	-	-	+
>C = C<	>CH - CH<	-	-	+	+
-C ≡ C-	-CH = CH-	-	-	+	+
1° RX	RH	+	-	-	+
2° RX	RH	+	+	-	+
3° RX	RH	-	+	-	+

Chart-2

Name	Reagent	Function
Wolf Kishner Reduction	(i) N ₂ H ₄ / (ii) KOH, Δ	
Clemenson Reduction	Zn-Hg / HCl	
Mozingo Reduction	 Dry HCl, followed by Raney Ni	
Stephen's Reduction	SnCl ₂ / HCl followed by H ₃ O ⁺	R-C≡N → R-CH = O
MPV Reduction		
Hydroboration Reduction	B ₂ H ₆ / AcOH, H ₂ O	 
Bouvoult Blank Reduction	Na / EtOH	R-COO-R → RCH ₂ OH + ROH
Transfer Hydrogenation	N ₂ H ₄ / H ₂ O ₂	

Rosenmund Reduction	$\text{H}_2, \text{Pd-BaSO}_4$	$\text{R}-\overset{\text{O}}{\underset{\text{ }}{\text{C}}}-\text{Cl} \longrightarrow \text{R}-\overset{\text{O}}{\underset{\text{ }}{\text{C}}}-\text{H}$ $^*-\text{C}\equiv\text{C}- \longrightarrow \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ -\text{C}=\text{C}- \\ \quad \\ \text{H} \quad \text{H} \end{array}$
Birch Reduction	$\text{Na} / \text{Liq. NH}_3$	$-\text{C}\equiv\text{C}- \longrightarrow \begin{array}{c} \text{H} \\ \\ -\text{C}=\text{C}- \\ \\ \text{H} \end{array}$
DIBAL-H (-78°C)	$\text{H}-\text{Al}\left(\begin{array}{c} \text{CH}_2-\text{CH}-\text{CH}_3 \\ \\ \text{CH}_3 \end{array}\right)_2$ followed by $\text{H}_2\text{O}^\oplus$	$\left. \begin{array}{l} -\text{COOR} \\ -\text{C}\equiv\text{N} \\ -\text{COCl} \\ \begin{array}{c} \text{O} \quad \text{O} \\ \quad \\ -\text{C}-\text{O}-\text{C}- \end{array} \end{array} \right\} \longrightarrow -\text{CHO}$ $-\text{CHO} \longrightarrow -\text{CH}_2\text{OH}$
Red phosphorus in presence of HI	Red P + HI	$\text{R}-\text{CO}_2\text{H} \longrightarrow \text{RCH}_3$ $\text{R}-\text{CH}=\text{O} \longrightarrow \text{RCH}_3$ $\text{R}-\overset{\text{O}}{\underset{\text{ }}{\text{C}}}-\text{R} \longrightarrow \text{RCH}_2\text{R}$ $\text{R}-\text{OH} \longrightarrow \text{R}-\text{H}$