

01

Periodic Table

Introduction:

The arrangement of all the known elements according to their properties in such a way that the elements with similar properties are grouped together in a tabular form is called periodic table.

Development of periodic table:

(A) Lavoisier's Classification:

(i) Lavoisier classified the elements simply in metals and non metals.

Metals are the ones which have the tendency of losing electrons.



Non-metals are the ones which have the tendency of gaining electrons.



(ii) Drawback or Limitation :

(a) As the number of elements increases, this classification became insufficient for the study of elements.

(b) There are few elements which have the properties of both metals as well as non-metals and they are called metalloids. Lavoisier could not decide where to place the metalloids.

(B) Dobereiner's triad rule [1817]

(i) He made groups of three elements having similar chemical properties called TRIAD.

(ii) In Dobereiner triad, atomic weight of middle element is nearly equal to the average atomic weight of first and third element.

	Element	At. Weight	Mean weight of first and last element
(i)	Li	7	
	Na	23	$\frac{7 + 39}{2} = 23$
	K	39	
(ii)	Ca	40	
	Sr	88	$\frac{40 + 137}{2} = 88.5$
	Ba	137	
(iii)	Cl	35.5	
	Br	80	$\frac{35.5 + 127}{2} = 81.25$
	I	127	
(iv)	S	23	
	Se	79	$\frac{32 + 127.6}{2} = 79.8$
	Te	127.6	

(iii) Other examples – (K, Rb, Cs), (P, As, Sb), (H, F, Cl), (Sc, Y, La)

- As only few elements were discovered at the time so only few such triads were available.
- Day by day as new elements are discovered, the rule could no longer be generalized.

Practice Question

- Q. The law of triads is applicable to -
 (1) C, N, O (2) H, O, N (3) Na, K, Rb (4) Cl, Br, I Ans. (4)
- Q. The law of triads is not applicable on -
 (1) Cl, Br, I (2) Na, K, Rb (3) S, Se, Te (4) Ca, Sr, Ba Ans. (2)

(C) Newlands law of octaves [1865]

- (i) He arranged the elements in the increasing order of their atomic mass and observed that properties of every 8th element was similar to the 1st one, like in the case of musical vowels notation.

Sa	Re	Ga	Ma	Pa	Dha	Ne	Sa
1	2	3	4	5	6	7	8

- (ii) At that time inert gases were not known.

Li	Be	B	C	N	O	F	
Na	Mg	Al	Si	P	S	Cl	
K	Ca						

- (iii) The properties of Li are similar to 8th element i.e. Na, Be are similar to Mg and so on.

Drawback or Limitation :

- (a) This rule is valid only upto Ca because after Ca due to presence of d-block elements there is difference of 18 elements instead of 8 elements.
- (b) After the discovery of Inert gas which when included in the periodic table becomes the 8th element from Alkali metal so this law had to be dropped.
- (c) This law found to be applicable only till calcium. It was not applicable to elements of higher atomic masses.

Practice Question

- Q. Which of the following set of elements obeys Newlands octave rule -
 (1) Na, K, Rb (2) F, Cl, Br (3) Be, Mg, Ca (4) B, Al, Ga Ans. (3)
- Q. For which of the pair Newlands octave rule is not applicable -
 (1) Li, Na (2) C, Si (3) Mg, Ca (4) Cl, Br Ans. (4)

(D) Lothar Meyer's curve [1869]

- (i) He plotted a curve between atomic weight and atomic volume of different elements.
- (ii) The following observation can be made from the curve -
- Most electropositive elements i.e. alkali metals (Li, Na, K, Rb, Cs etc.) occupy the peak positions on the curve.
 - Less electropositive i.e. alkaline earth metals (Be, Mg, Ca, Sr, Ba) occupy the descending position on the curve.
 - Metalloids (B, Si, As, Te, At etc.) and transition metals occupy bottom part of the curve.
 - Most electronegative i.e. halogens (F, Cl, Br, I) occupy the ascending position on the curve.

Note : Elements having similar properties occupy similar position on the curve.

Conclusion :


On the basis of this curve Lothar Meyer proposed that the physical properties of the elements are periodic function of their atomic weight and this become the base of Mendeleev's periodic table.

Illustration 1:

What is periodic function ?

Solution:

When the elements are arranged in the increasing order of their atomic mass (weight), elements having similar properties gets repeated after a regular interval.

**(E) Mendeleev's Periodic Table [1869]:**

- Mendeleev's periodic law – The physical and chemical properties of elements are the periodic function of their atomic weight.
- Characteristic of Mendeleev's periodic table –
 - It is based on atomic weight
 - 63 elements were known; noble gases were not discovered.
 - He was the first scientist to classify the elements in a systematic manner i.e. in horizontal rows and in vertical columns.
 - Horizontal rows are called periods and there were 7 periods in Mendeleev's Periodic table.
 - Vertical columns are called groups and there were 8 groups in Mendeleev's Periodic table.
 - Each group upto VIIth is divided into A & B subgroups. 'A' sub group elements are called normal elements and 'B' sub group elements are called transition elements.
 - The VIIIth group consists of 9 elements in three rows.
 - The elements belonging to same group exhibit similar properties.

(iii) Merits or advantages of Mendeleev's periodic table –

- (a) Study of elements – For the first time, all known elements were classified in groups according to their similar properties. So, study of the properties of elements became easier.
- (b) Prediction of new elements – It gave encouragement to the discovery of new elements as some gaps were left in it.

Sc (Scandium), Ga (Gallium), Ge (Germanium), Tc (Technetium)

were the elements whose position and properties were well defined by Mendeleev even before their discoveries and he left the blank spaces for them in his table.

e.g. - Blank space at atomic weight 72 in silicon group was called Eka silicon (means properties like silicon) and element (discovered later) was named Germanium.

Similarly other elements discovered after Mendeleev periodic table were :

Eka aluminium – Gallium (Ga)

Eka Boron – Scandium (Sc)

Eka Silicon – Germanium (Ge)

Eka Manganese – Technetium (Tc)

- (c) Correction of doubtful atomic weights – Corrections were done in atomic weight of some elements.

$$\text{Atomic Weight} = \text{Valency} \times \text{Equivalent weight}$$

Initially, it was found that equivalent weight of Be is 4.5 and it is trivalent (Valency = 3), so the weight of Be was 13.5 and there is no space in Mendeleev's table for this element. So, after correction, it was found that Be is actually divalent (Valency = 2). So, the weight of Be became $2 \times 4.5 = 9$ and there was a space between Li and B for this element in Mendeleev's table.

Corrections were done in atomic weight of elements are – U, Be, In, Au, Pt.

(iv) Demerits of Mendeleev's periodic table –

- (a) Position of hydrogen – Hydrogen resembles both, the alkali metals (IA) and the halogens (VIIA) in properties so Mendeleev could not decide where to place it.
- (b) Position of isotopes – As atomic weight of isotopes differs, they should have been placed in different position in Mendeleev's periodic table. But there was no such place for isotopes in Mendeleev's table.
- (c) Anomalous pairs of elements – There were some pair of elements which did not follow the increasing order of atomic weight.

eg : Ar and Co were placed before K and Ni respectively in the periodic table, but having higher atomic weights.


$\left(\begin{array}{cc} \text{Ar} & \text{K} \\ 39.9 & 39.1 \end{array} \right)$	$\left(\begin{array}{cc} \text{Te} & \text{I} \\ 127.5 & 127 \end{array} \right)$
$\left(\begin{array}{cc} \text{Th} & \text{Pa} \\ 232 & 231 \end{array} \right)$	$\left(\begin{array}{cc} \text{Co} & \text{Ni} \\ 58.9 & 58.6 \end{array} \right)$

- (d) Like elements were placed in different groups.

There were some elements like Platinum (Pt) and Gold (Au) which have similar properties but were placed in different groups in Mendeleev's table.

Pt	Au
VIII	IB

- (e) Unlike elements were placed in same group.




Cu, Ag and Au placed in 1st group along with Na, K etc. While they differ in their properties (Only similar in having ns^1 electronic configuration)

(f) It was not clear that 'lanthanides and Actinides' are related with IIIA group or IIIB group.

(g) Cause of periodicity : Why physical & chemical properties repeated in a group.

(F) Moseley's Experiment and Modern periodic law:

Moseley did an experiment in which he bombarded high speed electron on different metal surfaces and obtained characteristic X-rays.



Moseley's Experiment

He found out that $\sqrt{\nu} \propto Z$ (where ν = frequency) of characteristic X-rays from this experiment, Moseley concluded that the physical and chemical properties of the elements are periodic function of their atomic number. It means that when the elements are arranged in the increasing order of their atomic number, elements having similar properties gets repeated after a regular interval. This became the base of 'Modern Periodic Law'.

(i) **Modern periodic law** – The physical & chemical properties of elements are a periodic function of their atomic number.

(ii) Characteristics of modern periodic table –

(a) It was proposed by Moseley (1913).

(b) Modern periodic table is based on atomic number.

- (c) 9 vertical columns called groups.
- (d) These groups from I to VII are further subdivided into subgroups A and B, IA to VIIA, IB to VIIB, VIII and 0
- (e) Inert gases were introduced in periodic table by Ramsay in group 0.
- (f) 7 horizontal series called periods.

(G) Niels Bohr's Classification (Based on modified Mendeleev's periodic table)

Elements can be divided in four parts on the basis of electronic configuration.

(i) Inert gas elements –

- (a) The elements in which outermost shell or ultimate shell is completely filled up are called inert gas elements.
- (b) General electronic configuration is ns^2p^6 (Except He = $1s^2$)
- (c) Because of the most stable configuration, they are very less reactive. Hence known as inert gases.

(ii) Normal or Representative elements –

The elements in which outermost shell or ultimate shell is incomplete, and no. of electrons in the ultimate shell is less than eight while penultimate shell is complete are called normal or representative elements.


Or

The elements of s-block & p-block except inert gases are called normal or representative elements.

(iii) Transition elements –

The elements in which both ultimate (n) as well as penultimate shells (n – 1) are incomplete either in ground state or in any oxidation state are called transition elements.

According to this concept, Zn, Cd, Hg are not transition elements because they do not have incomplete penultimate shell either in ground state or in +2 oxidation state.




Note :

All transition elements are d-block elements but all d-block elements are not transition elements.

(iv) Inner transition elements –

- (a) The Elements in which all the three shells that is ultimate (n) penultimate (n-1) and pre or antepenultimate (n – 2) shell are incomplete are called inner transition elements.



- (b) There are 28 elements are known as inner transition elements.
- (c) Inner transition elements are divided into two series.

(i) Lanthanide series/Rare earth elements /Lanthanoids ($\text{Ce}_{58} - \text{Lu}_{71}$ 14 elements)

The first element of this series is Cerium, not Lanthanum.

In these elements, last electron enters into 4f subshell

They are present in IIIB group and 6th period of the periodic table.

Promethium (${}_{61}\text{Pm}$) is the only lanthanide which is synthetic and radioactive in nature.

(ii) Actinide series /Man made elements /Actinoids ($\text{Th}_{90} - \text{Lw}_{103}$ 14 elements)

The first element of this series is Thorium, not Actinium.

In these elements, last electron enters into 5f subshell.

They are present in IIIB group and 7th period of the periodic table.

All the actinides are radioactive in nature.

First three elements (Th, Pa, U) are found in nature while others are synthetic in nature.

After U₉₂ i.e. from ₉₃Np onwards elements are called transuranic elements because :

They are heavier than uranium.

They are derived from uranium by nuclear reactions.

Illustration 2:

Classify the following elements based on Bohr's concept :

(1) [Ar] 4s², 3d⁸

(2) 1s², 2s², 2p⁶, 3s², 3p⁶

(3) [Xe] 6s², 5d¹, 4f¹²

(4) [Rn] 7s², 6d⁴, 5f¹⁴

(5) [Kr] 5s², 4d¹⁰, 5p²

Solution:

(1) Transition

(2) Inert gas

(3) Inner transition

(4) Transition

(5) Normal element

Illustration 3:

The heaviest naturally occurring element is?


Solution:

Uranium.

Some Commonly Used Terms:


- Noble Gases** : Element of group 18 are called noble gases. These are also called as inert gases because their outermost ns and np orbitals are completely filled (except He has 1s²) and these gases are non-reactive in nature under ordinary conditions.
- Typical elements** : Elements of 3rd period (Na, Mg, Al, Si, P, S, Cl) are called typical elements because they represent the properties of other element of their respective group.
- Diagonal relationship** : Properties of elements of second period (Li, Be, B) resemble with the element of third period (Mg, Al, Si). This resemblance between properties of 2nd & 3rd period is called diagonal relationship.

2nd period elements show diagonal relationship with 3rd period elements because of same ionic potential (Ionic potential = Charge/Radius) value they show similarity in properties.



4. Bridge elements:

The typical elements of third period are also called bridge elements as the division between two subgroups A and B starts from these elements. In second group Mg acts as a bridge element. The properties of bridge element are somewhat mixed of the elements of two subgroups as magnesium shows similarities with alkaline earth metals (IIA) on one hand and with zinc metals (IIB) on the other.



Q. Which of the following elements belong to alkali metals ?

(1) $1s^2, 2s^2, 2p^2$

(2) $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2, 3d^{10}, 4p^6, 5s^1$

(3) $1s^2, 2s^2, 2p^5$

(4) None of these

Ans. (2)

Important Points:

(a) Normal elements present in all periods.

(b) Long form of modern periodic table can be divided into four portions.

1. Left portion – IA & IIA → s - block.

2. Right portion – IIIA to VII A + 0 group → p - block.

3. Middle portion – IIIB to VIIB + VIII + IB IIB → d - block.

4. Bottom portion – IIIB → f - block elements

(c) The group containing most electro positive elements – IA GROUP.

(d) The group containing most electro negative elements – VIIA GROUP

(e) The group containing maximum number of gaseous elements – ZERO (18th) GROUP

(f) "0" (zero) or 18 group have all the elements in gaseous form.

(g) The group in which elements have generally ZERO valency – (18th) GROUP ZERO

(h) In the periodic table –

Number of Gaseous elements – 11 (H, N, O, F, Cl + Noble gases)

Number of Liquid elements – 6 (Cs, Fr, Ga, Hg, Br, Uub)

Bromine is the only non-metal which exists in liquid form (at room temperature).

Number of Solid elements – 95 (if discovered elements are 112)

(i) No p-block elements in 1st period.

(j) 1st period has all the elements in gaseous form (H, He)

(k) 2nd period contains maximum number of gaseous elements. They are 4 : N, O, F, Ne

(l) IIIB/3rd group is called largest group having 32 elements including 14 Lanthanides and 14 Actinides

Sc

Y

La.....Lanthanides (14)

Ac.....Actinides (14)

Q. Properties of Li is similar to :

(1) Mg

(2) Na

(3) Both

(4) None of these

Ans. (3)

Q. 6th typical element is

Ans. Sulphur

Q. Lanthanides belong to which group of the table ?

(1) IIIA

(2) IIIB

(3) 3

(4) both 2 and 3

Ans. (4)

(H) Long form / Present form of Modern Periodic Table:

(It is also called as 'Bohr-Bury & Rang-Werner Periodic Table.)

(i) It is based on the Bohr-Bury concept of electron distribution.

(ii) This model was proposed by Rang & Werner.

(iii) It consists of 7 horizontal periods and 18 vertical columns (groups)

(iv) According to I. U. P. A. C., 18 vertical columns are named as 1st to 18th group.

(v) The co-relation between the groups in long form of periodic table and in modern periodic table are given below.

IA	IIA	IIIB	IVB	VB	VIB	VIIIB	VIIIB	VIIIB	VIIIB	IB	IIB	IIIA	IVA	VA	VIA	VIIA	0(Zero)
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18

(vi) Elements belonging to same group have same no. of electrons in the outermost shell so their properties are similar.

Description of periods (Based on modern periodic table):

Period	n	Sub shell	No. of elements	Element	Name of Period
1.	1	1s	2	${}_1\text{H} - {}_2\text{He}$	Shortest
2.	2	2s, 2p	8	${}_3\text{Li} - {}_{10}\text{Ne}$	Short
3.	3	3s, 3p	8	${}_{11}\text{Na} - {}_{18}\text{Ar}$	Short
4.	4	4s, 3d, 4p	18	${}_{19}\text{K} - {}_{36}\text{Kr}$	Long
5.	5	5s, 4d, 5p	18	${}_{37}\text{Rb} - {}_{58}\text{Xe}$	Long
6.	6	6s, 4f, 5d, 6p	32	${}_{55}\text{Cs} - {}_{86}\text{Rn}$	Longest
7.	7	7s, 5f, 6d, 7p	32	${}_{87}\text{Fr} - {}_{118}\text{Og}$	Incomplete

Conclusion

- Period number = outermost shell = maximum value of principle quantum number (n) in electronic configuration.
- Number of elements in a period = Number of electrons in a period

Description of Groups:

1st /Alkali metals

H = $1s^1$

Li = $1s^2, 2s^1$

Na = $1s^2, 2s^2, 2p^6, 3s^1$

K = $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^1$

General electronic configuration = ns^1

Number of valence shell $e^- = 1$

13th/Boron Family

B = $1s^2, 2s^2, 2p^1$

2nd/Alkaline earth metals

Be = $1s^2, 2s^2$

Mg = $1s^2, 2s^2, 2p^6, 3s^2$

Ca = $1s^2, 2s^2, 2p^6, 3s^2, 3p^6, 4s^2$



General electronic configuration = ns^2

(n = Number of shell)

Number of valence shell $e^- = 2$

14th/Carbon Family

C = $1s^2, 2s^2, 2p^2$






General electronic configuration = $ns^2 np^1$

Number of valence shell $e^- = 3$

15th/Nitrogen family/Pnictogen

(Used in fertilizer as urea)






General electronic configuration = $ns^2 np^3$

Number of valence shell $e^- = 5$



17th/Fluorine family/Halogens

(Salt forming)



General electronic configuration = $ns^2 np^5$

Number of valence shell $e^- = 7$






General electronic configuration = $ns^2 np^2$

Number of valence $e^- = 4$

16th/Oxygen family/Chalcogen

(Ore forming)





General electronic configuration = $ns^2 np^4$

Number of valence shell $e^- = 6$

18th/ Noble gas family /Inert gases /Noble gases

(Less reactive)



General electronic configuration = $ns^2 np^6$ (except He)

Number of valence shell $e^- = 8$

Metals, Non-Metals & Metalloids:

Apart from classifying elements into s, p, d and f-blocks, there is yet another broad classification of elements based on their properties. The elements can be broadly classified into

(a) Metals:

Majority of the elements in periodic table are metals and appears on the left side of the periodic table.

Properties:

- (i) These are usually solid at room temperature [exception - mercury]
- (ii) They have high melting and boiling point [exception Gallium & Caesium have very low melting point (303 K and 302 K respectively)]
- (iii) They are good conductor of heat and electricity.
- (iv) They are malleable (can be flattened into thin sheets by hammering) and ductile (can be drawn into wires)

(b) Non-Metals:

These are placed at the top right-hand side of periodic table. As we move horizontally along a period, the property of elements changes from metallic (on left) to non-metallic (on the right).

Properties :

- (i) These are usually solids or gases at room temperature.
- (ii) They have low melting point and boiling point (exceptions : Boron, Carbon).
- (iii) Most Non-metallic solids are brittle and are neither malleable nor ductile.

(c) Metalloids (Semi-metals) :

Properties of these elements show the characteristics of both metals and non-metals. Mainly Silicon (Si), Germanium (Ge), Arsenic (As), Antimony (Sb) and Tellurium (Te) are metalloids.

Estimating Position of an Element from its Electronic Configuration:

The last electron enters in which subshell gives idea of its block.

[☞ Think : $1s^1$ and $1s^2$ belongs to which block]

Period number = Principal quantum number of valence shell electron in ground state electronic configuration.

Group number for s block = number of valence shell electrons

Group number for p block = 10 + number of valence shell electrons

Group number for d block = number of $[ns + (n-1) d]$ electrons

Group number for f-block = 3

[☞ Use these carefully while locating the position.]

- Merits of long form of periodic table –**

- Position of isotopes – Atomic No. of isotopes are similar, so different isotopes can be placed at same place in periodic table.
- (Ar – K) (Co – Ni) (Te – I) are now in increasing order of atomic number.
- Lanthanides and actinides are in IIIB group or group 3.
- In this periodic table 2 diagonal lines separates out metals, metalloids and non-metals.
(metalloids Si, Ge, As, Sb, Te)
- Elements of same group have same general formula of electronic configuration of outer most shell.

- Demerits of long form of periodic table –**

- Position of hydrogen is still controversial.
- 'He' is an inert gas but it has different electronic configuration than other inert gas elements.
- Lanthanides and actinides are still not placed in main frame.
- Isotopes have different physical properties but have same place in periodic table.

Practice Question

- Q. Each period in the periodic table starts with a subshell of new shell and ends with –
 (1) Small subshell (2) Next higher shell Ans. (3)
 (3) p-subshell of the same shell (4) Different subshell of the same subshell
- Q. The number of elements in long period is –
 (1) 18 (2) 8 (3) 32 (4) 10 Ans. (1)
- Q. Long form of the periodic table is based on –
 (1) Atomic size (2) Atomic number (3) Atomic mass (4) Atomic volume Ans. (2)
- Q. Maximum work on periodic table related with atomic number was given by –
 (1) Moseley (2) Mendeleev (3) Aston (4) Hund Ans. (1)

Classification on the Basis of Sub shell in which last e^- enters:**(i) s - block elements –**

- In these elements last electron enters in s - subshell.
- Group – IA(1) and IIA(2) group
- Period – 1 to 7th
- Electronic configuration – ns^{1-2}
 IA group $n = 1$ to 7
 IIA group $n = 2$ to 7
- Total s - block elements are (14) (including H and He)
- Total s - block metals are (12) (excluding H)
- He is a s-block element as last e^- enters in s-subshell.

General characteristics of s-block elements :

- (a) s-block elements are soft metal, that is why these elements have low melting and boiling point.
- (b) Their oxides and hydroxides are basic in nature.
- (c) They act as reducing agent as they have tendency to donate e^- .
- (d) They have very low ionisation energy and highly electropositive so, they form ionic compound.
- (e) Flame of these elements have the property of showing different colour.

(ii) p - block elements –

- (a) Last e^- enters in p - sub shell
- (b) Group – IIIA (13) to VIIA (17) + 0 group (18) (except He)
- (c) He is a s-block element but placed in p-block zero or 18th group as it is a noble gas element.
- (d) Period – 2nd to 7th
- (e) Electronic configuration – $ns^2 np^{1-6}$
- (f) Total p - block elements – (36)

Characteristic of p- block elements :

- (a) p-block possesses all three kind of elements i.e. metals, non-metals and metalloids.
- (b) Oxides of non-metals are acidic in nature.
- (c) They form covalent compounds mostly.
- (d) They are oxidising in nature.

(iii) d - block elements –

- (a) Last e^- enters in $(n - 1)$ d subshell
- (b) Group – IIIB – VIIB, VIII, IB, IIB or group 3 to 12 (IUPAC)
- (c) Period – 4th to 7th
- (d) Electronic configuration – $(n - 1) d^{1-10} ns^{1 \text{ or } 2}$
- (e) Total d - block elements – (40) (If 112 elements are included in periodic table)
- (f) IIB elements (Zn, Cd, Hg) are d-block elements but not transition elements.

General characteristics of d - block elements –

- (a) They all are metals, which are very hard having high melting and boiling point (except Hg).
- (b) Elements of this block situated in between 's' and p-block elements. So, the character of this block elements lie between s and p-block elements
- (c) They show variable oxidation - state. eg. $Mn \Rightarrow Mn^{2+}, Mn^{3+}, Mn^{4+}, Mn^{5+}, Mn^{6+}, Mn^{7+}$
- (d) They form ionic and covalent bond both.
- (e) They are good conductor of heat and electricity and form complex compounds.
- (f) Metals, which have unpaired electrons show para-magnetism.
- (g) They form alloys and most of the elements act as catalyst.

(iv) f - block elements –

- (a) Differentiating e^- enters in $(n - 2)$ f subshell.
- (b) Group – IIIB or 3rd
- (c) Period – 6th and 7th
- (d) From at. no. 58 - 71, 6th period ; Lanthanide series $4f^{1-14} 5d^0 \text{ or } 1 6s^2$
90 - 103 | 7th period ; Actinide series $5f^{1-14} 6d^{0-2} 7s^2$
- (e) Total number of f block elements – (28)
- (f) All the actinides are radioactive elements.
- (g) Transuranic actinides are man made elements. (Np_{93} – Lw_{103})
- (h) Lanthanides are found rarely on earth so these are called rare earth metals.

General characteristics of f-block elements –

- (a) All the f-block elements are heavy metals.
- (b) They have high melting and boiling point.
- (c) The most common oxidation state of these elements is +3.
- (d) The elements of 5f series are radioactive in nature.

Determination of period, block and group of an element –

(i) Period No. – The period no. of the element can be predicted from the principal quantum no. (n) of the valence shell.

e.g. electronic configuration of iodine is :

$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^5$. Therefore, the period number of iodine is 5 as the valence shell configuration is $5s^2 5p^5$.

(A) When atomic number is given:

Step I : $71 \geq Z \geq 58 \Rightarrow$ Lanthanoids (6th Period) } f-block
 $103 \geq Z \geq 90 \Rightarrow$ Actinoids (7th Period)

Group number = IIIB or Group 3 (largest group of periodic tables)

Step II : $Z = 104$ to 118 (Period number = 7)

Group number = last two digits in atomic number of elements

Example: $Z = 104$

Group no. = 4

Step III: Group number = $18 + \text{given atomic number} - \text{atomic number of next noble gas}$

If the value of this formula is negative then use 32 instead of 18 in formula.

(B) When electronic configuration is given

Period number (n) = number of outermost shell/Highest shell number.

(ii) Block No. – The type of orbital which receives the last electron known as block no.

e.g. An element 'X' has its electronic configuration as $1s^2 2s^2 2p^6 3s^2 3p^4 4s^2 3d^8$. As the last electron enters in the d-orbital, therefore it is a d-block element.

Block identification:

- If np electron present then p - block ($ns^2 np^{1-6}$)
group number = $12 + np$ electrons
- If np electron absent then s/f/d block
 - If $(n-2)f^0 (n-1)d^0 ns^{1-2} = s$ block
group number = ns electrons
 - If $(n-2)f^{1-14} (n-1)d^{0-2} ns^2 = f$ block
group number = IIIB
- If any other configuration or $(n-1)d^{1-10} ns^{0-2}$ (d-block)
group number = (n-1)d electron + ns electron

(iii) Group No. – It is predicted from the number of electrons in the valence shell and penultimate shell as follows.

- (a) For s-block elements, group number is equal to the number of electrons in the valence shell after the noble gas core.
e.g. An element 'Y' having electronic configuration $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2$ or $[\text{Ar}] 4s^2$ has two electrons in valence shell and it is a s-block element. Therefore, it belongs to group 2.
- (b) For p-block element, the group number is equal to 10 + number of electrons in valence shell.
e.g. An element 'Z' with electronic configuration as $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^3$ has five electrons in its valence shell and belongs to p-block.
Therefore, its group number is $10 + 5 = 15$. It belongs to group VA of the Mendeleev's periodic table.
- (c) For d-block elements group number is equal to the number of electrons in $(n-1)d$ sub-shell and valence shell.
e.g. An element 'A' having electronic configuration as $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$. so, its group number will be $10 + 1 = 11$.

Predicting atomic no. of the successive member in a group or family –

Knowing the atomic no. of the first member of a group, we can write the atomic no. of the subsequent elements by adding given magic no. For example

H ₁	1 + 2 = 3	Li	3 + 8 = 11	Na	11 + 8 = 19	K	19 + 18 = 37	Rb	37 + 18 = 55	Cs	55 + 32 = 87
Magic no.	2		8		8		18		18		32

(ii) Other than group IA-

Group	1	2	3	4, 5, 6, 7, 8, 9, 10, 11, 12	13, 14, 15, 16, 17	18
	IA	IIA	IIIB		IIIA IVA VA VIA VIIA	'0'gp
1 st	2	-				8
2 nd	8	8			8	8
3 rd	8	8			18	18
4 th	18	18	18	18	18	18
5 th	18	18	18	32	32	32
6 th	18	18	18	32	32	32
7 th	32	32	32	32	-	-

Lanthanide series
Total elements = 14

Practice Question

- Q. In the periodic table the elements $Z = 30$ is expected to be present in –
(1) IA series (2) IB group (3) IIA group (4) IIB group Ans. (4)
- Q. Elements A, B, C, D and E have the following electronic configurations :-
(A) $1s^2, 2s^2 2p^1$ (B) $1s^2, 2s^2 2p^6, 3s^2 3p^1$ (C) $1s^2, 2s^2 2p^6, 3s^2 3p^3$
(D) $1s^2, 2s^2 2p^6, 3s^2 3p^5$ (E) $1s^2, 2s^2 2p^6, 3s^2 3p^6, 4s^1$
The elements which belong to the same group of the periodic table are :-
(1) (A) and (C) (2) (A) and (B) (3) (A) and (D) (4) (A) and (E) Ans. (2)

Q. Uranium is a member of –

(1) Transition series (2) III-period (3) Actinide series (4) VI-period Ans. (3)

Q. In modern periodic table 6th period contain elements –

(1) 8 (2) 18 (3) 10 (4) 32 Ans. (4)

Nomenclature of Elements with atomic number > 100:

(a) IUPAC gave names to elements above atomic number 100 as follows –

Digit	0	1	2	3	4	5	6	7	8	9
Name	nil	un	bi	tri	quad	pent	hex	sept	oct	enn
Abbreviation	n	u	b	t	q	p	h	s	o	e

(b) In all the elements suffix is – ium.

Ex. Atomic No.	IUPAC Name	Symbol	Elemental Name	Symbol
101	Unnilunium	Unu	Mendelevium	Md
102	Unnilbium	Unb	Nobelium	No
103	Unniltrium	Unt	Lawrencium	Lr
104	Unnilquadium	Unq	Rutherfordium	Rf
105	Unnilpentium	Unp	Dubnium	Db
106	Unnilhexium	Unh	Seaborgium	Sg
107	Unnilseptium	Uns	Bohrium	Bh
108	Unniloctium	Uno	Hassium	Hs
109	Unnilennium	Une	Meitnerium	Mt
110	Ununnilium	Uun	Darmstadtium	Ds

Screening effect or Shielding effect (σ) and effective nuclear charge (Z_{eff}):

- Valence shell e^- suffer force of attraction due to nucleus and force of repulsion due to inner shell electrons.
- The decrease in force of attraction on valence e^- due to inner shell e^- is called screening effect or shielding effect. (i.e. total repulsive force is called shielding effect.)
- Due to screening effect valence shell e^- experiences less force of attraction exerted by nucleus.
i.e. total attraction force experienced by valence electrons represented by a number is Z_{eff} .
- There is a reduction in nuclear charge due to screening effect. Reduced nuclear charge is called effective nuclear charge.
- If nuclear charge = Z , effective nuclear charge = Z_{eff} , σ (Sigma) = Screening constant or shielding constant. So, $Z_{\text{eff}} = (Z - \sigma)$

Slater's rule to know screening or shielding constant (σ)

- For single electron species $\sigma = 0$
- Screening effect (S.E.) for two e^- species
Ex. In He ($1s^2$)
Screening effect of one $1s e^-$. where $\sigma = 0.30$
 $\therefore Z_{\text{eff}} = Z - \sigma = 2 - 0.30 = 1.7$
- Screening effect of each ns and np (Outermost shell) electrons is 0.35
- Screening effect of each $(n - 1)$ penultimate shell s, p, d electrons are 0.85
- Screening effect of each $(n - 2)$ and below all the e^- present in s, p, d, f is 1.0
From top to bottom in a group Z_{eff} remain constant
- Screening effect of each nd & nf electron is 0.35 and for all the e^- below nd & nf is 1.

Group	Element	Li	Na	K	Rb	Cs	Fr
	Z_{eff}	1.30	2.20	2.20	2.20	2.20	2.20
Period	Element	Be	B	C	N	O	F
	Z_{eff}	1.95	2.6	3.25	3.90	4.55	5.20

- For same shell shielding effect has the order as $s > p > d > f$ (due to penetration effect)
 - Z_{eff} for different ions of an element
 $Z_{\text{eff}} \uparrow$ as Positive charge $\uparrow$, and $Z_{\text{eff}} \downarrow$ as Negative charge $\uparrow$
 Ex. $\text{N}^{\oplus} > \text{N} > \text{N}^{\ominus} = Z_{\text{eff}}$
 - Z_{eff} for isoelectronic species (When arranged in increasing order of atomic no.)
 $Z_{\text{eff}} \uparrow$ as Atomic no. $\uparrow$
 Ex. $\text{H}^{\ominus} < \text{Li}^{\oplus} < \text{Be}^{2\oplus} < \text{B}^{3\oplus}$ ($2e^{\ominus}$ species)
 $\text{N}^{3\ominus} < \text{O}^{2\ominus} < \text{F}^{\ominus} < \text{Na}^{\oplus} < \text{Mg}^{2\oplus}$ ($10e^{\ominus}$ species)

Periodicity in Periodic Properties:

- The regular gradation in properties from top to bottom in a group and from left to right in a period is called periodicity in properties.
- In a period, the ultimate shell remains same, but the number of electrons gradually increases.
- In a group, the number of electrons in the ultimate shell remains same, but the values of n increases.

Cause of Periodicity

- The cause of periodicity in properties is due to the same outermost shell electronic configuration repeating over regular intervals.
- In the periodic table, elements with similar properties occur at intervals of 2, 8, 8, 18, 18 and 32.

Periodic Properties:

Atomic Radius:

- It is distance between outermost $e^{\ominus}$ and nucleus.
- Half of the nuclear distance between two atoms is defined as atomic radius.
- X-ray diffraction, $e^{\ominus}$ diffraction method and nuclear magnetic resonance (NMR) spectrum methods are used to determine internuclear distance or bond length.

$$\text{Atomic radius} = \frac{\text{Inter nuclear distance}}{2}$$

- Atomic radius depends on the type of chemical bond between atoms in a molecule. These are :

(i) Covalent radius (ii) Metallic radius (iii) Van der Waals radius

(i) Covalent radius :

Covalent radius is the half of the internuclear distance between two singly bonded atoms of the same element.

e.g. If internuclear distance of $\text{A}-\text{A}(\text{A}_2)$ molecule is $(d_{\text{A}-\text{A}})$ and covalent radius is r_{A} then

$$d_{\text{A}-\text{A}} = r_{\text{A}} + r_{\text{A}} \quad \text{or} \quad 2r_{\text{A}}$$

$$r_{\text{A}} = \frac{d_{\text{A}-\text{A}}}{2}$$

e.g. - In Cl_2 molecule, internuclear distance is $1.98\text{\AA}$ so $r_{\text{Cl}} = \frac{1.98\text{\AA}}{2} = 0.99\text{\AA}$


(ii) Metallic radius :

- Half of the internuclear distance between two adjacent metallic atoms in crystalline lattice structure.
- Metallic radius $>$ Covalent radius

(iii) Van der Waals radius:

- Those atoms (like noble gases) which are not bonded with each other and for gaseous molecules Van der Waals radius is measured.
- Half of the distance between the nuclei of adjacently placed atoms in solid state of a noble gas/gaseous molecules is Van der Waals radius.

- (c) Van der Waals radius > Covalent radius.
 (d) Inert gas has only Van der Waals radius.
 (e) In molecules of nonmetals both covalent and Van der Waals radius exists.
 (f) For a same atom $VWR > \text{Metallic } R > \text{Covalent } R$



For a same element

Covalent radius < Metallic radius < Van der Waals radius

(iv) Factors affecting atomic size are:

- | | |
|---|---|
| (a) Atomic size ↓ when Effective nuclear charge ↑ | Li > Be > B > C > N > O > F |
| (b) Atomic size ↑ when No. of shells ↑ | Li < Na < K < Rb < Cs |
| (c) Atomic size ↓ when Screening effect ↓ | Li > Be > B > C > N > O > F |
| (d) Atomic size ↓ when Magnitude of +ve charge ↑ | Mn > Mn ²⁺ > Mn ³⁺ > Mn ⁴⁺ |
| Atomic size ↑ when Magnitude of -ve charge ↑ | O < O ⁻ < O ²⁻ |

(v) Periodic variation of atomic size:

- (i) Across a period – It decreases from left to right in a period as nuclear charge increases
 e.g. Li > Be > B > C > N > O > F < Ne
 e.g. Ar > Na > Cl
- (ii) In a group – It increases from top to bottom in a group as number of shell increases
 e.g. Li < Na < K < Rb < Cs
 e.g. Be < Li < Na

(vi) Lanthanide and Actinide Contraction :-

- (a) Outermost electronic configuration of inner transition elements is
 $(n-2)f^{1-14}, (n-1)s^2p^6d^{0-1}, ns^2$ ($n = 6$ or 7)
- (b) e^- enters in $(n-2)$ f orbitals
- (c) Mutual screening effect of e^- is very less, because of complex structure of f-orbital
- (d) Due to poor shielding of 4f and 5f electrons, effective nuclear charge (Z_{eff}) increases in lanthanides and actinides so atomic size of these elements moving from left to right slightly decreases. It is known as lanthanide contraction in 4f series and actinide contraction in 5f series.
 Actinide contraction is stronger than lanthanide contraction because size of 5f orbitals is larger than 4f orbital so shielding power of 5f is less effective than that of 4f.

Practice Questions

Q. From the given set of species, point out the species from each set having least atomic radius :

- (1) O^{2-} , F^- , Na^+ (2) Ni, Cu, Zn (3) Li, Be, Mg (4) He, Li^+ , H^-

Correct answer is :-

- (1) O^{2-} , Cu, Li, H^- (2) Na^+ , Ni, Be, Li^+
 (3) F^- , Zn, Mg, He (4) Na^+ , Cu, Be, He Ans. (2)

Q. Which of the following pairs of elements have almost similar atomic radii :

- (1) Zr, Hf (2) Mo, W (3) Co, Ni (4) All Ans. (1)

Q. If the ionic radii of K^+ and F^- are nearly the same (i.e. 1.34 Å) then the atomic radii of K and F respectively are :

- (1) 1.34 Å, 1.34 Å (2) 0.72 Å, 1.96 Å
 (3) 1.96 Å, 0.72 Å (4) 1.96 Å, 1.34 Å Ans. (2)

Ionic Radius :

(A) Cationic radius:

(a) When a neutral atom loses e^- it converts into cation (+ve charged ion)

Atomic radius > Cationic radius

because after losing e^- no. of e^- reduces, but no. of protons remains same, due to this Z_{eff} increases, hence electrons are pulled towards nucleus and atomic radius decreases, moreover after losing all the electrons from the outer most shell, penultimate shell becomes ultimate shell which is nearer to nucleus so size decreases.

(b) Size of cation $\uparrow$ when Magnitude of the (+)ve charge $\downarrow$ or $Z_{eff} \downarrow$

e.g. $Fe > Fe^{2+} > Fe^{3+}$

$Pb^{2+} > Pb^{4+}$

$Mn > Mn^{2+} > Mn^{3+} > Mn^{4+} > Mn^{5+} > Mn^{6+} > Mn^{7+}$

(B) Anionic radius :-

(a) When neutral atom gains e^- it converts into an anion.

Anionic radius > Atomic radius

(b) In an anion e^- are more than protons so effective nuclear charge reduces, and inter electronic repulsion increases, which also increases screening effect. So, distance between e^- and nucleus increases and size of anion also increases.

Size of anion $\uparrow$ when Magnitude of the (-)ve charge $\uparrow$ or $Z_{eff} \downarrow$

e.g. Z of fluorine is 9

	F	F^-
Proton	9	9
e^-	9	10

so $\frac{Z}{e} = \frac{9}{9} = 1$; As Z_{eff} of F^- is less than F so size of $F^- > F$

(C) Size of iso protonic species:

- Those species having same no. of protons but different number of electrons i.e. ions formed by the same atom.
- for isoprotonic species the size increases with increase in electrons

	N^+	N	N^-	N^{2-}	N^{3-}
Z	7	7	7	7	7
e	6	7	8	9	10

Order of radius : ($N^{\oplus} < N < N^{\ominus} < N^{2\ominus} < N^{3\ominus}$)

($H^{\oplus} < H < H^{\ominus}$), ($I^{\oplus} < I < I^{\ominus}$)

For same atom, Cationic radius < Atomic radius < Anionic radius

(D) Size of iso electronic species:

- Those species having same no. of e^{-} but different nuclear charge forms isoelectronic series.
- for isoelectronic species the atomic radius decreases with increase in nuclear charge


	$S^{2\ominus}$	$Cl^{\ominus}$	$K^{\oplus}$	$Ca^{2\oplus}$
Z	16	17	19	20
e	18	18	18	18

Order of radius : ($S^{2\ominus} > Cl^{\ominus} > K^{\oplus} > Ca^{2\oplus}$),

($N^{3\ominus} > O^{2\ominus} > F^{\ominus} > Na^{\oplus} > Mg^{2\oplus} > Al^{3\oplus}$), ($Na^{\oplus} < Ne < F^{\ominus}$)

Note :

In the direction of arrow ($\rightarrow$) ionic size increases.



Atomic radius trends for d and f block elements :

In a period :

For transition elements of 3d series

- From $_{21}Sc$ to $_{30}Zn$ in first transition series Z_{eff} continuously increases, but atomic size does not decrease continuously.
- From $_{21}Sc$ to $_{26}Fe$ size decreases
Nuclear charge > Screening effect
- From $_{26}Fe$ to $_{28}Ni$ size almost constant
Nuclear charge $\sim$ Screening effect
- From $_{28}Ni$ to $_{30}Zn$ size increases. As nuclear charge is less effective than screening effect.

Final order in 3d series : $Sc > Ti > V > Cr < Mn > Fe \geq Co \approx Ni < Cu < Zn$

In a group :

- The atomic radius of elements increases on moving down the first transition series (3d) to second transition series (4d). This is due to the increases in number of shells with the increase in atomic number.
- The atomic radii of second (4d) and third (5d) transition series in a group is almost same except $Y(39)$ and $La(57)$

(iii) In transition series 1st, 2nd and 3rd


size increases down the groups

III_B Sc
 $\downarrow$ Y
 $\downarrow$ La

IV_B Ti
 $\downarrow$ Zr
 $\downarrow$ Hf

Almost equal due to lanthanide contraction

Radii $3d < 4d \approx 5d$ except IIIrd B.



$$\text{Sc} < \text{Y} < \text{La}$$

$$\text{Cu} < \text{Au} < \text{Ag}$$

$$\text{Ti} < \text{Zr} \approx \text{Hf}$$

For inner transition elements:

As we move along the lanthanide series, there is a decrease in atomic as well as ionic radius. The decrease in size is regular in ions but not so regular in atoms. This is due to lanthanide contraction*.

Exceptions :




- (1) Noble gases have largest atomic sizes [Van der Waals radii] in their respective periods. However, their covalent radii are smaller (only Xe show covalent radii)
- (2) Size of Al > Ga, [Z_{eff} increasing] (d-contraction or scandide contraction)
- (3) Size of Hf & Zr are same (lanthanide contraction)

Ionisation Potential Or Ionization Energy Or Ionization Enthalpy:

- (i) **Minimum energy** required to remove most loosely held outer most shell e^- in ground state from an isolated gaseous atom is known as Ionisation Energy.
 (Isolated $\rightarrow$ Without any bonding with other atom)

(ii) Successive I.E.

- (a) For an atom M, successive ionisation energies are as follows -



$$\text{IE}_3 > \text{IE}_2 > \text{IE}_1$$

- (b) e^- cannot be removed from solid state of an atom, it has to convert in gaseous form, Energy required for conversion from solid state to gaseous state is called Sublimation energy.
- (c) IE is always an endothermic process ($\Delta H = +ve$)
- (d) It is measured in eV/atom (electron volt/atom) or kCal/mole or kJ/mole

(iii) Factors affecting ionisation energy:

(A) Atomic size : Larger the atomic size, smaller is the I.E. It is due to that the size of atom increases the outermost electrons e^- farther away from the nucleus and nucleus loses the attraction on that electrons and hence can be easily removed. (Generally, down the group I.E. decreases)

I.E. $\uparrow$ when Atomic size $\downarrow$

(B) Effective nuclear charge (Z_{eff}) : I.E. increases with the increase in nuclear charge between outermost electrons and nucleus. (Generally, Along the period I.E. increases)

I.E. $\uparrow$ when Effective nuclear charge $\uparrow$

(C) Screening effect : Higher is the screening effect on the outer most electrons causes less attraction from the nucleus and can be easily removed, which is leading to the lower value of I.E.

I.E. $\uparrow$ when Screening effect $\downarrow$

(D) Penetration power of sub shells : (Subshells of the same shell)

(a) Order of attraction of subshells towards nucleus. (Penetration power) is - $s > p > d > f$

(b) As s-subshell is more closer to nucleus so more energy will be required to remove e^- in comparison to p, d & f.

(c) Order of decreasing IE is $s > p > d > f$

Example IE_1 of Be $>$ IE_1 of B
 $(1s^2 2s^2)$ $(1s^2 2s^2 2p^1)$

After loosing one e^- , B attains electronic configuration of Be, so 2nd IE of B is more than Be.

IE_2 of B $>$ Be

(E) Stability of half filled and fully filled orbitals :

(a) Half filled p^3, d^5, f^7 or fully filled s^2, p^6, d^{10}, f^{14} are more stable than others so it requires more energy.
 e.g. In C, N, O, F

IE_1 order is $C < O < N < F$

(b) Because of half filled p-orbitals in N, its ionisation energy (stability) is higher than O.

– IE_1 order $Na < Al < Mg$

(c) Because s-orbital in Mg is completely filled and its penetration power is also higher than p-orbital (Al).

IE_2 order $Mg^+ < Al^+ < Na^+$
 $(2,8,1) \quad (2,8,2) \quad (2,8)$

(F) Oxidation state :

I.E. $\uparrow$ when oxidation state of an atom $\uparrow$

Ion with high oxidation state will have high ionisation energy.

e.g. $Fe^{3+} > Fe^{2+} > Fe$

(iv) Periodicity of IE :

In a group : Size increase so IE decrease.

$Li > Na > K > Rb > Cs$

$Sc > Y > La$



Size increases

IE decreases

Exception :

- IE of Al < I.E. of Ga (While I.E. decreases down the group it is due to Transition contraction)
- IE of Zr < IE of Hf (While I.E. should decrease down the group, it is due to lanthanide contraction)

4d 5d

- $3d > 4d < 5d$ for group 7, 8, 9, 11 and 12

$3d < 4d < 5d$ for group 4, 5, 6 and 10

In a period :- In a period atomic size decreases and Z_{eff} increases so removal of electron becomes difficult and IE increases.

Li Be B C N O F Ne

→
 Z_{eff} increases,
atomic size decreases,
IE increases.

IE = Ne > F > N > O > C > Be > B > Li

I.E. Order in s-block :

IE = H > Be > Mg > Ca > Sr > Li > Ba > Na > K > Rb > Cs

Practice Questions

Q. In which of the following have higher difference in the value of IInd and IIIrd I.E. –

(1) $1s^2 2s^2 2p^6 3s^2 2p^1$

(2) $1s^2 2s^2 2p^6 3s^2$

(3) $1s^2 2s^2 2p^6 3s^1$

(4) $1s^2 2s^2 2p^6 3s^2 3p^2$

Ans. (2)

Q. What is the correct order of I.E. –

(1) $5d > 4d$

(2) $3d < 4d$

(3) $4d \sim 5d$

(4) All

Ans. (1)

Q. Amongst the following elements (whose electronic configurations are given below) the one having highest ionisation energy is :-

(1) $[\text{Ne}]3s^2 3p^1$

(2) $[\text{Ne}]3s^2 3p^3$

(3) $[\text{Ne}]3s^2 3p^2$

(4) $3d^{10}, 4s^2 4p^3$

Ans. (2)

Q. In C, N, O, F which of the following order is correct for I.E. –

(1) $F > O > C > N$

(2) $O > F > N > C$

(3) $F > N > O > C$

(4) $N > F > O > C$

Ans. (3)

Q. Which of the following elements has the lowest ionisation energy ?

(1) Na

(2) K

(3) Mg

(4) Al

Ans. (2)

(v) Application of ionisation energy:**(A) Metallic and non metallic character:**

Metallic nature ↑ when I.E. ↓

Metallic → I.E. Low (Na, K, Rb etc.)

non metallic → I.E. High (F, Cl, Br etc.)

(B) Reactivity of metals:

Reactivity of metals ↑ when I.E. ↓


Reactivity increases down the group as I.E. decreases.

(C) Reducing character :


Reducing nature ↑ when I.E. ↓

(a) IA group has minimum I.E. so they are strong reducing agents and order of reducing nature.

(Li < Na < K < Rb < Cs)



Variation of first ionization energy ($I.E.$)₁ with atomic number for elements with $Z = 1$ to 60



First ionization energy ($I.E.$)₁ of elements of the second period as a function of atomic number (Z)

Periodicity of Ionisation energy in d-block elements:

1st, 2nd, 3rd IE's are increasing from left to right for 1st Transition series, but not regularly.





For 2nd IE $Cr > Fe > Mn$ and $Cu > Zn$

For 3rd IE $Mn > Cr > Fe$ and Zn has highest.

Element	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Atomic number	21	22	23	24	25	26	27	28	29	30
Electronic configuration										
M	$3d^1 4s^2$	$3d^2 4s^2$	$3d^3 4s^2$	$3d^5 4s^1$	$3d^5 4s^2$	$3d^6 4s^2$	$3d^7 4s^2$	$3d^8 4s^1$	$3d^{10} 4s^1$	$3d^{10} 4s^2$
M^{2+}	$3d^1$	$3d^2$	$3d^3$	$3d^4$	$3d^5$	$3d^6$	$3d^7$	$3d^8$	$3d^9$	$3d^{10}$
M^{3+}	[Ar]	$3d^1$	$3d^2$	$3d^3$	$3d^4$	$3d^5$	$3d^6$	$3d^7$	-	-
Enthalpy of atomisation $\Delta_0 H / \text{kJ mol}^{-1}$										
	326	473	515	397	281	416	425	430	339	126
Ionisation Enthalpy, $\Delta_0 H / \text{kJ mol}^{-1}$										
$\Delta_0 H$ I	631	656	650	653	717	762	758	736	745	906
II	1235	1309	1414	1592	1509	161	1644	1752	1958	1734
III	2393	2657	2833	2990	3260	2962	3243	3402	3556	3829





Practice Questions

Q. In which of the following the energy change corresponds to first ionisation energy only :-



Ans. (1)

Q. The correct order of decreasing second ionization energy of Li, Be, Ne, C, B



Ans. (4)

Q. Which of the following statement defines the correct representation of second ionisation enthalpy -

(1) The energy required to remove the electron of a unipositive atom

(2) The energy required to remove the any second electron of an atom

(3) The energy required to remove the second most loosely bound electron

(4) The energy required to remove two electron of an atom.


Ans. (1)

Q. Match the column.

Column-I	Column-II	
Valence electronic configuration	Successive ionisation energies	
(a) ns^1	(p) 19, 27, 36, 48, 270	
(b) ns^2	(q) 16, 28, 34, 260	
(c) $ns^2 np^1$	(r) 18, 26, 230, 250	
(d) $ns^2 np^2$	(s) 14, 200, 220, 240	Ans. (a-s ; b-r ; c-q ; d-p)

Electron Affinity and Electron gain enthalpy:

(i) The amount of energy released when electron is added to the valence shell of an isolated gaseous atom known as electron affinity.




here

E. A. = electron affinity

$\Delta_{eg}H$ = electron gain enthalpy

Note : $\Delta_{eg}H = -E. A.$ (At T = absolute zero)


(ii) Mostly energy is released in the process of adding first e^{-} to neutral atom.




(iii) Generally, $\Delta_{eg_1}H$ is exothermic ($\Delta_{eg_1}H = -ve$)

(iv) $|\Delta_{eg_1}H|$ of neutral atom is equal to magnitude of I.E. of its anion.

{Energy evolved in adding e^{-} to convert neutral atom to anion} = {Energy absorbed to remove e^{-} from anion}



(v) On adding 2nd electron in anion $X^{\ominus}$



$\Delta_{eg_2}H$ (2nd electron gain enthalpy) = +ve

Energy has to be given to introduce 2nd electron in an anion, because of repulsion between negative charge on anion ($X^{\ominus}$) and 2nd electron. So $\Delta_{eg_2}H = +ve$.

(vi) Factors affecting electron affinity :

(a) Atomic size –

E.A. $\uparrow$ when Atomic size $\downarrow$

(b) Screening effect –

E.A. $\uparrow$ when screening effect $\downarrow$

(c) Effective nuclear charge (Z_{eff}) –

E.A. $\uparrow$ when $Z_{eff} \uparrow$



(d) Stability of completely filled or half-filled orbitals –

Electron affinity of elements having fully filled or half-filled configuration is very less or nearly zero and, in some elements, energy is given to add any electron in their valence shell. It is because of stability of fully filled or half-filled configuration.



(vii) In period

- (a) Electron affinity increases along the period due to increase in Z_{eff} and decrease in atomic size.
 (b) Halogens possess maximum electron affinity due to small size and maximum Z_{eff} and after gaining one electron. They attain a stable inert gas configuration.
 (c) Electron affinity of 2nd group, N, and noble gases is zero.
 (d) In 2nd period value of electron affinity of C is higher than N, because of half-filled orbitals present in valence shell of N, which is more stable.
 (e) In 3rd period value of electron affinity of Si is higher than P, because of half-filled orbitals present in P.
 (f) The value of electron affinity of inert gases is zero due to fully filled p-orbitals.



Li	Be	B	C	N	O	F	Ne
Na	Mg	Al	Si	P	S	Cl	Ar
	Zero			Less			zero

**(viii) In Group –**

- (a) Electron affinity decreases down the group. As the size increases and attraction on additional electron is less.
 (b) Electron affinity of 3rd period element is greater than electron affinity of 2nd period elements of the respective group.



Due to small size of fluorine, electron density around the nucleus increases. The incoming electron suffers more repulsion. In case of chlorine electron density decreases due to large size, decreasing order of electron affinity of halogen. $\text{Cl} > \text{F} > \text{Br} > \text{I}$



N & P have low electron affinity due to stable half filled configuration.

(ix) Application of Electron affinity :-

- (a) Oxidising nature $\uparrow$ when E.A. $\uparrow$

But F has more oxidising power than Cl because F has more standard reduction potential.

- (b) Reactivity of non-metal $\uparrow$ when E.A. $\uparrow$

Electron affinity (Data Table) (eV/atom):

1							18
H							He
+0.754		13	14	15	16	17	-0.5
Li		B	C	N	O	F	Ne
+0.618		$\leq +0.277$	+1.263	-0.073	+1.461	+3.399	-1.2
Na		Al	Si	P	S	Cl	Ar
+0.548		+0.441	+1.385	+0.747	+2.077	+3.617	-1.0
K		Ga	Ge	As	Se	Br	Kr
+0.502		+0.03	+1.2	+0.81	+2.021	+3.365	-1.0
Rb		In	Sn	Sb	Te	I	Xe
+0.486		+0.3	+1.2	+1.07	+1.971	+3.059	-0.8

Practice Questions

- Q. Which of the following element has the lowest value of electron affinity -
 (1) Carbon (2) Oxygen (3) Fluorine (4) Neon Ans. (4)
- Q. Which of the following element has the highest value of electron affinity -
 (1) Fluorine (2) Chlorine (3) Sulphur (4) Oxygen Ans. (2)

Electronegativity (EN):

- (i) The tendency of an atom to attract shared pair of electrons towards itself is called electronegativity.
 (ii) EN and EA both have tendency to attract electrons but electron affinity is for isolated atoms. Where as electronegativity is for bonded atoms.
 (iii) A polar covalent bond of A – B may be broken
 as (a) $A - B \longrightarrow A^{\ominus} : + B^{\oplus}$ (EN A > EN B)
 or (b) $A - B \longrightarrow A^{\oplus} + :B^{\ominus}$ (EN A < EN B)
 depending on their tendency to attract bonded electron.
 (iv) There is no unit of electronegativity as EN is tendency to attract electrons of a bonded atom not an energy
 (v) Pauling explained it first time.
 (vi) EN of elements are as follows –

Electronegativity of Some Elements (on Pauling's Scale)

H 2.1						
Li 1.0	Be 1.5	B 2.0	C 2.5	N 3.0	O 3.5	F 4.0
Na 0.9	Mg 1.2	Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0
K 0.8	Ca 1.0	Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8
Rb 0.8	Sr 1.0	In 1.7	Sn 1.8	Sb 1.9	Te 2.1	I 2.5
Cs 0.7	Ba 0.9	Tl 1.8	Pb 1.9	Bi 1.9	Po 2.0	At 2.2
Fr 0.7	Ra 0.9					

In **Pauling's Scale**, elements having almost same electronegativity are-

$C \approx S \approx I \approx 2.5$
 $N = Cl = 3.0$
 $P = H = 2.1$
 $Cs = Fr = 0.7$
 $Be = Al = 1.5$

Note :

- Small atoms are normally having more EN than larger atoms
- Electronegativity of F > Cl but Electron affinity of Cl > F

(vii) Factors Affecting electronegativity:

(A) Atomic size :

E.N. $\uparrow$ when Atomic size $\downarrow$

(B) Effective nuclear charge (Z_{eff}) :-

E.N. $\uparrow$ when Z_{eff} $\uparrow$

(C) Oxidation state :

E.N. $\uparrow$ when oxidation state $\uparrow$

$Mn^{2\oplus} < Mn^{4\oplus} < Mn^{7\oplus}$

$O^{\ominus} < O < O^{\oplus}$

$Fe < Fe^{2\oplus} < Fe^{3\oplus}$

- As atomic radius decreases by increasing oxidation state of cation species, EN increases.
- In anionic species, the order of electronegativity is $O^{2\ominus} < O^{\ominus} < O$

(D) Electronegativity does not depend on filled or half-filled orbitals, because it is a tendency to attract bonded electron, not to gain electron from outside.

(viii) Periodic table & Electronegativity:

- (a) Electronegativity decreases down the group.
 (b) In period on moving from left to right EN increases.

Exceptions:

- '0' group – Electronegativity of '0' group is always zero, because inert gas do not form molecule.
- In IIIA group, after Al value of electronegativity increases down the group, because of scandide contraction and lanthanide contraction.
 EN of Ga > EN of Al

(ix) Application of electronegativity :

(A) Metallic and non metallic nature :-

Low electronegativity $\longrightarrow$ Metals


High electronegativity $\longrightarrow$ Non metals

Metallic character increases down the group but decreases along a period.

(B) Bond length :-

Bond length $\downarrow$ when $\Delta E.N. \uparrow$

Here ΔEN = difference in electronegativities of bonded atoms



- HF has minimum bond length because of much difference in the electronegativities of H and F.

Effect of E.N. on bond length (Schomaker and Stevenson law):

If in a diatomic molecule electronegativities of A – B have more difference. Then actual bond length will be reduced.

As per Schomaker & Stevenson- The reduction in bond length depends on the difference in electronegativities of atoms by following manner -

$$d_{A-B} = r_A + r_B - 0.09 |x_A - x_B|$$

Here x_A is E.N. of A & x_B is E.N. of B

e.g. - If bond length of $F_2 = 1.44 \text{ \AA}$, Bond length of $H_2 = 0.74 \text{ \AA}$.

Find out the bond length of H – F ? (EN of F is 4.0, EN of H is 2.1)


Solution - $d_{H-F} = r_F + r_H - 0.09 |X_F - X_H|$

$$\therefore r_F = 1.44 / 2 = 0.72 \text{ \AA}, r_H = 0.72 / 2 = 0.37 \text{ \AA}$$

$$\therefore d_{H-F} = 0.72 + 0.37 - 0.09 (4.0 - 2.1) \\ = 1.09 - (0.09 \times 1.9) = 1.09 - 0.171 = 0.919 \text{ \AA}$$

(C) Bond energy :- By increasing ΔEN bond length decreases and hence bond energy increases

Bond energy $\propto$ Electronegativity difference



(D) Acidic strength of hydra-acids :

Bond energy (Strength) $\propto$ stability of molecule.

- Order of stability of hydra acids is
 $HF > HCl > HBr > HI$ and so order of their acidic strength will be –
 $HF < HCl < HBr < HI$

- In VA group -

NH ₃	↓	* Thermal stability decreases
PH ₃		* Basic character decreases
AsH ₃		* Acidic character increases

In PH₃ and AsH₃ there is less difference in the electronegativities of X_A and X_B, so their bond energy decreases and hence acidic character (loosing H⁺ ion) increases.

(E) Nature of bonds -

It can be determined by Hanny & Smith formula -

$$\text{Ionic \%} = 16 |X_A - X_B| + 3.5 (X_A - X_B)^2$$

Here X_A = Electronegativity of A

X_B = Electronegativity of B

If X_A - X_B > 2.1 Ionic % > 50% i.e. Ionic bond

If X_A - X_B < 2.1 Ionic % < 50% i.e. covalent bond

- If X_A = X_B; then A - B will be non-polar,

e.g. H—H, F—F

- If X_A > X_B and difference of EN is small then A^{δ-} — B^{δ+} bond will be polar covalent

e.g. H₂O (H^{δ+} — O^{δ-} — H^{δ+})

- If X_A >> X_B and X_A - X_B ΔEN is high then A⁻ — B⁺ bond will be polar ionic

e.g. Na⁺Cl⁻

(F) Nature of oxides and hydroxides:

(a) Along a period, acidic nature of oxides and hydroxides increases.

(b) Down the group basic nature of oxides and hydroxides increases

(c) Types of oxides :

- Generally, oxides of non-metals are acidic in nature. E.g. CO₂, N₂O₃, B₂O₃ etc.

- Generally, oxides of metals are basic in nature. E.g. CaO, MgO, Na₂O etc.

- Oxides of some element are amphoteric in nature





amphoteric oxides are ZnO, Al₂O₃, Ga₂O₃, BeO, PbO, PbO₂, SnO, SnO₂, Sb₂O₃, As₂O₃, V₂O₅, MnO₂, PtO₂ etc.

- CO, H₂O, NO, N₂O are neutral oxides.

(d) Acidic strength of oxide and oxyacid







(i) When central atom is diff.

EN of C.A. ↑, Acidic nature of oxide and oxyacid ↑



(ii) When central atom is same




Oxidation state of C.A. $\uparrow$, EN of C.A. $\uparrow$, Acidic nature of oxide and oxyacid $\uparrow$



(G) Nature of hydroxides :-


(a) If $X_A - X_O > X_O - X_H$

then AO bond will be more polar and will break up as



So, hydroxide is basic

In ClOH -



So, hydroxide is acidic

(H) Nomenclature of inorganic compounds :-

Prefix — less electronegative element

Suffix — More electronegative element

e.g. Cl_2O (Right) OCl_2 (Wrong)

In Dichloroxide the EN of Cl is less than 'O' That's why chloro for Cl is in prefix position.

OF_2 Oxygen difluoride

ICl Iodine chloride

NH_3 Exception (Here H is less electronegative but it is suffix)

(x) Electronegativity Scale :-

(A) Pauling Scale (Bond energy scale) :-

$$0.208\sqrt{\Delta_{AB}} = X_A - X_B, \text{ When B.E. in Kcal/mol}$$

Here Δ_{AB} is resonance energy of AB molecule ; $\Delta_{AB} = \text{BE}_{A-B} - \sqrt{\text{BE}_{A-A} \times \text{BE}_{B-B}}$

BE $\rightarrow$ Bond Energy.

This equation gives difference in electronegativity values.

(B) Mulliken scale :- According to Mulliken electronegativity is average value of IE and EA of an element,

$$X_m = \frac{\text{IE} + \text{EA}}{2}$$

Relation between Pauling scale & Mulliken scale

$$X_p = \text{Electronegativity given by Pauling } X_p = \frac{X_m}{2.8}$$

X_m = Electronegativity given by mulliken

- If IE and EA are given in eV, then electronegativity by Mulliken on Pauling scale will be

$$\text{Pauling EN of an atom} = \frac{\text{IE} + \text{EA}}{5.6}$$

- If IE and EA are given in kcal/mole then

$$\text{EN of an atom} = \frac{\text{IE} + \text{EA}}{2 \times 62.5}$$

(C) Allred Rochow's scale :- As per Rochow, electronegativity is the force by which nucleus of an atom attracts electron which are on the covalent radius.

$$(X_{\text{AR}}) \text{ Electronegativity} = \frac{0.359 Z_{\text{eff}}}{r_{\text{covalent}}^2}$$

$$\text{Relation between } X_{\text{P}} \text{ \& } X_{\text{AR}} \text{ or } X_{\text{P}} = X_{\text{AR}} + 0.744$$

PRACTICE QUESTIONSS

Q. The electronegativities of F and H are 4.0 and 2.1 respectively. The percent ionic character in H and F bond is

- (1) 43 (2) 34 (3) 94 (4) 39 Ans. (1)

Q. Which will be the correct order of electronegativity –

- (1) I < Br < F < Cl (2) F > Cl > Br > I (3) I > Br > F > Cl (4) F < Cl < Br < I Ans. (2)

Properties Dependent on Electronegativity:

- | | |
|--------------------------------------|------------------------|
| (1) % ionic character | (2) Strength of bond |
| (3) Bond Length | (4) Nature of hydrides |
| (5) Nature of oxides and hydroxides. | |

Difference between EN and EA

Electronegativity	Electron Affinity
- Tendency of an atom in a molecule to attract the bonded electrons	- Energy released when an electron is added to neutral isolated gaseous atom
- Relative value of an atom	- Absolute value of an atom
- It regularly changes in a period or group	- It does not change regularly
- It has no unit	- It is measured in eV/atom or kJ mol ⁻¹ or kCal mole ⁻¹