

04

The d and f-Block Elements

The d-Block Elements

The d-block contains the elements of groups 3-12 of the periodic table which are placed in the periods 4, 5, 6 and 7.

Electronic configuration :

General Electronic configuration of d-block elements : $(n-1)d^{1-10} ns^{1-2}$ [except for Pd]

There are mainly four series of the d-block elements, 3d series (Sc to Zn), 4d series (Y to Cd), 5d series (La and Hf to Hg) and 6d series (Ac and Rf to Cn).

1 st Series (3d-series)										
Z	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
	21	22	23	24	25	26	27	28	29	30
4s	2	2	2	1	2	2	2	2	1	2
3d	1	2	3	5	5	6	7	8	10	10
2 nd Series (4d-series)										
Z	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd
	39	40	41	42	43	44	45	46	47	48
5s	2	2	1	1	1	1	1	0	1	2
4d	1	2	4	5	6	7	8	10	10	10
3 rd Series (5d-series)										
Z	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg
	57	72	73	74	75	76	77	78	79	80
6s	2	2	2	2	2	2	2	1	1	2
5d	1	2	3	4	5	6	7	9	10	10
4 th Series (6d-series)										
Z	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Uub
	89	104	105	106	107	108	109	110	111	112
7s	2	2	2	2	2	2	2	2	1	2
6d	1	2	3	4	5	6	7	8	10	10

Transition Elements :

The name transition metals are often used to refer to the elements of the d-block.

As per old convention, originally the name 'Transition Metals' was given because their chemical properties were transitional between s and p-block elements and due to that they are placed between s-block and p-block elements in the periodic table.

As per new convention of IUPAC, the d-block elements which has partially filled d-subshell in their ground state as well as in their common oxidation state, are called 'Transition Metals'.

Since Zn group has d^{10} configuration in their ground state as well as in +2 oxidation state, they are not considered as transition elements but they are d-block elements.

General Characteristics :

(i) **Physical state** : All the d-block elements exist as solid. [except Hg : liquid at room temperature]

(ii) **Metallic character** : They are all metal and good conductor of heat & electricity.

(iii) **Lattice Structures of Transition Metals**


Lattice Structures of Transition Metals

Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
hcp (bcc)	hcp (bcc)	bcc	bcc	X (bcc, ccp)	bcc (hcp)	ccp (hcp)	ccp	ccp	X (hcp)
Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd
hcp (bcc)	hcp (bcc)	bcc	bcc	hcp	hcp	ccp	ccp	ccp	X (hcp)
La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg
hcp (ccp, bcc)	hcp (bcc)	bcc	bcc	hcp	hcp	ccp	ccp	ccp	X

(bcc = body centred cubic; hcp = hexagonal close packed; ccp = cubic close packed; X = a typical metal structure)


(iv) **Variation in atomic radius :**

In a period from left to right in 3d-series atomic size decreases along the period with some irregularities.



Element		Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Metallic Radii (in pm)	M	164	147	135	129	137	126	125	125	128	137
		Decreases from Sc to Cr				Exceptionally larger size	Remains almost same for Fe, Co, Ni			Increases again in Cu and Zn	
Ionic Radii (in pm)	M ²⁺	-	-	79	82	82	77	74	70	73	75
	M ³⁺	73	67	64	62	65	65	61	60	-	-

In a group from 3d to 4d series elements atomic radius increases regularly but from 4d to 5d series element (after La) atomic radius does not increase significantly due to “Lanthanide Contraction”.



Trends in atomic radii of transition elements

Note:

- (i) Atomic Radii order (Group 4 to 12) : $3d < 4d \leq 5d$
 (ii) Atomic Radii regularly increases in Group 3 from Sc to La. ($Sc < Y < La$)

Lanthanide Contraction :

In 5d series elements (after La) 4f subshell is completely filled due to Lanthanoid series elements. Due to poorest shielding of 4f electrons over 5d and 6s electrons and significant increase in nuclear charge from 4d to 5d series, atomic radius of 5d series elements does not increase significantly and contraction in size is observed. This contraction in size is known as "Lanthanide Contraction" or "f-contraction".

Actinide Contraction :

Same as Lanthanoid contraction, Actinide Contraction is observed in 6d series.

Note - Actinide contraction is more effective as compared to Lanthanide contraction.

(v) Density :

- (a) The atomic volume of the transition elements is lower as compared to that of the s-block elements, so their density is comparatively high ($d = M/V$)
 (b) **In a period (3d-series)** density data is as follows :

Element	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Density/g cm ⁻³	3.43	4.1	6.07	7.19	7.21	7.8	8.7	8.9	8.9	7.1

- (c) **In all the groups** (except 3rd Group) generally, there is normal increase in density from 3d to 4d series, but from 4d to 5d density increases nearly twice due to Lanthanide Contraction.

Ex. : $Ti < Zr << Hf$

Note:

- (i) In 3d-series Cu & Ni (as per NCERT) has the highest density and Sc has the lowest density.
 (ii) Os (22.57 gm cm⁻³) and Ir (22.61 gm cm⁻³) have highest density among all the d-block elements.

Metallic Bond Strength :

Generally, strength of metallic bond depends directly on no. of unpaired electrons present in the metal atom thus more will be the unpaired electrons more will be the interaction between atoms and more will be the metallic bond strength.

$$\text{Strength of metallic bond} \propto \text{No. of unpaired electrons}$$

Note:


- (i) Cr, Mo, W are hardest in their series – strong metallic bond (highest no. of unpaired electrons)
 (ii) Zn, Cd, Hg(I) are softest in their series – weak metallic bond (least no. of unpaired electrons)
 (iii) Generally, in a Group metallic bond strength increases [except Zn, Cd, Hg(I)] that's why W is the hardest among all d-block metals.

Melting Point :

Generally, the d-block elements show comparatively high melting points.

Generally, melting point of a metal depends directly on metallic bond strength (no. of unpaired electrons)

$$\text{Melting Point} \propto \text{Metallic Bond Strength (no. of unpaired electrons)}$$



Trends in melting points of transition elements

Note:

- (i) Group 6 elements have highest M.P. in their respective periods (due to max. no. of unpaired electrons) and Group 12 elements have lowest M.P. in their respective periods (due to unavailability of unpaired electrons).

Cr } Highest M.P. in their period
 Mo } due to max. no. of unpaired
 W } electrons are involved in
 } metallic bonding

Zn } Lowest M.P. in their period
 Cd } due to unavailability of unpaired
 Hg } electrons weak metallic bonding
 } is shown


- (ii) Among all d-block elements W has the highest melting point while Hg has the lowest.

- (iii) In Mn and Tc melting point decreases due to their exceptional crystal structure.

Enthalpy of Atomization ($\Delta_a H^\circ$):

In chemistry, the enthalpy of atomisation is the energy required for total separation of all the atoms to convert them into isolated gaseous atoms in a chemical substance.


For metals it is the energy required to separate all metal atoms present in the metal lattice by breaking all the metallic bonds and converting them into isolated gaseous atoms.



Metal lattice $[M(s)]$

Generally, Enthalpy of atomization depends on strength of metallic bonding (no. of unpaired electrons), melting point and boiling point of the metal.

Element	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Enthalpy of atomisation, $\Delta_a H^\ominus$ (in kJ mol^{-1})	326	473	515	397	281	416	425	430	339	126



Trends in enthalpies of atomization of transition elements

Note:

- (i) In 3d series : V has the highest Enthalpy of atomization and Zn has the lowest.
- (ii) Among all d-block elements, W has the highest Enthalpy of atomization.

Ionisation Enthalpies ($\Delta_i H$) :

There is an increase in ionisation enthalpy along each series of the transition elements from left to right due to an increase in nuclear charge but do not increase as regular as in the case of non-transition elements moving from left to right in a period.

Element		Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Ionisation Enthalpy $\Delta_i H$ (kJ mol ⁻¹)	(IE ₁)	631	656	650	653	717	762	758	736	745	906
	(IE ₂)	1235	1309	1414	1592	1509	1561	1644	1752	1958	1734
	(IE ₃)	2393	2657	2833	2990	3260	2962	3243	3402	3556	3829

Note:

- IE₁, IE₂ and IE₃ are increasing from left to right for 3d-series, but not regularly.
- For 2nd IE : Cr > Fe > Mn and Cu > Zn
- For 3rd IE : Mn > Cr > Fe and Zn has the highest 3rd IE.

Variable oxidation states:**Variable oxidation states possible:**

- The elements which give the greatest number of oxidation states occur in or near the middle of the series. Manganese, for example, exhibits all the oxidation states from +2 to +7.
- The lesser number of oxidation states at the extreme ends stems from either too few electrons to lose or share (Sc, Ti) or too many d electrons (hence fewer orbitals available in which to share electrons with others) for higher valence (Cu, Zn).
- Thus, early in the series scandium (II) is virtually unknown and titanium (IV) is more stable than Ti(III) or Ti(II).
- At the other end, the only oxidation state of zinc is +2 (no d electrons are involved).
- The maximum oxidation states of reasonable stability correspond in value to the sum of the s and d electrons upto manganese (Ti^{IV}O₂, V^VO₂⁺, Cr^{VI}O₄²⁻, Mn^{VII}O₄⁻) followed by a rather abrupt decrease in stability of higher oxidation states, so that the typical species to follow are Fe^{II,III}, Co^{II,III}, Ni^{II}, Cu^{I,II}, Zn^{II}.

- (6) The variability of oxidation states, a characteristic of transition elements, arises out of incomplete filling of d orbitals in such a way that their oxidation states differ from each other by unity, e.g., V^{II} , V^{III} , V^{IV} , V^V .
- (7) This is in contrast with the variability of oxidation states of non-transition elements where oxidation states normally differ by a unit of two.
- (8) An interesting feature in the variability of oxidation states of the d-block elements are noticed among the groups (groups 4 through 10).
- (9) In group 6, Mo (VI) and W(VI) are found to be more stable than Cr (VI). Thus Cr (VI) in the form of dichromate in acidic medium is a strong oxidising agent, whereas MoO_3 and WO_3 are not.
- (10) Low oxidation states are found when a complex compound has ligands capable of π -acceptor character in addition to the σ -bonding. For example, in $Ni(CO)_4$ and $Fe(CO)_5$, the oxidation state of nickel and iron is zero.
- (11) As the oxidation number of a metal increases, ionic character decreases. In the case of Mn, Mn_2O_7 is a covalent green oil. Even CrO_3 and V_2O_5 have low melting points. In these higher oxides, the acidic character is predominant. Thus, Mn_2O_7 gives $HMnO_4$ and CrO_3 gives H_2CrO_4 and $H_2Cr_2O_7$. V_2O_5 is, however, amphoteric though mainly acidic and it gives VO_4^{3-} as well as VO_2^+ salts. In vanadium there is gradual change from the basic V_2O_3 to less basic V_2O_4 and to amphoteric V_2O_5 . V_2O_4 dissolves in acids to give VO^{2+} salts. Similarly, V_2O_5 reacts with alkalis as well as acids to give VO_4^{3-} and VO_2^+ respectively. The well characterised CrO is basic but Cr_2O_3 is amphoteric.

Oxidation states of the 1st transition series:

Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
			+1					+1	
	+2	+2	+2	+2	+2	+2	+2	+2	+2
+3	+3	+3	+3	+3	+3	+3	+3		
	+4	+4	+4	+4	+4	+4	+4		
		+5	+5	+5					
			+6	+6	+6				
				+7					

****most common ones are in bold text**

Stability of higher oxidation states :

Trends in stability of higher oxidation state :


- (1) Table shows the stable halides of the 3d series of transition metals. The highest oxidation numbers are achieved in TiX_4 (tetrahalides), VF_5 and CrF_6 . The +7 state for Mn is not represented in simple halides but MnO_3F is known, and beyond Mn no metal has a trihalide except FeX_3 and CoF_3 .
- (2) The ability of fluorine to stabilize the highest oxidation state is due to either higher lattice energy as in the case of CoF_3 , or higher bond enthalpy terms for the higher covalent compounds, e.g., VF_5 and CrF_6 .
- (3) Although V^V is represented only by VF_5 , the other halides, however, undergo hydrolysis to give oxyhalides, VOX_3 .
- (4) Another feature of fluorides is their instability in the low oxidation states e.g., VX_2 ($X = Cl, Br$ or I)

Formulas of halides of 3d-metals

Oxidation Number									
+6			CrF ₆						
+5		VX ₅	CrF ₅						
+4	TiX ₄	VX ₄ ^I	CrX ₄	MnF ₄					
+3	TiX ₃	VX ₃	CrX ₃	MnF ₃	FeX ₃ ¹	CoF ₃			
+2	TiX ₂ ^{III}	VX ₂	CrX ₂	MnX ₂	FeX ₂	CoX ₂	NiX ₂	CuX ₂ ^{III}	ZnX ₂
+1								CuX ^{III}	

Key : X = F → I ; X^I = F → Br ; X^{II} = F → Cl ; X^{III} = Cl → I

and the same applies to CuX. On the other hand, all Cu(II) halides are known except the iodide. In this case, Cu²⁺ oxidises I⁻ to I₂:



(5) However, many copper (I) compounds are unstable in aqueous solution and undergo disproportionation.



(6) The stability of Cu²⁺ (aq) rather than Cu⁺ (aq) is due to the much more negative $\Delta_{\text{hyd}}H^{\circ}$ of Cu²⁺ (aq) than Cu⁺, which more than compensates for the second ionisation enthalpy of Cu.

(7) The ability of oxygen to stabilize the highest oxidation state is demonstrated in the oxides.


(8) The highest oxidation number in the oxides coincides with the group number and is attained in Sc₂O₃ to Mn₂O₇.

(9) Beyond Group 7, no higher oxides of Fe above Fe₂O₃, are known, although ferrates (VI) (FeO₄)²⁻, are formed in alkaline media but they readily decompose to Fe₂O₃ and O₂.

(10) Besides the oxides, oxo cations stabilize V^V as VO₂⁺, V^{IV} as VO²⁺ and Ti^{IV} as TiO²⁺

(11) The ability of oxygen to stabilize these high oxidation states exceeds that of fluorine. Thus, the highest Mn fluoride is MnF₄ whereas the highest oxide is Mn₂O₇. The ability of oxygen to form multiple bonds to metals explains its superiority.

(12) In the covalent oxide Mn₂O₇, each Mn is tetrahedrally surrounded by O's including a Mn–O–Mn bridge.



(13) The tetrahedral [MO₄]ⁿ⁻ ions are known for V^V, Cr^{VI}, Mn^V, Mn^{VI} and Mn^{VII}.


Oxidation Number	Groups									
	3	4	5	6	7	8	9	10	11	12
+7					Mn ₂ O ₇					
+6				CrO ₃						
+5			V ₂ O ₅							
+4		TiO ₂	V ₂ O ₄	CrO ₂	MnO ₂					
+3	Sc ₂ O ₃	Ti ₂ O ₃	V ₂ O ₃	Cr ₂ O ₃	Mn ₂ O ₃	Fe ₂ O ₃				
					Mn ₃ O ₄ [*]	Fe ₃ O ₄ [*]	Co ₃ O ₄ [*]			
+2		TiO	VO	[CrO]	MnO	FeO	CoO	NiO	CuO	ZnO
+1									Cu ₂ O	

* mixed oxides

Standard Electrode Potentials :

Electrode potential : There is no regular trend in these values. This is attributed to the irregular variation of ionization enthalpies ($IE_1 + IE_2$), hydration energies and the sublimation energies in the period.

Element		Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Standard electrode potential $E^\ominus$ (in V)	M^{2+}/M	-	-1.63	-1.18	-0.90	-1.18	-0.44	-0.28	-0.25	+0.34	-0.76
	M^{3+}/M^{2+}	-	-0.37	-0.26	-0.41	+1.57	+0.77	+1.97	-	-	-



Electronic Configurations and some other Properties of the First Series of Transition Elements

Element		Sc	Ti	v	Cr	Mn	Fe	Co	Ni	Cu	Zn
Atomic number (Z)		21	22	23	24	25	26	27	28	29	30
Electronic configuration	M	$3d^1 4s^2$	$3d^2 4s^2$	$3d^3 4s^2$	$3d^5 4s^1$	$3d^5 4s^2$	$3d^6 4s^2$	$3d^7 4s^2$	$3d^8 4s^2$	$3d^{10} 4s^1$	$3d^{10} 4s^2$
	$M^\oplus$	$3d^1 4s^1$	$3d^2 4s^1$	$3d^3 4s^1$	$3d^5$	$3d^5 4s^1$	$3d^6 4s^1$	$3d^7 4s^1$	$3d^8 4s^1$	$3d^{10}$	$3d^{10} 4s^1$
	M^{2+}	$3d^1$	$3d^2$	$3d^3$	$3d^4$	$3d^5$	$3d^6$	$3d^7$	$3d^8$	$3d^9$	$3d^{10}$
	M^{3+}	[Ar]	$3d^1$	$3d^2$	$3d^3$	$3d^4$	$3d^5$	$3d^6$	$3d^7$	-	-
Enthalpy of atomisation, $\Delta_a H^\ominus$ (in kJ mol ⁻¹)		326	473	515	397	281	416	425	430	339	126
Ionisation Enthalpy, $\Delta_i H$ (in kJ mol ⁻¹)	(IE_1)	631	656	650	653	717	762	758	736	745	906
	(IE_2)	1235	1309	1414	1592	1509	1561	1644	1752	1958	1734
	(IE_3)	2393	2657	2833	2990	3260	2962	3243	3402	3556	3829
Metallic Radii (in pm)		164	147	135	129	137	126	125	125	128	137
Ionic radii (in pm)	M^{2+}	-	-	79	82	82	77	74	70	73	75
	M^{3+}	73	67	64	62	65	65	61	60	-	-
Standard Electrode Potential $E^\ominus$ (in V)	M^{2+}/M	-	-1.63	-1.18	-0.90	-1.18	-0.44	-0.28	-0.25	+0.34	-0.76
	M^{3+}/M^{2+}	-	-0.37	-0.26	-0.41	+1.57	+0.77	+1.97	-	-	-
Density (in g cm ⁻³)		3.43	4.1	6.07	7.19	7.21	7.8	8.7	8.9	8.9	7.1

Magnetic Properties :**Magnetic character**

- Magnetic character is measured in terms of spin only magnetic moment $\mu = \sqrt{n(n+2)}$
here n = no. of unpaired electron.
- Metals, which do not have any unpaired electrons show diamagnetism and which have unpaired electrons show para-magnetism.

Diamagnetism

- Substances which are weakly repelled by an external magnetic field are said to be diamagnetic.

Para-magnetism

- Substances which are weakly attracted towards an external magnetic field are said to be para-magnetic.

Ferromagnetism

- Substances very strongly attracted towards an external magnetic field are said to be ferromagnetic.
- Ferromagnetism is the extreme form of para-magnetism.
- Ferromagnetic materials $\Rightarrow$ Fe, Co, Ni etc.

Colours of Ions in Aqueous Solutions :

When an electron from a lower energy d-orbital is excited to a higher energy d-orbital (in split d-orbitals), the energy of excitation corresponds to the frequency of light absorbed. This frequency generally lies in the visible region. The colour observed corresponds to the complementary colour of the light absorbed.

In aqueous solutions the colours of the ions observed are listed in table below

$\text{Ti}^{4\oplus} \rightarrow$	colourless	$\text{Sc}^{3\oplus} \rightarrow$	colourless
$\text{V}^{4\oplus} \rightarrow$	blue	$\text{Ti}^{3\oplus} \rightarrow$	purple
$\text{V}^{2\oplus} \rightarrow$	violet	$\text{V}^{3\oplus} \rightarrow$	green
$\text{Cr}^{3\oplus} \rightarrow$	violet	$\text{Cr}^{2\oplus} \rightarrow$	blue
$\text{Mn}^{2\oplus} \rightarrow$	pink	$\text{Mn}^{3\oplus} \rightarrow$	violet
$\text{Fe}^{3\oplus} \rightarrow$	yellow	$\text{Fe}^{2\oplus} \rightarrow$	light green
$\text{Ni}^{2\oplus} \rightarrow$	green	$\text{Co}^{2\oplus} \rightarrow$	pink
$\text{Zn}^{2\oplus} \rightarrow$	colourless	$\text{Cu}^{2\oplus} \rightarrow$	blue

Note:

Solutions of ions having d^0 and d^{10} configurations are colourless while d^1 to d^9 may show colour.

Complex Formation :

The transition metals form a large number of complex compounds. This is due to the comparatively smaller sizes of the metal ions, their high ionic charges and the availability of d orbitals for bond formation.

Complex compounds are those in which the metal ion binds a number of anions or neutral molecules giving complex species with characteristic properties.

A few examples are: $[\text{Fe}(\text{CN})_6]^{3-}$, $[\text{Fe}(\text{CN})_6]^{4-}$, $[\text{Cu}(\text{NH}_3)_4]^{2+}$, $[\text{PtCl}_4]^{2-}$ etc.

Catalytic Properties :


- The transition metals and their compounds are known for their catalytic activity. This activity is due to their ability to adopt multiple oxidation states and to form complexes.

For example: Vanadium(V) oxide (in Contact Process), finely divided iron (in Haber's Process), and nickel (in Catalytic Hydrogenation).



- Catalysts at a solid surface involve the formation of bonds between reactant molecules and atoms of the surface of the catalyst (first row transition metals utilise 3d and 4s electrons for bonding).

- (3) This has the effect of increasing the concentration of the reactants at the catalyst surface and also weakening of the bonds in the reacting molecules (the activation energy is lowering).
 (4) Also, because the transition metal ions can change their oxidation states, they become more effective as catalysts.

For example, iron (III) catalyses the reaction between iodide and per sulphate ions.



An explanation of this catalytic action can be given as:



Catalyst

Used in Process

TiCl ₃	→	Used as the Ziegler-Natta catalyst in the production of polythene.
V ₂ O ₅	→	Convert SO ₂ to SO ₃ in the contact process for making H ₂ SO ₄
MnO ₂	→	Used as a catalyst to decompose KClO ₃ to give O ₂
Fe	→	Promoted iron is used in the Haber-Bosch process for making NH ₃
FeCl ₃	→	Used in the production of CCl ₄ from CS ₂ and Cl ₂
PdCl ₂	→	Wacker process for converting C ₂ H ₄ + H ₂ O + PdCl ₂ to CH ₃ CHO + 2HCl + Pd.
Pd	→	Used for hydrogenation (e.g. phenol to cyclohexanone).
Pt/PtO	→	Adams catalyst, used for reductions.
Pt/Rh	→	Formerly used in the Ostwald process for making HNO ₃ to oxidize NH ₃ to NO
Cu	→	Is used in the direct process for manufacture of (CH ₃) ₂ SiCl ₂ used to make silicones.
Cu/V	→	Oxidation of cyclohexanol/cyclohexanone mixture to adipic acid which is used to make nylon-66
CuCl ₂	→	Deacon process of making Cl ₂ from HCl
Ni	→	Raney nickel, numerous reduction processes (e.g. manufacture of hexamethylenediamine, production of H ₂ from NH ₃ , reducing anthraquinone to anthraquinol in the production of H ₂ O ₂)
FeSO ₄ + H ₂ O ₂	→	Used as Fenton's reagent for oxidizing alcohols to aldehydes.

Formation of Interstitial Compounds:

Interstitial compounds are those which are formed when small atoms like H, B, C or N are trapped inside the crystal lattices of metals.

The principal physical and chemical characteristics of these compounds are as follows:

- They have high melting points, higher than those of pure metals.
- They are very hard (some borides approach diamond in hardness).
- They retain metallic conductivity.
- They are chemically inert.

Alloy Formation :

An alloy is a blend of metals prepared by mixing the components. Alloys may be homogeneous solid solutions in which the atoms of one metal are distributed randomly among the atoms of the other. Such alloys are formed by atoms with metallic radii that are within about 15 percent of each other. Because of similar radii and other characteristics of transition metals, alloys are readily formed by these metals. The alloys so formed are hard and have often high melting points.


The best known are ferrous alloys: Chromium, Vanadium, Tungsten, Molybdenum and Manganese are used to produce a variety of steels and stainless steel.

Alloys of transition metals with non-transition metals such as brass (Copper-Zinc) and bronze (Copper-Tin), are also of considerable industrial importance.


Some Important Compounds of Transition Elements and their (Physical and Chemical) properties ($K_2Cr_2O_7$, K_2CrO_4 , $KMnO_4$, K_2MnO_4):

Chromate – Dichromate :

Preparation :




[Lime (CaO) added with Na_2CO_3 which keeps the mass porous so that air has access to all parts and prevents fusion]





Then $Na_2Cr_2O_7$ is crystallised out as $Na_2Cr_2O_7 \cdot 2H_2O$ on evaporation.
(red crystal)

How to get $K_2Cr_2O_7$:





$NaCl$ crystallises out first and filtered off. Then $K_2Cr_2O_7$ crystallised out on cooling





The chromates and dichromates are interconvertible in aqueous solution depending upon pH of the solution. The oxidation state of chromium in chromate and dichromate is the same.




The structures of chromate ion, CrO_4^{2-} and the dichromate ion, $Cr_2O_7^{2-}$ are shown below. The chromate ion is tetrahedral whereas the dichromate ion consists of two tetrahedra sharing one corner with $Cr-O-Cr$ bond angle of 126° . Sodium and potassium dichromate are strong oxidising agents; the sodium salt has a greater solubility in water and is extensively used as an oxidising agent in organic chemistry. Potassium dichromate is used as a primary standard in volumetric analysis. In acidic solution, its oxidising action can be represented as follows:



Thus, acidified potassium dichromate will oxidise iodides to iodine, sulphides to sulphur, tin (II) to tin (IV) and iron (II) salts to iron (III). The half-reactions are noted below:



The full ionic equation may be obtained by adding the half-reaction for potassium dichromate to the half-reaction for the reducing agent, for e.g.,



* Similarities between hexavalent Cr & S-compounds :

(i) SO_3 & $\text{CrO}_3 \rightarrow$ both acidic.

(ii) $\text{S} \rightarrow \text{SO}_4^{2-}, \text{S}_2\text{O}_7^{2-}$, $\text{Cr} \rightarrow \text{CrO}_4^{2-}, \text{Cr}_2\text{O}_7^{2-}$

(iii) CrO_4^{2-} & SO_4^{2-} are isomorphous

(iv) SO_2Cl_2 & $\text{CrO}_2\text{Cl}_2 \xrightarrow{\text{OH}^-} \text{SO}_4^{2-}$ & CrO_4^{2-} respectively.

(v) SO_3Cl^- & $\text{CrO}_3\text{Cl}^- \xrightarrow{\text{OH}^-} \text{SO}_4^{2-}$ & CrO_4^{2-}

(vi) CrO_3 & $\text{b}(\text{SO}_3)$ has same structure $\begin{array}{c} \text{O} \\ || \\ -\text{Cr}-\text{O}-\text{Cr}-\text{O}-\text{Cr}- \\ || \quad || \quad || \\ \text{O} \quad \text{O} \quad \text{O} \end{array}$

Illustration 1 :

In laboratory $\text{K}_2\text{Cr}_2\text{O}_7$ is used mainly not $\text{Na}_2\text{Cr}_2\text{O}_7$. Why?

Solution:

$\text{Na}_2\text{Cr}_2\text{O}_7$ is deliquescent enough and changes its concentration and cannot be taken as primary standard solution whereas $\text{K}_2\text{Cr}_2\text{O}_7$ has no water of crystallisation and not deliquescent.


Illustration 2 :

How to standardise $\text{Na}_2\text{S}_2\text{O}_3$ solution in iodometry?

Solution:

$\text{K}_2\text{Cr}_2\text{O}_7$ is primary standard $\Rightarrow$ strength is known by weighing the salt in chemical balance and dissolving in measured amount of water.


Then in acidic solution add. KI




This I_2 is liberated can be estimated with $\text{S}_2\text{O}_3^{2-}$.

Manganate & Permanganate :

Preparation of Manganate (MnO_4^{2-}) and permanganate (MnO_4^-) :



Reaction involved :



Reagents used other than O_2 (air) :





Note- In presence of KClO_3 & KNO_3 the above reaction is faster because these two on decomposition provides O_2 easily.


The above green solution is quite stable in alkali, but in pure water and in presence of acids, depositing MnO_2 and giving a purple solution of permanganate.




Conversion of MnO_4^{2-} into MnO_4^-




But in the above method $\frac{1}{3}$ of Mn is lost as MnO_2 but when oxidised either by Cl_2 or by O_3 , unwanted MnO_2 does not form and no loss of Mn found as MnO_2 .



or




Commercial Method :



Note:

(i) Manganate can be obtained back from permanganate by boiling KMnO_4 with KOH .



(ii) In laboratory conversion of Mn^{2+} to MnO_4^- is done by :

(i) PbO_2

(ii) $\text{Pb}_3\text{O}_4 + \text{HNO}_3$


(iii) $\text{Pb}_2\text{O}_3 + \text{HNO}_3$

(iv) $\text{NaBiO}_3 / \text{H}^+$


(v) $(\text{NH}_4)_2\text{S}_2\text{O}_8 / \text{H}^+$

(vi) $\text{KIO}_4 / \text{H}^+$

Structures of Manganate and Permanganate ions:




Tetrahedral
manganate
(green) ion



Tetrahedral
permanganate
(purple) ion


Oxidising property of KMnO_4 in acidic medium:



- (iii) $\text{MnO}_4^- + \text{H}_2\text{O}_2 + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{O}_2 + \text{H}_2\text{O}$
 (iv) $\text{MnO}_4^- + \text{SO}_2 \xrightarrow{\text{H}^+} \text{Mn}^{2+} + \text{H}_2\text{SO}_4$
 (v) $\text{MnO}_4^- + \text{NO}_2^- + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{NO}_3^- + \text{H}_2\text{O}$
 (vi) $\text{MnO}_4^- + \text{H}_2\text{C}_2\text{O}_4 + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$
 (vii) $\text{MnO}_4^- + \text{H}_2\text{S} \rightarrow \text{Mn}^{2+} + \text{S} \downarrow + \text{H}_2\text{O}$
 (viii) $\text{MnO}_4^- + \text{S}_2\text{O}_3^{2-} \rightarrow \text{Mn}^{2+} + \text{S} \downarrow + \text{SO}_4^{2-}$



Note:

- (i) It is not a primary standard since it is difficult to get it in a high degree of purity and free from traces of MnO_2 .
 (ii) It is slowly reduced to MnO_2 especially in presence of light or acid





Hence it should be kept in dark bottles and standardise just before use.

- (iii) Permanganate titrations in presence of hydrochloric acid are unsatisfactory since hydrochloric acid is oxidised to chlorine.

**Oxidising property of KMnO_4 in neutral or faintly alkaline solution:**


- (i) $2\text{KMnO}_4 + \text{H}_2\text{O} + \text{KI} \rightarrow 2\text{MnO}_2 + 2\text{KOH} + \text{KIO}_3$
 (ii) $2\text{KMnO}_4 + 3\text{KHCO}_2 \rightarrow 2\text{MnO}_2 + \text{KHCO}_3 + 2\text{K}_2\text{CO}_3 + \text{H}_2\text{O}$
 (iii) $2\text{KMnO}_4 + 3\text{H}_2\text{O}_2 \rightarrow 2\text{KOH} + 2\text{MnO}_2 + 2\text{H}_2\text{O} + 3\text{O}_2$
 (iv) $8\text{MnO}_4^- + 3\text{S}_2\text{O}_3^{2-} + \text{H}_2\text{O} \rightarrow 8\text{MnO}_2 + 6\text{SO}_4^{2-} + 2\text{OH}^-$



In absence of Zn^{2+} ions, some of the Mn^{2+} ion may escape, oxidation through the formation of insoluble $\text{Mn}^{\text{II}}[\text{Mn}^{\text{IV}}\text{O}_3]$ manganous permanganite.

Uses of KMnO_4 :

Besides its use in analytical chemistry, potassium permanganate is used as a favourite oxidant in preparative organic chemistry. Its uses for the bleaching of wool, cotton, silk and other textile fibres and for the decolourisation of oils are also dependent on its strong oxidising power.

Heating effect :

The f-Block Elements

Inner Transition Elements :

The elements in which the additional electron enters in $(n - 2)f$ orbitals are called **inner transition elements or f-block elements**.

Position in the periodic table :

The Lanthanides resemble with Yttrium in most of their properties. So, it became necessary to accommodate all the fifteen elements together at one place. This has been done by placing the first element, Lanthanum below Yttrium and placing the remaining fourteen elements separately in the lower part of the periodic table.

Lanthanide series ($Z = 58 - 71$) (Ce – Lu)

Actinide series ($Z = 90 - 103$) (Th – Lr)

Lanthanides (4f-series elements or Rare Earths or Lanthanoids or Lanthanons):

- (i) Lanthanides are reactive elements so do not found in free state in nature.
- (ii) Most important minerals for lighter Lanthanides are – Monazite, Cerites and for heavier Lanthanides – Gadolinite and Xenotime

Electronic configuration :

- (i) The general configuration of Lanthanides may be given as $4f^{1-14}5s^25p^65d^{0-1}6s^2$.

Atomic Number	Element	Symbol	Outer electronic configuration	
			Atomic	+3 ion
58	Cerium	Ce	$4f^1 5d^1 6s^2$	$4f^1$
59	Praseodymium	Pr	$4f^3 6s^2$	$4f^2$
60	Neodymium	Nd	$4f^4 6s^2$	$4f^3$
61	Promethium	Pm	$4f^5 6s^2$	$4f^4$
62	Samarium	Sm	$4f^6 6s^2$	$4f^5$
63	Europium	Eu	$4f^7 6s^2$	$4f^6$
64	Gadolinium	Gd	$4f^7 5d^1 6s^2$	$4f^7$
65	Terbium	Tb	$4f^9 6s^2$	$4f^8$
66	Dysprosium	Dy	$4f^{10} 6s^2$	$4f^9$
67	Holmium	Ho	$4f^{11} 6s^2$	$4f^{10}$
68	Erbium	Er	$4f^{12} 6s^2$	$4f^{11}$
69	Thulium	Tm	$4f^{13} 6s^2$	$4f^{12}$
70	Ytterbium	Yb	$4f^{14} 6s^2$	$4f^{13}$
71	Lutetium	Lu	$4f^{14} 5d^1 6s^2$	$4f^{14}$

- (ii) It is to be noted that filling of 4f orbitals in the atoms is not regular. A 5d electron appears in gadolinium ($Z = 64$) with an outer electronic configuration of $4f^7 5d^1 6s^2$ (and not $4f^8 6s^2$). This is because the 4f and 5d electrons are at about the same potential energy and that the atoms have a tendency to retain stable half-filled configuration.
- (iii) On the other hand, the filling of f-orbitals is regular in tripositive ions.
- (iv) After losing outer electrons, the f-orbitals shrink in size and became more stable.
- (v) **Pm** is the only synthetic radioactive Lanthanide.

Oxidation states :

Lanthanides	Oxidation	Actinides	Oxidation state
^{58}Ce	+3, +4	^{90}Th	+4
^{59}Pr	+3, (+4)	^{91}Pa	(+4), +5
^{60}Nd	+3	^{92}U	(+3), (+4), (+5), +6
^{61}Pm	+3	^{93}Np	(+3), (+4), +5, (+6), (+7)
^{62}Sm	(+2), +3	^{94}Pu	(+3), +4, (+5), (+6), (+7)
^{63}Eu	+2, +3	^{95}Am	+2, (+3), (+4), (+5), (+6)
^{64}Gd	+3	^{96}Cm	+3, (+4)
^{65}Tb	+3, +4	^{97}Bk	+3, (+4)
^{66}Dy	+3, (+4)	^{98}Cf	+3
^{67}Ho	+3	^{99}Es	+3
^{68}Er	(+2), +3	^{100}Fm	+3
^{69}Tm	(+2), +3	^{101}Md	+3
^{70}Yb	+2, +3	^{102}No	+3
^{71}Lu	+3	^{103}Lr	+3

(Oxidation states in brackets are unstable states)

- (i) The Lanthanides contain two s electrons in the outermost shell, they are therefore expected to exhibit a characteristic oxidation state of +2. But for the Lanthanides, the +3 oxidation is common.
- (ii) This corresponds to the use of two outermost electrons ($6s^2$) along with one inner electron. The inner electron used is a 5d electron (in La, Gd and Lu), or one of the 4f electron if no 5d electrons present.
- (iii) All the Lanthanides attain +3 oxidation state and only **Cerium, Praseodymium, and Terbium** exhibit **higher oxidation state (+4)**. **Eu and Yb** exhibit +2 oxidation state.
- (iv) Oxidation states +2 and +4 occur particularly when they lead to –
- (a) A noble gas configuration **Ex.** Ce^{4+} (f^0). The formation of Ce^{IV} is favoured by its noble gas configuration, but it is a strong oxidant reverting to the common +3 state. The E° value for $\text{Ce}^{4+}/\text{Ce}^{3+}$ is +1.74 V which suggests that it can oxidise water. However, the reaction rate is very slow and hence Ce (IV) is a good analytical reagent.
- (b) A half-filled 'f' orbital **Ex.** Eu^{2+} , (f^7), Pr, Nd, Tb and Dy also exhibit +4 state but only in oxides, MO_2 . Eu^{2+} is formed by losing the two s electrons and its f^7 configuration accounts for the formation of this ion. However, Eu^{2+} is a strong reducing agent changing to the common +3 state. Similarly, Yb^{2+} which has f^{14} configuration is a reductant.
- (c) A completely filled 'f' orbital **Ex.** Yb^{2+} (f^{14})
- (v) Therefore, in higher oxidation state, they act as oxidising while in lower state as reducing agents.

Magnetic properties :

- (i) In tri-positive Lanthanide ions the number of unpaired electrons regularly increases from Lanthanum to Gadolinium (0 to 7) and then continuously decreases upto Lutetium (7 to 0).
- (ii) Lanthanum and Lutetium ions are diamagnetic, while all other tri-positive Lanthanide ions are paramagnetic. (Exception – Neodymium is the most paramagnetic Lanthanide).
- (iii) Ce^{4+} and Yb^{2+} are also diamagnetic ions.

Colour :

- (i) The Lanthanide ions have unpaired electrons in their 4f orbitals. Thus, these ions absorb visible region of light and undergo f-f transition and hence exhibit colour.
- (ii) The colour exhibited depends on the number of unpaired electrons in the 4f orbitals.
- (iii) The ions often with $4f^n$ configuration have similar colour to those ions having $4f^{14-n}$ configuration.
- (iv) Lanthanide ions having $4f^0$, $4f^{14}$ are colourless.

Other Properties :

- All the Lanthanides are silvery white soft metals and tarnish rapidly in air.
- The hardness increases with increasing atomic number, samarium being steel hard. Their melting points range between 1000 to 1200 K but samarium melts at 1623 K.
- They have typical metallic structure and are good conductors of heat and electricity.
- Highly dense metals with high melting points do not show any regular trend.
- **Ionisation Energies :** Lanthanides have fairly low ionisation energies comparable to alkaline earth metals.
- **Electro positive character :** High due to low I.P.
- **Complex formation :** Do not have much tendency to form complexes due to low charge density because of their large size. Lu^{3+} is smallest in size can only form complex.
- **Reducing Agent :** They readily lose electrons so are good reducing agent.
- **Alloy :** Alloys of Lanthanides with Fe are called **Misch metals**, which consists of a Lanthanide metal (~ 95%) and iron (~ 5%) and traces of S, C, Ca and Al.
- **Basic Nature :** $\text{La}(\text{OH})_3$ is most basic in nature while $\text{Lu}(\text{OH})_3$ least basic.
- **Carbide :** Lanthanides form MC_2 type carbide with carbon, which on hydrolysis gives C_2H_2 .
- The Lanthanide elements Eu and Yb dissolve directly in very high concentration in liquid ammonia.

Lanthanide contraction :

- (i) In the Lanthanide series with increasing atomic number, in general, there is a progressive decrease in the size from Lanthanum to Lutetium and regular decreases in ionic size from La^{3+} to Lu^{3+} . This decrease in size is known as Lanthanide contraction.
- (ii) The general electronic configuration of these elements is $4f^{1-14} 5d^{0-1} 6s^2$. In these elements the added electron enters the deep-seated f-orbitals.
- (iii) Due to very poor shielding effect of (n-2)f electrons, they exert very little screening effect on the outermost $6s^2$ electrons and therefore experiences considerable pull by the nucleus. Hence, with increasing atomic number, the enhanced nuclear charge leads to contraction in the size of atoms and ions.
- (iv) The atomic volumes of Europium and Ytterbium are unexpectedly large. The large atomic size of Eu and Yb suggest weaker bonding in the solid elements. Both these elements have only two electrons extra than the stable configurations (half filled, f^7 , and completely filled, f^{14}), hence they utilise two electrons in metallic bonding as in the case with Barium.

Effects of Lanthanide Contraction :

Close resemblance of Lanthanides : The general decrease in the sizes of the Lanthanides with an increase in their nuclear charges result in a small increase in their ionisation energies. Hence their basic and ionic nature gradually decreases from La to Lu. This also explains the variations in properties such as increased tendency for hydrolysis and formation of complex salts and decreased thermal stability, solubility of their salts.

Similarity of Yttrium with Lanthanides : The properties of Yttrium are so similar to the Lanthanides that it is considered more a member of the Lanthanide series than a congener of scandium.

Anomalous behaviour of post-Lanthanides : The following anomalies may be observed in the behaviour of post-Lanthanide elements.


(a) Atomic size : The ionic radii of Zr^{4+} is about 9% more than Ti^{4+} . Similar trend is not maintained on passing from the second to third transition series. The ionic radius of Hf^{4+} , instead of increasing (because of inclusion of one more electronic shell), decreases (or is virtually equal to Zr^{4+}) as a consequence of the Lanthanide contraction. This explains the close similarities between the members of the second and third transition series than between the elements of the first and second series.

(b) Ionisation potential and electronegativity : The effect of Lanthanide contraction is also seen in the increase in the ionisation potential values and electronegativities of the elements of the third transition series, contrary to the general trend. Because of the Lanthanide contraction, the post-Lanthanide elements have stronger positive field and thus the electrons are held more tightly.

The greater effective nuclear charge of the former makes them more electronegative than the latter.

(c) High density : Because of Lanthanide contraction, the atomic sizes of the post Lanthanide elements become very small. Consequently, the packing of atoms in their metallic crystals become so much compact that their densities are very high. The densities of the third transition series elements are almost double to those of the second series elements.

(d) Chemical reactions of the Lanthanides :



Application of Lanthanides :

Cerium is most useful element in the Lanthanides –

- Ceramic application – CeO_2 , La_2O_3 , Nd_2O_3 and Pr_2O_3 are used as decolourizing agents for glasses.
- CeS (M.P. – 2000°C) is used in the manufacture of a special type of crucibles and refractories.
- Lanthanide compounds like cerium molybdate, cerium tungstate is used as paints and dyes.
- In textile and leather industries (Ce salts).
- Misch metal is pyrophoric and is used in cigarette & gas lighter.

Actinides (5f-series elements or Actinoids or Actinons) :

- The elements in which the extra electron enters 5f-orbitals of $(n - 2)^{\text{th}}$ main shell are known as Actinides.
- The man-made eleven elements ${}_{93}\text{Np} - {}_{103}\text{Lr}$ are placed beyond uranium in the periodic table and are collectively called trans-uranium elements.
- Th, Pa and U first three Actinides are natural elements.

Electronic configuration :

The general configuration of Actinides may be given as $5f^{1-14} 6d^{0-1} 7s^2$.

Atomic No.	Elements	Symbol	Electronic Configuration
90	Thorium	Th	$5f^0 6d^2 7s^2$
91	Protactinium	Pa	$5f^2 6d^1 7s^2$
92	Uranium	U	$5f^3 6d^1 7s^2$
93	Neptunium	Np	$5f^4 6d^1 7s^2$
94	Plutonium	Pu	$5f^6 6d^0 7s^2$
95	Americium	Am	$5f^7 6d^0 7s^2$
96	Curium	Cm	$5f^7 6d^1 7s^2$
97	Berkelium	Bk	$5f^9 6d^0 7s^2$
98	Californium	Cf	$5f^{10} 6d^0 7s^2$
99	Einsteinium	Es	$5f^{11} 6d^0 7s^2$
100	Fermium	Fm	$5f^{12} 6d^0 7s^2$
101	Mendelevium	Md	$5f^{13} 6d^0 7s^2$
102	Nobelium	No	$5f^{14} 6d^0 7s^2$
103	Lawrencium	Lr	$5f^{14} 6d^1 7s^2$

Oxidation states :

- (i) In Lanthanides and Actinides +3 oxidation is the most common for both of the series of elements.
- (ii) This oxidation state becomes increasingly more stable as the atomic number increases in the Actinide series.
- (iii) Highest oxidation states in the Actinides is +7 exhibited by ${}_{93}\text{Np}$ & ${}_{94}\text{Pu}$, it is unstable.
- (iv) Highest stable oxidation state is +6 shown by ${}_{92}\text{U}$.

Oxidation States of Actinoids

Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
	3	3	3	3	3	3	3	3	3	3	3	3	3
4	4	4	4	4	4	4	4						
	5	5	5	5	5								
		6	6	6	6								
			7	7									

Other Properties :

- **Physical appearance :** Actinides are silvery white metals. They get tarnished when exposed to the attack of alkalis.
- **Density :** All the Actinides except **thorium** and **americium** have high densities.
- **Colour :** Actinide ions are generally coloured. The colour of Actinide ions depends upon the number of 5f-electrons. The ions containing no unpaired 5f-electrons (exactly full filled f-subshell) are colourless, as expected.

- **Ionisation energies** : Ionisation energies values of Actinides are low.
- **Electropositive character** : All the known Actinide metals are **highly electropositive**. They resemble Lanthanide series in this respect.
- **Melting Boiling properties** : They have **high melting and boiling points**. They do not follow regular gradation of melting or boiling points with increase in atomic number.
- **Magnetic properties** : The Actinide elements are paramagnetic due to the presence of unpaired electrons.
- **Radioactive nature** : All the Actinides are radioactive in nature.
- **Actinide contraction** : The size of atom/cation decrease regularly along the Actinides series. The steady decrease in ionic radii with increase in atomic number is referred to as **actinide contraction**. This is due to poor shielding of 5f-electrons.

Comparison of Lanthanides and Actinides :

Points of Resemblance :

- Both Lanthanides and Actinides show a dominant oxidation state of +3.
- Both are electropositive and act as strong reducing agents.
- Cations with unpaired electrons in both of them are paramagnetic.
- Most of the cations of Lanthanides and Actinides are coloured.
- Both of them show a steady decrease in their ionic radii along the series. Thus, Lanthanides show **Lanthanide contraction** and Actinides show **Actinide contraction**.

Difference between Lanthanides & Actinides :

Lanthanides		Actinides	
1.	Besides the most common oxidation state of +3 Lanthanides show +2 and +4 oxidation state in case of certain elements.	1.	Besides the most common oxidation state of +3, actinides show +4, +5 and +6 oxidation states in case of certain elements.
2.	Lanthanides have less tendency towards complex formation.	2.	Actinides have a stronger tendency towards complex formation.
3.	Except promethium, they are non-radioactive.	3.	All the Actinides are radioactive.
4.	Oxides and hydroxide of Lanthanides are less basic.	4.	Oxides and hydroxides of Actinides are more basic.

Some important uses of Actinides are as follows –

Thorium : Thorium is used in atomic reactors as fuel rods and in the treatment of cancer.

Uranium : Uranium is used as nuclear fuel. Its salts are used in glass industry (for imparting green colour). textile industry and also in medicines.

Plutonium : Plutonium is used as fuel for atomic reactors as well as in atomic bombs.