

District Fact Sheet — National Family Health Survey-5 (2019-21)

District: Kurnool

HOUSEHOLD CHARACTERISTICS

- 7. Population living in households with electricity (%) 98.2
- 8. Population living in households with an improved drinking-water source¹ (%) 92.4
- 9. Population living in households that use an improved sanitation facility² (%) 71.6
- 10. Households using clean fuel for cooking³ (%) 61.0
- 11. Households using iodized salt (%) 88.5
- 12. Households with any usual member covered under a health insurance/financing scheme (%) 64.3
- 13. Children age 5 years who attended pre-primary school during the school year 2019-20 (%) 29.7
- 39. Average out-of-pocket expenditure per delivery in a public health facility (Rs.) 3510

NUTRITIONAL STATUS OF CHILDREN

- 73. Children under 5 years who are stunted (height-for-age)¹⁸ (%) 29.4
- 74. Children under 5 years who are wasted (weight-for-height)¹⁸ (%) 17.8
- 75. Children under 5 years who are severely wasted (weight-for-height)¹⁹ (%) 5.9
- 76. Children under 5 years who are underweight (weight-for-age)¹⁸ (%) 31.2

BLOOD SUGAR LEVEL AND BLOOD PRESSURE AMONG ADULTS

Women age 15 years and above:

- 86. Blood sugar level - high (141-160 mg/dl)²³ (%) 5.2
- 87. Blood sugar level - very high (>160 mg/dl)²³ (%) 1.9
- 88. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level²³ (%) 10.5
- 92. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%) 7.6
- 94. Elevated blood pressure (Systolic $\geq$ 140 mm of Hg and/or Diastolic $\geq$ 90 mm of Hg) or taking medicine to control blood pressure (%) 16.1

Men age 15 years and above:

- 89. Blood sugar level - high (141-160 mg/dl)²³ (%) 6.4
- 90. Blood sugar level - very high (>160 mg/dl)²³ (%) 2.6
- 91. Blood sugar level - high or very high (>140 mg/dl) or taking medicine to control blood sugar level²³ (%) 13.9
- 95. Mildly elevated blood pressure (Systolic 140-159 mm of Hg and/or Diastolic 90-99 mm of Hg) (%) 9.1
- 97. Elevated blood pressure (Systolic $\geq$ 140 mm of Hg and/or Diastolic $\geq$ 90 mm of Hg) or taking medicine to control blood pressure (%) 21.1