

06 Oct '26

HW: Listen to previous class videos again
(on Geometry)

HW: Check previous HWs


^a
c

Perimeter of a circle:



If you take a point 'C' on a plane (2D)

and find all the points 'P' that are equidistant
to C, ($\overline{CP} = \text{radius} = r$) those points together form a

circle.



Let's take two circles with radii (plural of radius)
 r_1 and r_2 , and measure their
perimeters P_1 and P_2 .



If $r_1 > r_2$ then $P_1 > P_2$.

Let's measure in each case,

$$\frac{P}{r}, \frac{P_1}{r_1}, \frac{P_2}{r_2} \text{ etc.}$$

You can measure ' r ' using a ruler/tape.
How can you measure ' P '?
(perimeter)

Take a thread and spread it around the circle and mark the length. Then measure the thread's length using a tape.

Hb: Take 20 different radii, draw circles with those radii. Make a table of ' r ', ' P ', $\frac{P}{r}$ in each case.

Eg:

	r	P	$\frac{P}{r}$
(1)	2.3	P_1	$\frac{P_1}{2.3}$
(2)	4	P_2	$\frac{P_2}{4}$
⋮			

(Measure ' P ' in each case using a thread & tape; Compute $\frac{P}{r}$)

Take a picture of all the circles + threads.
Note $r, P, \frac{P}{r}$ on/next to the circles.

To draw a circle of radius ' r ', place your pencil in a compass, fix the distance / compass length to match ' r ' (using a tape / scale), then keep the pointed end of the compass at the center of your circle and draw the circle with the pencil (make sure the compass length doesn't change)

