## CS 105: DIC on Discrete Structures

Graph theory
Basic terminology, Eulerian walks

Lecture 25 Oct 16 2023

# Topic 3: Graph theory

Last topic of this course

Graphs and their properties!

## Topic 3: Graph theory


## Last topic of this course


Graphs and their properties!

### Textbook Reference

- ▶ Introduction to Graph Theory,  $2^{nd}$  Ed., by Douglas West.
- ► Low cost Indian edition available, published by PHI Learning Private Ltd.

).





## Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: e = vu means e is an edge between v and u  $(u \neq v)$ .





### Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: e = vu means e is an edge between v and u  $(u \neq v)$ .

### General Definition

A graph G is a triple V, E, R where V is a set of vertices, E is a set of edges and  $R \subseteq E \times V \times V$  is a relation.





### Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: e = vu means e is an edge between v and u  $(u \neq v)$ .

### General Definition

A graph G is a triple V, E, R where V is a set of vertices, E is a set of edges and  $R \subseteq E \times V \times V$  is a relation.

We will consider only finite graphs (i.e., |V|, |E| are finite) and often simple graphs. Also, we assume |V| > 0.


- ▶ The degree d(v) of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e.,  $|\{vw \in E \mid w \in V\}|$ . A vertex of degree 0 is called an isolated vertex.
- ▶ A walk is a sequence of vertices  $v_1, ..., v_k$  such that  $\forall i \in \{1, ..., k-1\}, (v_i, v_{i+1}) \in E$ . The vertices  $v_1$  and  $v_k$  are called the end-points and others are called internal vertices.

- ▶ The degree d(v) of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e.,  $|\{vw \in E \mid w \in V\}|$ . A vertex of degree 0 is called an isolated vertex.
- A walk is a sequence of vertices  $v_1, \ldots v_k$  such that  $\forall i \in \{1, \ldots k-1\}, (v_i, v_{i+1}) \in E$ . The vertices  $v_1$  and  $v_k$  are called the end-points and others are called internal vertices.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.

- ▶ The degree d(v) of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e.,  $|\{vw \in E \mid w \in V\}|$ . A vertex of degree 0 is called an isolated vertex.
- ▶ A walk is a sequence of vertices  $v_1, ..., v_k$  such that  $\forall i \in \{1, ..., k-1\}, (v_i, v_{i+1}) \in E$ . The vertices  $v_1$  and  $v_k$  are called the end-points and others are called internal vertices.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.
- ▶ A graph is connected if there is a walk between every two vertices.


- ▶ The degree d(v) of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e.,  $|\{vw \in E \mid w \in V\}|$ . A vertex of degree 0 is called an isolated vertex.
- ▶ A walk is a sequence of vertices  $v_1, ..., v_k$  such that  $\forall i \in \{1, ..., k-1\}, (v_i, v_{i+1}) \in E$ . The vertices  $v_1$  and  $v_k$  are called the end-points and others are called internal vertices.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.
- ► A graph is connected if there is a walk between every two vertices.
- ▶ The length of a walk is the number of edges in it.

# Basic terminology: Monday Morning Quiz Part 1



- 1. Give examples for each of the following in the above graph:
  - 1.1 All vertices of degree 3
  - 1.2 A walk of length 5
  - 1.3 A closed walk of length 6

# Basic terminology: Monday Morning Quiz Part 1



- 1. Give examples for each of the following in the above graph:
  - 1.1 All vertices of degree 3
  - 1.2 A walk of length 5
  - 1.3 A closed walk of length 6
- 2. True or False: (a) If a graph is not connected, it must have an isolated vertex.
  - (b) A graph is connected iff some vertex has an edge to every other vertex.

### More definitions

▶ A path is a walk in which no vertex is repeated. Recall, its length is the number of edges in it.

## More definitions


- ▶ A path is a walk in which no vertex is repeated. Recall, its length is the number of edges in it.
- ► A trail is a walk in which no edge is repeated.

### More definitions

- ▶ A path is a walk in which no vertex is repeated. Recall, its length is the number of edges in it.
- ▶ A trail is a walk in which no edge is repeated.
- ▶ A cycle is a (non-empty) closed walk whose internal vertices are all distinct from each other and from the end-point and edges are also distinct.

## More definitions

- A path is a walk in which no vertex is repeated. Recall, its length is the number of edges in it.
- ▶ A trail is a walk in which no edge is repeated.
- ▶ A cycle is a (non-empty) closed walk whose internal vertices are all distinct from each other and from the end-point and edges are also distinct.



- 1. Give examples for each of the following in the above graph:
  - 1.1 A path of length 5
  - 1.2 A cycle of length 6
- 2. True or False: Every path is a walk, every cycle is a closed

## Prove or disprove

If every vertex of a graph G has degree at least 2, then G contains a cycle.

A path P is said to be maximal in G if it is not contained in any longer path. In a finite graph maximal paths exist.

## Prove or disprove

If every vertex of a graph G has degree at least 2, then G contains a cycle.

A path P is said to be maximal in G if it is not contained in any longer path. In a finite graph maximal paths exist.

## Prove or disprove

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Proof:



ightharpoonup Let P be a maximal path in G. It starts from some u.

A path P is said to be maximal in G if it is not contained in any longer path. In a finite graph maximal paths exist.

## Prove or disprove

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ightharpoonup Let P be a maximal path in G. It starts from some u.
- $\triangleright$  Since P cannot be extended, every nbr of u is already in P.

A path P is said to be maximal in G if it is not contained in any longer path. In a finite graph maximal paths exist.

### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Proof:



- ightharpoonup Let P be a maximal path in G. It starts from some u.
- $\triangleright$  Since P cannot be extended, every nbr of u is already in P.
- ▶ As  $d(u) \ge 2$ ,  $\exists v \text{ in } P \text{ such that } uv \notin P$ .
- ightharpoonup Thus we have the cycle  $u \dots vu$ .

A path P is said to be maximal in G if it is not contained in any longer path. In a finite graph maximal paths exist.

### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Proof:



- ightharpoonup Let P be a maximal path in G. It starts from some u.
- $\triangleright$  Since P cannot be extended, every nbr of u is already in P.
- ▶ As  $d(u) \ge 2$ ,  $\exists v \text{ in } P \text{ such that } uv \notin P$ .
- ightharpoonup Thus we have the cycle  $u \dots vu$ .

Is this true if G were infinite?

### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ightharpoonup Let P be a maximal path in G. It starts from some u.
- ightharpoonup Since P cannot be extended, every nbr of u is already in P.
- ▶ As  $d(u) \ge 2$ ,  $\exists v \text{ in } P \text{ such that } uv \notin P$ .
- ightharpoonup Thus we have the cycle  $u \dots vu$ .

Is this true if G were infinite?

No! Consider  $V = \mathbb{Z}$ ,  $E = \{ij : |i - j| = 1\}$ .

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

- ightharpoonup If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

- ightharpoonup If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

- ightharpoonup If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof:

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

- ightharpoonup If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof: ( $\Longrightarrow$ ) Suppose G is Eulerian.

- ► Every vertex has even degree.
  - each passage through a vertex uses two edges (in and out).
  - ▶ at the first vertex first edge is paired with last.

### Definition

A graph is called Eulerian if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an Eulerian walk.

- ightharpoonup If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof: ( $\Longrightarrow$ ) Suppose G is Eulerian.

- ► Every vertex has even degree.
  - each passage through a vertex uses two edges (in and out).
  - ▶ at the first vertex first edge is paired with last.
- Any two edges are in the same walk implies graph is connected (unless it has isolated vertices).

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof ( $\Leftarrow$ ): By induction on number of edges m in G.

ightharpoonup Base case: m=1.

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof ( $\Leftarrow$ ): By induction on number of edges m in G.

▶ Induction step: m > 1. By Lemma G has a cycle C.

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - ightharpoonup Delete all edges in cycle C and (new) isolated vertices.

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - $\triangleright$  Delete all edges in cycle C and (new) isolated vertices.
  - ightharpoonup Get  $G_1, \ldots, G_k$ . Each  $G_i$  is
    - connected
    - ightharpoonup has < m edges.
    - ▶ all its vertices have even degree (why? degree of any vertex was even and removing C, reduces each vertex degree by 0 or 2.)

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - $\triangleright$  Delete all edges in cycle C and (new) isolated vertices.
  - ightharpoonup Get  $G_1, \ldots, G_k$ . Each  $G_i$  is
    - connected
    - ightharpoonup has < m edges.
    - ▶ all its vertices have even degree (why? degree of any vertex was even and removing C, reduces each vertex degree by 0 or 2.)

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - ightharpoonup Delete all edges in cycle C and (new) isolated vertices.
  - $\triangleright$  By indn hyp, each  $G_i$  is Eulerian, has a Eulerian walk.

#### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - ightharpoonup Delete all edges in cycle C and (new) isolated vertices.
  - $\triangleright$  By ind hyp, each  $G_i$  is Eulerian, has a Eulerian walk.
  - Combine the cycle C and the Eulerian walk of  $G_1, \ldots, G_k$  to produce an Eulerian walk of G (how?).

### Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

### Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

- ▶ Induction step: m > 1. By Lemma G has a cycle C.
  - ightharpoonup Delete all edges in cycle C and (new) isolated vertices.
  - $\triangleright$  By ind hyp, each  $G_i$  is Eulerian, has a Eulerian walk.
  - Combine the cycle C and the Eulerian walk of  $G_1, \ldots, G_k$  to produce an Eulerian walk of G (how?).
    - ▶ Traverse along cycle C in G and when some  $G_i$  is entered for first time, detour along an Eulerian walk of  $G_i$ .
    - ▶ This walk ends at vertex where we started detour.
    - ▶ When we complete traversal of *C* in this way, we have completed an Eulerian walk on *G*.