

CS 105: DIC on Discrete Structures

Instructor : S. Akshay

Sept 25, 2023

Lecture 18 – Counting and Combinatorics
Recurrence Relations (Contd.)

Summary and what's next

Part 1: Proofs and basic mathematical structures

Part 2: Counting and Combinatorics

- ▶ Basics of counting
 - ▶ Product principle
 - ▶ Sum principle
 - ▶ Bijection principle
 - ▶ Double counting
- ▶ Subsets, partitions, Permutations and combinations
 1. Binomial coefficients and Binomial theorem
 2. Pascal's triangle
 3. Permutations and combinations with repetitions
 4. Estimating $n!$

Summary and what's next

Part 1: Proofs and basic mathematical structures

Part 2: Counting and Combinatorics

- ▶ Basics of counting
 - ▶ Product principle
 - ▶ Sum principle
 - ▶ Bijection principle
 - ▶ Double counting
- ▶ Subsets, partitions, Permutations and combinations
 1. Binomial coefficients and Binomial theorem
 2. Pascal's triangle
 3. Permutations and combinations with repetitions
 4. Estimating $n!$
- ▶ Recurrence relations and generating functions
- ▶ Pigeonhole Principle and its extensions

Next: Recurrence relations and generating functions

Definition

- ▶ A **recurrence relation** for a sequence is an equation that expresses its n^{th} term using one or more of the previous terms of the sequence.
- ▶ A **linear recurrence relation** is of the form

$$u_n = a_{k-1}u_{n-1} + \dots + a_1u_{n-k+1} + a_0u_{n-k}$$

where $a_0, \dots, a_{k-1} \in \mathbb{R}, k \in \mathbb{N}$ are constants.

Next: Recurrence relations and generating functions

Definition

- ▶ A **recurrence relation** for a sequence is an equation that expresses its n^{th} term using one or more of the previous terms of the sequence.
- ▶ A **linear recurrence relation** is of the form

$$u_n = a_{k-1}u_{n-1} + \dots + a_1u_{n-k+1} + a_0u_{n-k}$$

where $a_0, \dots, a_{k-1} \in \mathbb{R}, k \in \mathbb{N}$ are constants.

- ▶ k is called the **degree/depth** of the sequence.
- ▶ The first few (e.g., k elements u_0, \dots, u_{k-1}) are **initial conditions** and they determine the whole sequence.

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 8?

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 8?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 8?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 8?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$
- ▶ $n = 4 : (((a + b) + c) + d), ((a + b) + (c + d)), ((a + (b + c)) + d), \dots$

More Examples of Recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ How many such bit strings are there of length 2, 3, 4?
- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 8?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$
- ▶ $n = 4 : (((a + b) + c) + d), ((a + b) + (c + d)), ((a + (b + c)) + d), \dots$
In general, let $C(n)$ be the number of ways of doing this.

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
- ▶ k can be anything from 1 till $n - 1$

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
- ▶ k can be anything from 1 till $n - 1$
- ▶ Thus,
$$C(n) = \sum_{i=1}^{n-1} C(i)C(n - i) \text{ for } n > 1$$


Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
- ▶ k can be anything from 1 till $n - 1$
- ▶ Thus,
$$C(n) = \sum_{i=1}^{n-1} C(i)C(n - i) \text{ for } n > 1$$
- ▶ Initial conditions are $C(0) = C(1) = 1$.
- ▶ This sequence are called Catalan numbers...


How do we **solve** such recurrences? We start with the Fibonacci sequence.

An aside: find the Fibonacci sequence!



- ▶ $F(n) = F(n-1) + F(n-2)$.
- ▶ $1, 1, 2, 3, 5, 8, 13, \dots$
- ▶ Can you observe the sum of which terms in the Pascal's triangle gives rise to the terms of the Fibonacci sequence?

An aside: find the Fibonacci sequence!



- ▶ $F(n) = F(n-1) + F(n-2)$.
- ▶ 1, 1, 2, 3, 5, 8, 13,
- ▶ Can you observe the sum of which terms in the Pascal's triangle gives rise to the terms of the Fibonacci sequence?

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation...

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} + F_{n-2}$, $G_n = G_{n-1} + G_{n-2}$ and $H_n = aF_n + bG_n$, then what is a recurrence for H_n ?

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} + F_{n-2}$, $G_n = G_{n-1} + G_{n-2}$ and $H_n = aF_n + bG_n$, then $H_n = H_{n-1} + H_{n-2}$

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} + F_{n-2}$, $G_n = G_{n-1} + G_{n-2}$ and $H_n = aF_n + bG_n$, then $H_n = H_{n-1} + H_{n-2}$

Proof method 1 (for linear recurrences: try α^n !)

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it!

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. How do we get a and b ?

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. Use F_0 and F_1 – initial conditions: $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. Use F_0 and F_1 – initial conditions: $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n.$$

Solving general linear recurrence sequences

- (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:
$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (this can be fixed!) and non-linear recurrences.