CS105 (DIC on Discrete Structures) Problem set 8

- Attempt all questions.
- Apart from things proved in lecture, you cannot assume anything as "obvious". Either quote previously proved results or provide clear justification for each statement.
- 1. Show that if there are 30 students in a class, at least two have last names beginning with the same letter.
- 2. In CS105 this semester there are 186 registered students. They have to be given grades, which must be of course one of following: AA, AB, BB, BC, CC, CD, DD, FR.
 - (a) Can we be sure that at least 10 students have got the same grade? Why or why not?
 - (b) What is the minimum number of students required in the course to be sure that at least 15 will have the same grade?
- 3. Show that among any n+1 positive integers not exceeding 2n there must be an integer that divides one of the other integers.
- 4. Consider the number of strings x_n of length n over the set $\{0, 1, 2\}$ such that any two consecutive numbers in the string differ by at most 1. Thus 01221 is such a string of length 5, but 01200 is not.
 - (a) Write a recurrence relation for x_n .
 - (b) What are x_1 , x_2 and x_7 ?
 - (c) Solve the recurrence to get a closed form solution for x_n in terms of n.
- 5. How many ways can a convex *n*-sided polygon be cut into triangles by adding non-intersecting diagonals (i.e., connecting vertices with non-crossing lines)? Write a recurrence and solve it!
- 6. How many integral solutions does $x_1+x_2+x+3=11$ have where $0 \le x_1 \le 3, 0 \le x_2 \le 4, 0 \le x_3 \le 6$?

- 7. Recall that a derangement is a permutation of objects that leaves no object in its original position. Let D_n denote the number of derangements of $\{1,\ldots,n\}$. Thus, D_n is the number of ways in which letters numbered 1 to n can be put into boxes numbered from 1 to n such that for any i, the letter i is not in box i. We wrote a recurrence for this in the earlier problem sheet. Here is a question on derangements via Principle of Inclusion Exclusion (PIE):
 - (a) Obtain a closed form solution for D_n in terms of n, using PIE.
 - (b) What is $\lim_{n\to\infty} \frac{D_n}{n!}$?
- 8. A bijection $f:\{1,\ldots,2n\}\to\{1,\ldots,2n\}$ is said to be *cool* if there exists $i\in\{1,\ldots,2n-1\}$ such that |f(i)-f(i+1)|=n. Prove that, for each n, more than half the bijections from $\{1,\ldots,2n\}$ to $\{1,\ldots,2n\}$ are cool. (hint: use the Principle of Inclusion and Exclusion!)