

Missing Data - Assignment 1

Aga, Nisse, Ruben

2024-02-28

Contents

1	Introduction (Ruben)	1
2	Methodology (Aga)	2
2.1	Dataset	2
3	Introduction (Ruben)	2
4	Methodology (Aga)	2
4.1	Dataset	2
4.2	Data variables	3
4.3	Data processing methodology	3
5	EDA Results (Nisse)	4
5.1	Descriptive statistics	4
5.2	Correlations	6
5.3	Outliers	7
5.4	Relations	8
6	Missing data problem (Aga)	8
6.1	Missing data and response Patterns	8
6.2	Comparison of the two different models in terms of missing data treatment !!! (Ruben)	14
6.3	Conclusion in terms of answering RQ (Nisse)	14
7	Appendix	14

1 Introduction (Ruben)

1.0.1 RQ (Ruben)

Alcohol consumption led to 2.8 million deaths in 2016 and accounts for almost 10% of global deaths in people aged 15-49 years. Higher levels of alcohol consumption leads to a higher risk of mortality and the only level of

alcohol consumption that minimizes this risk is zero alcohol consumption (study). This means that drinking any alcohol at all increases the risk of mortality. All of this makes it vital to know and understand the factors behind alcohol consumption.

Moore et al. 2005 found that age, sex, ethnicity, marital status, education level and household income were all either negatively or positively associated with alcohol consumption. In Garnett et al. 2022 it was found that the level of depression was negatively associated with alcohol consumption. Considering that the only safe level of alcohol consumption is zero, it is important to understand what factors predict regular consumption of alcohol instead of only looking at the amount of alcohol consumption as in the two studies mentioned above.

The research question of this study is whether the occurrence of regular alcohol consumption (12 or more in a year) can be predicted by the variables: depression level, age, sex, ethnicity (white vs other), marital status and household income. It is expected that age and depression level will be negatively correlated with the occurrence of alcohol consumption while household income will be positively correlated (sources). It is also expected that being male (vs female) and white (vs other ethnicities) will be positively correlated with the regular occurrence of alcohol consumption (source). Regarding marital status it is expected that the married status will be positively associated with the regular occurrence of alcohol consumption compared to others but that is based on a study where the factor marital status consisted of just married and other (source), while in this study more categories are considered.

2 Methodology (Aga)

2.1 Dataset

The dataset used is a subset of the data collected in the National Health and Nutrition Examination Survey (NHANES). The survey is a part of annual program that investigates the health and nutrition of a representative sample of people in the United States. The data we used contains information about 525 individuals that has been collected for the NHANES 2007-2008. This is a subset of the 12,946 individuals in that years' survey sample, out of which 78.4% was interviewed and 75.4% was examined in mobile examination centers. The NHANES is further subdivided into themed sections, such as the Alcohol Use questionnaire, that have separate documentation that will be referred to later.

The used dataset contains a wide range of variables related, to the health of the individuals. We further subset the data by only including variables relevant to the study (demographics, alcohol use and answers to depression screener questions). The selected variables are further described in Variable Description section.

«««< HEAD

3 Introduction (Ruben)

3.0.1 RQ (Ruben)

4 Methodology (Aga)

4.1 Dataset

The dataset used is a subset of the data collected in the National Health and Nutrition Examination Survey (NHANES). The survey is a part of annual program that investigates the health and nutrition of a representative sample of people in the United States. The data we used contains information about 525 individuals that has been collected for the NHANES 2007-2008. This is a subset of the 12,946 individuals in that years' survey sample, out of which 78.4% was interviewed and 75.4% was examined in mobile examination centers.

The used dataset contains a wide range of variables related to the health of the individuals. We further subset the data by only including variables relevant to the study (demographics, alcohol use and answers to depression screener questions). The selected variables are further described in Variable Description section.

4.2 Data variables

Table 1: Variable descriptions

Role	Variable	Name	Type	Characteristics	Target
Outcome	Drink regularly	drink_regularly	Categorical	Binary, yes and no	m/f, age 20-150
Predictor	Sex	sex	Categorical	Binary, male and female	m/f, age 0-150
Predictor	Age	age	Numeric	Discrete	m/f, age 0-150
Predictor	Ethnicity	ethnicity	Categorical	Nominal, 5 categories	m/f, age 0-150
Predictor	Education	marital	Categorical	Nominal, 5 categories	m/f, age 20-150
Predictor	Marital status	marital	Categorical	Nominal, 5 categories	m/f, age 20-150
Predictor	Household income	household_income	Categorical	Nominal, 12 categories	m/f, age 0-150
Predictor	No interest in activity	dep1	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Feeling depressed	dep2	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Sleeping issues	dep3	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Feeling tired	dep4	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Eating issues	dep5	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Feeling bad about yourself	dep6	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Concentrating issues	dep7	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Moving and speaking issues	dep8	Categorical	Ordinal, 1-3 scale	m/f, age 18-150
Predictor	Suicidal thoughts	dep9	Categorical	Ordinal, 1-3 scale	m/f, age 18-150

Table 1 lists the variables used in our subset selection, which will be utilised for the model in question. The predictor variables [*dep1...dep9*] are sourced from the same Depression Screener, where respondents of age 18 to 150 were ought to assign a number (1 to 3) regarding their mental and physical state within the last 2 weeks. Multiple signs of depression were measured this way, which can later be combined to indicate an overall level of depression.

The demographic variables - that being **sex**, **age**, **ethnicity**, **education** and **household_income** - were taken from the same screening component as well. The following should be noted, regarding these demographic variables:

- The variable **age** is topcoded at the value 80 for the respondents who were older than 80 years.
- The variable **education** was targeted at respondents of age 20 to 150, thus excluding younger participants. This is due to the fact that this question includes responses such as **AA degree** and **College Graduate**.
- Similarly, the variable **marital** was also targeted at respondents of age 20 to 150.
- The variable **household_income** is ordinal, rather than continuous.

As for the remaining demographic variables, namely **sex**, **age**, **ethnicity** and **household_income**, these are retrieved from target age 0 to 150.

Finally, the **drink_regularly** variable was obtained from an Alcohol Use questionnaire targeted at ages 20 and up.

4.3 Data processing methodology

After arriving at the subset dataset, as Exploratory Data Analysis was performed. The distributions of the variables were investigated and presence of missing data was found. The important findings of this step are

presented in ‘EDA Results’ section. It is important to note, that after the uneven distribution in ‘marital’ categories was discovered and the relationship of the categories with other variables was investigated, it was decided to combine the not-married categories into one. Thus, a less complex, binary variable was created.

Following the EDA, the missing data problem was addressed. The extend, distribution and patterns of missingness were investigated.

4.3.1 Model methodology

- what we investigated and why

5 EDA Results (Nisse)

5.1 Descriptive statistics

Table 4 from the appendix shows summary statistics for each of the variables within the dataset; including mean values, standard deviations, IQR statistics and data range values. For categorical variables, a list of possible categories and their respective proportions is provided to replace the continuous summary statistics. Lastly, the column N shows the amount of cases with present data - that being non-NA values.

In general, it is worthy to note that all variables are interpreted as a factor, excluding **age** and the multiple depression levels of **dep**. Table 1 suggests that **dep** should in fact be categorical ordinal and should therefore be cast as a factor. That said - and as mentioned in Section 4 - keeping the levels of **dep** as a numerical continuous datatype is beneficial for our missing data problem.

Our dichotomous outcome variable **drink_regularly** has 307 cases of “yes” and 107 cases of “no”, having a outcome balance of 69% and 31% respectively. This outcome ratio could be considered imbalanced, which could affect the accuracy of our logistic regression model. Additionally, the total amount of value entries (N) of 446 suggests that 79 cases contain values outside of the set of possible binary values - most likely being missing values. Table 2 further confirms this.

Table 2: Drink regularly value distributions

drink_regularly	n
yes	307
no	139
NA	79

Predictor **sex** is a dichotomous variable with a balanced frequency distribution across the values “male” and “female”, that being 48% and 52% respectively. 525 entries contain one of these values, suggesting that the variable does not contain missing data.

The predictor **age** is the only continuous variable present within the sub-selected dataset. Although the survey was targeted at respondents of age 0-150 for most variables - the documentation even mentioning the topcoded entries for age 80+ - the dataset seems to only contain cases of people between the age of 20 and 69. Moreover, Figure 1 suggests a uniform distribution of the age variable. Like **sex**, **age** has 525 cases with non-NA values, hence the variable does not contain missing data values.

Predictor **ethnicity** has 5 categories, with **non-hispanic_white** being the most prevalent category with 220 cases (42% of total), all the while **other** is the most infrequent category with 25 cases (5%). A total of 525 cases contain **ethnicity** data, once again suggesting no presence of missing data values within this variable.




Figure 1: Distribution of age

Predictor `education` is similar to `ethnicity`, having 5 categories. That said, the frequencies of said categories seem to be less out of proportion, with `some_college` being the most frequent category with 155 (30%) cases. Whilst the data was retrieved from respondents of ages 20 and higher, we do not observe any missing data values within the variable. This can be further justified by the fact that the minimum `age` within our dataset is 20, therefore foregoing possible issues with younger participants being unable to provide data for this specific variable.

The predictor variable `marital` contains 6 categories. Prior to combining the `marital` categories, the category value `married` exceeded the the average frequency of other categories (that being 49.2) by a large margin - as can be seen in Figure 2. Specifically, 279 cases (53%) were `married`, with the other 5 categories made up the remaining 47% of the cases. `never_married` was the most frequent among those other 5 categories with 102 (19%) cases, whilst `separated` was the most infrequent category with only 14 (3%) cases.




Figure 2: Distribution of marital status categories before pre-processing

5.2 Correlations

```
## Warning in chisq.test(data$marital, data$drink_regularly): Chi-squared
## approximation may be incorrect

##
## Pearson's Chi-squared test
##
## data: data$marital and data$drink_regularly
```

Table 3: Contingency table of marital vs. drink regularly


	yes	no
married	175	81
widowed	8	11
divorced	48	14
separated	8	4
never_married	45	23
living_with_partner	23	6

```
## X-squared = 10.218, df = 5, p-value = 0.06928
```

```
##                X^2 df P(> X^2)
## Likelihood Ratio  9.8952  5 0.078259
## Pearson          10.2183  5 0.069280
##
## Phi-Coefficient   : NA
## Contingency Coeff.: 0.15
## Cramer's V        : 0.151
```

5.3 Outliers

For our only continuous predictor `age`, a boxplot shows potential outliers if datapoints fall outside of the interquartile range (whiskers of the plot). Figure ?? shows the distribution of `age` in said boxplot, revealing no possible outliers. This is to be expected, since `age` was uniformly distributed as mentioned in Section 5.1.



As for our categorical predictors,

Figure 3 shows the frequencies per `marital` category after the pre-processing step of combining the other values into one.

5.4 Relations

6 Missing data problem (Aga)

6.1 Missing data and response Patterns

Firstly, we investigate the overall distribution of missing data in our dataset:






Figure 3: Distribution of marital status categories after pre-processing



As can be seen on the graph above, 8.2% of the data is missing. The missing values occur in the outcome variable 'drink_regularly' and in the responses to questions 'dep2', 'dep3', 'dep5', 'dep6', 'dep8' and 'dep9' that create the depression score variable. 15% of responses are missing for the predictor variable. 25% of the responses are missing for the individual depression questions 2, 3, 5 and 6, 10% are missing for question 8 and 14% for question 9.

We further investigate the missing data patterns by looking at the response patterns:

```
#Creates a graph with all of the response patterns in the dataset and their frequency
md.pattern(data, rotate = TRUE)
```



```
##      id sex age ethnicity education marital household_income dep1 dep4 dep7 dep8
## 263  1  1  1         1         1         1         1  1  1  1  1  1
## 72   1  1  1         1         1         1         1  1  1  1  1  1
## 45   1  1  1         1         1         1         1  1  1  1  1  1
## 22   1  1  1         1         1         1         1  1  1  1  1  1
## 43   1  1  1         1         1         1         1  1  1  1  1  1
## 19   1  1  1         1         1         1         1  1  1  1  1  1
## 7    1  1  1         1         1         1         1  1  1  1  1  1
## 2    1  1  1         1         1         1         1  1  1  1  1  1
## 29   1  1  1         1         1         1         1  1  1  1  0
## 15   1  1  1         1         1         1         1  1  1  1  0
## 2    1  1  1         1         1         1         1  1  1  1  0
## 1    1  1  1         1         1         1         1  1  1  1  0
## 5    1  1  1         1         1         1         1  1  1  1  0
##      0  0  0         0         0         0         0  0  0  0  0  52
##      dep9 drink_regularly dep2 dep3 dep5 dep6
## 263    1                1  1  1  1  1  0
## 72     1                1  0  0  0  0  4
## 45     1                0  1  1  1  1  1
## 22     1                0  0  0  0  0  5
## 43     0                1  1  1  1  1  1
## 19     0                1  0  0  0  0  5
## 7      0                0  1  1  1  1  2
## 2      0                0  0  0  0  0  6
## 29     1                1  1  1  1  1  1
## 15     1                1  0  0  0  0  5
```

```
## 2      1      0      1      1      1      1      2
## 1      1      0      0      0      0      0      6
## 5      0      1      1      1      1      1      2
##      76     79    131    131    131    131    731
```

This figure reveals that there are 13 distinct response patterns in the dataset and the missingness pattern is not monotone. The most frequent pattern is no missing entries, with 263 cases. It is important to note that the depression questions 2, 3, 5 and 6 are always either all present or all missing. It is very probable that the reason for item non-response for the depression items is the same, since there are no cases of only some of them missing.

Based on the missingness pattern of the depression items, 41% of the overall depression score includes at least one missing value.

```
# Creating vectors that indicate if a value is missing in a given variable. Since the pattern in depres

mdrink <- is.na(data$drink_regularly)
mdep <- is.na(data$dep2)

# Testing dependency between missing value in var1 and values of var2. Null hypothesis: no dependency.

out1 <- t.test(age ~ mdrink, data = data)
out1$statistic
```

6.1.0.1 Testing dependency of missing values

```
##          t
## 19.31658
```

```
out1$p.value
```

```
## [1] 3.099076e-45
```

```
# Should this be on data1 or data?
mcar_test(data)
```

```
## # A tibble: 1 x 4
##   statistic    df p.value missing.patterns
##   <dbl> <dbl>   <dbl>         <int>
## 1    465.   164       0             13
```

Thus, the missing values are definitely not missing at random.

- what's the missing data mechanism?

6.1.1 Result models with deletion and imputation (Nisse)

- formula
- table with coefficients and pval (make sure to exponential the coefficients for easier interpretation)
- Interpretation of model result

```
miceOut <- mice(data, defaultMethod = c("norm.predict", "logreg", "polyreg", "polr"), m = 1, maxit = 1)
```

```
##
## iter imp variable
## 1 1 drink_regularly dep2 dep3 dep5 dep6 dep8 dep9
```

```
## Warning: Number of logged events: 1
```

```
reg_imp_data <- complete(miceOut)
summary(reg_imp_data)
```

```
##          id      drink_regularly      sex      age
## Min.   :41531   yes:362          male :254   Min.   :20.00
## 1st Qu.:43912   no :163          female:271 1st Qu.:33.00
## Median :46357
## Mean    :46470
## 3rd Qu.:48934
## Max.    :51610
##
##          ethnicity      education      marital
## mexican_american : 95   no_high_school : 58   Length:525
## other_hispanic    : 61   some_high_school:101   Class :character
## non-hispanic_white:220   high_school_grad:123   Mode  :character
## non-hispanic_black:124   some_college    :155
## other              : 25   college_grad    : 88
##
##          household_income      dep1      dep2      dep3
## 100000+      : 76   Min.    :0.0000   Min.    : -0.2767   Min.    : -0.24575
## 25000:34999: 59   1st Qu.:0.0000   1st Qu.: 0.0000   1st Qu.: 0.00000
## 20000:24999: 52   Median :0.0000   Median : 0.0000   Median : 0.04826
## 35000:44999: 51   Mean    :0.4095   Mean    : 0.3513   Mean    : 0.66590
## 75000:99999: 49   3rd Qu.:1.0000   3rd Qu.: 0.7156   3rd Qu.: 1.00000
## 10000:14999: 45   Max.    :3.0000   Max.    : 3.0000   Max.    : 3.00000
## (Other)      :193
##          dep4      dep5      dep6      dep7
## Min.    :0.0000   Min.    : -0.2370   Min.    : -0.1288   Min.    :0.0000
## 1st Qu.:0.0000   1st Qu.: 0.0000   1st Qu.: 0.0000   1st Qu.:0.0000
## Median :1.0000   Median : 0.0000   Median : 0.0000   Median :0.0000
## Mean    :0.7562   Mean    : 0.3416   Mean    : 0.3206   Mean    :0.3238
## 3rd Qu.:1.0000   3rd Qu.: 0.5244   3rd Qu.: 0.4460   3rd Qu.:0.0000
## Max.    :3.0000   Max.    : 3.0000   Max.    : 3.0161   Max.    :3.0000
##
##          dep8      dep9
## Min.    : -0.2251   Min.    : -0.33496
## 1st Qu.: 0.0000    1st Qu.: 0.00000
```

```
## Median : 0.0000 Median : 0.00000
## Mean   : 0.2042 Mean   : 0.06535
## 3rd Qu.: 0.0000 3rd Qu.: 0.00000
## Max.    : 3.0000 Max.    : 3.00000
##
```

6.2 Comparison of the two different models in terms of missing data treatment !!! (Ruben)

6.3 Conclusion in terms of answering RQ (Nisse)

7 Appendix

Table 4: Summary statistics

Variable	N	Mean	Std. Dev.	Min	Pctl. 25	Pctl. 75	Max
id	525	46470	2898	41531	43912	48934	51610
drink_regularly	446						
... yes	307	69%					
... no	139	31%					
sex	525						
... male	254	48%					
... female	271	52%					
age	525	45	14	20	33	57	69
ethnicity	525						
... mexican_american	95	18%					
... other_hispanic	61	12%					
... non-hispanic_white	220	42%					
... non-hispanic_black	124	24%					
... other	25	5%					
education	525						
... no_high_school	58	11%					
... some_high_school	101	19%					
... high_school_grad	123	23%					
... some_college	155	30%					
... college_grad	88	17%					
marital	525						
... married	279	53%					
... other	246	47%					
household_income	525						
... 0:4999	13	2%					
... 5000:9999	24	5%					
... 10000:14999	45	9%					
... 15000:19999	40	8%					
... 20000:24999	52	10%					
... 25000:34999	59	11%					
... 35000:44999	51	10%					
... 45000:54999	44	8%					
... 55000:64999	35	7%					
... 65000:74999	37	7%					
... 75000:99999	49	9%					
... 100000+	76	14%					
dep1	525	0.41	0.79	0	0	1	3
dep2	394	0.28	0.58	0	0	0	3
dep3	394	0.53	0.86	0	0	1	3
dep4	525	0.76	0.9	0	0	1	3
dep5	394	0.31	0.7	0	0	0	3
dep6	394	0.2	0.56	0	0	0	3
dep7	525	0.32	0.71	0	0	0	3
dep8	473	0.2	0.59	0	0	0	3
dep9	449	0.067	0.37	0	0	0	3