

Exercises in Introductory Physics I

Exercise Sheet 9
due to 18.12.23, 11:59 AM


Rotation

1. The moment of inertia of a sphere (mass M , radius R) as depicted in the figure below (a) is given by:


$$I = \int a^2 \rho(\vec{r}) dV = \int a^2 dm$$

For a sphere with a constant density ρ this can be expressed in spherical coordinates:

$$I = \rho \int_0^R \int_0^\pi \int_0^{2\pi} a^3 r d\varphi d\vartheta dr$$



(a) Moment of inertia of a sphere



(b) Moment of inertia of a sphere with holes (dark grey)

- a) Show that the Moment of Inertia I of an isotropic sphere is (2P)


$$I = \frac{2}{5} MR^2$$

Hint: Show all steps of the calculation. While there are multiple ways to solve this, one might encounter the following integral, for which the solution is given:

$$\int_0^\pi (\sin(\vartheta))^3 d\vartheta = \frac{4}{3}$$


- b) In the figure (b), a sphere (radius R , density ρ) with 4 spherical holes (radius $r = \frac{R}{4}$, the holes are displaced from the centre of the sphere in x - and y -direction) is shown. Calculate the moment of inertia for a rotation around the central y -axis for this object (1P).

2. A wheel of radius r and moment of inertia I about its axis is fixed at the top of an inclined plane (inclination Θ) as shown in the figure below. A string is wrapped around the wheel and its free end supports a block of mass M which can slide on the plane without friction. Given that at $t = 0$ the system is at rest at a height h , how long does it take until it reaches the bottom ?(2P)




3. A thin rod with a length $l_r = 0.4$ m and a mass $m_r = 1$ kg is allowed to turn around its centre. At the beginning it is not moving. Then it is hit at one end by a projectile (mass $m_1 = 0.01$ kg, speed $v_1 = 200$ m \cdot s $^{-1}$) which is going perpendicular to the rod and the rotation axis. The projectile sticks to the rod.

Hint: The mass of the projectile should not be neglected.



- Calculate the angular speed with which the rod starts to move immediately after being hit by the projectile. (1P)
- Calculate the energy that is dissipated during the collision. (1P)

4. A tethered sphere (mass $m_1 = 1 \text{ kg}$) slides frictionless on a horizontal table surface on a circular trajectory of radius r_0 and period $T = 3 \text{ s}$ as depicted in the figure above. The sphere is connected via a massless string to a second sphere (mass $m_2 = 2 \text{ kg}$) through a small hole in the center of the circular trajectory of the first sphere.



- For the equilibrium situation at which $r_0 = \text{const}$, calculate the radius r_0 . (1P)
- Now the mass of the second sphere is increased by 1 kg . What will happen to the velocity of the first sphere? Explain your answer with respect to the acting forces. (1P)
- What are the final radius r_{final} and angular velocity ω_{final} of the sphere on the table when the system reaches equilibrium again? (1P)

Note: Neglect friction. Assume that the second sphere does not reach the ground.