

## Section 6.3: Cardinality

CS1231S Discrete Structures

Wong Tin Lok

National University of Singapore

11 August 2020

No one shall be able to drive us  
from the paradise that Cantor  
created for us.

David Hilbert

Imagine set theory's having been  
invented by a satirist as a kind of  
parody on mathematics.

Ludwig Wittgenstein

---

Tell me your view on infinity at  
<https://pollev.com/wtl>.

## What we saw

- ▶ equality of functions
- ▶ composition of functions
- ▶ injections, surjections, bijections
- ▶ inverses

### Definition 6.2.13

Let  $f: A \rightarrow B$ . Then  $g: B \rightarrow A$  is an *inverse* of  $f$  if

$$\forall x \in A \quad \forall y \in B \quad (y = f(x) \Leftrightarrow x = g(y)).$$

### Theorem 6.2.18

A function is bijective if and only if it has an inverse.

## Now

how one can compare sizes of infinite sets

No one shall be able to drive us from the paradise that Cantor created for us.

David Hilbert

Imagine set theory's having been invented by a satirist as a kind of parody on mathematics.

Ludwig Wittgenstein

---

Tell me your view on infinity at  
<https://pollev.com/wtl>.

## Equality of cardinality

### Definition 5.2.3



- (1) A set is *finite* if it has finitely many (distinct) elements. It is *infinite* if it is not finite.
- (2) Suppose  $A$  is a finite set. The *cardinality* of  $A$ , or the *size* of  $A$ , is the number of (distinct) elements in  $A$ .

### Definition 6.3.1 (Cantor)

- (1) Two sets  $A, B$  are said to have the *same cardinality* if there is a bijection  $A \rightarrow B$ .
- (2) A set is *countable* if it is finite or it has the same cardinality as  $\mathbb{Z}_{\geq 0}$ .

### Note 6.3.2

An infinite set  $B$  is countable if and only if there is a sequence  $b_0, b_1, b_2, \dots \in B$  in which every element of  $B$  appears exactly once.



## Examples of countable sets

### Example 6.3.3

(1)  $\mathbb{Z}_{\geq 0}$  is countable because each element of  $\mathbb{Z}_{\geq 0}$  is listed exactly once in the sequence  
 $0, 1, 2, 3, 4, \dots$

(2) The set  $E = \{2x : x \in \mathbb{Z}_{\geq 0}\}$  is countable because each element of  $E$  is listed exactly once in the sequence

$0, 2, 4, 6, 8, 10, 12, \dots$

only in infinite

Note that  $E \subsetneq \mathbb{Z}_{\geq 0}$ , but  $E$  and  $\mathbb{Z}_{\geq 0}$  have the same cardinality.

(3)  $\mathbb{Z}$  is countable because each element of  $\mathbb{Z}$  is listed exactly once in the sequence  
 $0, 1, -1, 2, -2, 3, -3, \dots$

Note that  $\mathbb{Z}$  is the union of two disjoint infinite sets  $\mathbb{Z}_{\geq 0}$  and  $\mathbb{Z}^-$ , but  $\mathbb{Z}$  and  $\mathbb{Z}_{\geq 0}$  have the same cardinality.

### Note 6.3.2

An infinite set  $B$  is countable if and only if being countable can still be infinite

there is a sequence  $b_0, b_1, b_2, \dots \in B$  in which every element of  $B$  appears exactly once.

## Countable cardinalities are the smallest cardinalities

### Proposition 6.3.4

Any subset  $A$  of a countable set  $B$  is countable.

Proof     **splitting by cases**

1. If  $A$  is finite, then  $A$  is countable by definition.

2. So suppose  $A$  is infinite.

2.1. Then  $B$  is infinite too as  $A \subseteq B$ .

2.2. Use the countability of  $B$  to find a sequence  $b_0, b_1, b_2, \dots$   
in which every element of  $B$  appears exactly once.

2.3. Taking away those terms in the sequence that are not in  $A$ , we are left with a subsequence in which every element of  $A$  appears exactly once.

2.4. So  $A$  is countable. □

either finite

countably infinite

countable

$\vdots$   
 $\bullet$   
 $\bullet$

} infinite

$\vdots$

2 •

1 •

0 •

### Note 6.3.2

An infinite set  $B$  is countable if and only if

there is a sequence  $b_0, b_1, b_2, \dots \in B$  in which every element of  $B$  appears exactly once.





# Uncountability of the power of a countable infinite set

## Theorem 6.3.8 (Cantor 1891)

Let  $A$  be a countable infinite set. Then  $\mathcal{P}(A)$  is not countable.

## Corollary 6.3.9

$\mathbb{R}$  is not countable.

diagonal element	$a_0$	$a_1$	$a_2$	$a_3$	$a_4$	$\dots$
$B_0$	$\notin$	$\in$	$\notin$	$\notin$	$\notin$	$\dots$
$B_1$	$\in$	$\notin$	$\in$	$\notin$	$\in$	$\dots$
$B_2$	$\notin$	$\in$	$\in$	$\notin$	$\in$	$\dots$
$B_3$	$\notin$	$\notin$	$\in$	$\notin$	$\notin$	$\dots$
$B_4$	$\in$	$\notin$	$\in$	$\in$	$\notin$	$\dots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\ddots$
<hr/>						
$B$	$\in$	$\in$	$\notin$	$\in$	$\in$	$\dots$

## Definition 5.2.1

$\mathcal{P}(A)$  denotes the set of all subsets of  $A$ .

- ▶  $B_0, B_1, B_2, \dots \in \mathcal{P}(A)$ .
- ▶  $a_0, a_1, a_2, \dots$  contains every element of  $A$  exactly once.
- ▶  $B \in \mathcal{P}(A)$  but  $B$  does not appear in  $B_0, B_1, B_2, \dots$ .

## Note 6.3.2

An infinite set  $B$  is countable if and only if

there is a sequence  $b_0, b_1, b_2, \dots \in B$  in which every element of  $B$  appears exactly once.





# Summary

## What we saw

- ▶ one way to compare the sizes of infinite sets
- ▶ the cardinality of  $\mathbb{Z}_{\geq 0}$  is the smallest amongst those of infinite sets
- ▶ infinite sets can have different cardinalities

No one shall be able to drive us from the paradise  
that Cantor created for us. David Hilbert



## Questions

- (1) When does one set have a smaller cardinality than another?
- (2) What is the cardinality of an infinite set? Search for “cardinal numbers”.
- (3) Is there a subset of  $\mathcal{P}(\mathbb{Z}_{\geq 0})$  that is not countable but not of the same cardinality as  $\mathcal{P}(\mathbb{Z}_{\geq 0})$ ? Search for “continuum hypothesis”.

## Next

### Induction