available at www.sciencedirect.com
journal homepage: www.europeanurology.com

European Association of Urology

Corrigendum

Corrigendum to "Association of Declining Prostate-specific Antigen Levels with Clinical Outcomes in Patients with Metastatic Castration-resistant Prostate Cancer Receiving [^{177}Lu]Lu-PSMA-617 in the Phase 3 VISION Trial" [Eur. Urol. 86 (2024) 552–562]

Andrew J. Armstrong [a], *Oliver Sartor* [b], *Johann de Bono* [c], *Kim Chi* [d], *Karim Fizazi* [e], *Bernd J. Krause* [f], *Ken Herrmann* [g], *Kambiz Rahbar* [h], *Scott T. Tagawa* [i], *Fred Saad* [j], *Tomasz M. Beer* [k], *Jiwen Wu* [l], *Osvaldo Mirante* [m], *Michael J. Morris* [n,*]

[a] *Duke Cancer Institute Center for Prostate and Urologic Cancers, Departments of Medicine, Surgery, Pharmacology and Cancer Biology, Duke University, Durham, NC, USA;* [b] *Department of Medical Oncology, Mayo Clinic, Rochester, MN, USA;* [c] *The Institute of Cancer Research and The Royal Marsden Hospital, London, UK;* [d] *Division of Medical Oncology, University of British Columbia, Vancouver, BC, Canada;* [e] *Cancer Medicine Department, Gustave Roussy Institute, University of Paris Saclay, Villejuif, France;* [f] *Department of Nuclear Medicine, Rostock University Medical Center, Rostock, Germany;* [g] *Clinic for Nuclear Medicine, University Hospital Essen, Essen, Germany;* [h] *Department of Nuclear Medicine, University Hospital Münster, Münster, Germany;* [i] *Division of Hematology and Medical Oncology, Weill Cornell Medicine, New York, NY, USA;* [j] *Urology Department, University of Montreal Hospital Center, University of Montreal, Montreal, QC, Canada;* [k] *Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA;* [l] *Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA;* [m] *Advanced Accelerator Applications, a Novartis company, Geneva, Switzerland;* [n] *Genitourinary Oncology Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA*

The authors regret a minor data error in the reporting of best radiographic responses in Supplementary Table 5 in the Supplementary Material. The percentage of patients with an evaluable response who had progressive disease (PD) in the 'Increase' subgroup for best overall PSA decline from baseline in the ^{177}Lu-PSMA-617 arm should be 31% not 40%. The number of patients with PD (n = 22) and the denominator (n = 71) remain correct. The data in the Table should therefore read as '22 (31)' not '22 (40)'. The authors also note that the rows in this Table are slightly misaligned vertically but are presented in the correct order.

DOI of original article: https://doi.org/10.1016/j.eururo.2024.08.021
* Corresponding author at: Genitourinary Oncology Service, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY 10065, USA. Tel. +1 646-422-4469.
E-mail address: morrism@mskcc.org (M.J. Morris).

https://doi.org/10.1016/j.eururo.2025.03.008
0302-2838/© 2025 The Author(s). Published by Elsevier B.V. on behalf of European Association of Urology.

Supplementary Table 5 – Efficacy outcomes by best overall PSA decline from baseline in the ^{177}Lu-PSMA-617 arm

	Best overall PSA decline from baseline in the ^{177}Lu-PSMA-617 arm			
	Increase	**≥0–<50% decline**	**≥50–<90% decline**	**≥90% decline**
rPFS[a]				
Events, *n*/*n*	79/96	40/54	69/88	41/106
Median, mo (95% CI)	3.1 (2.4, 3.7)	6.0 (4.2, 7.8)	8.8 (8.5, 10.8)	19.7 (17.0, 20.6)
HR[d] (95% CI)	–	0.41 (0.28, 0.62)	0.20 (0.14, 0.30)	0.04 (0.02, 0.08)
OS[b]				
Events, *n*/*n*	127/146	64/80	75/112	38/157
Median, mo (95% CI)	8.4 (7.3, 9.7)	12.0 (10.3, 14.0)	15.0 (13.9, 17.8)	NE (26.8, NE)
HR[d] (95% CI)	–	0.59 (0.44, 0.80)	0.42 (0.32, 0.56)	0.10 (0.07, 0.15)
Best radiographic response[c]				
Patients with evaluable response, *n*	71	49	78	96
ORR, *n* (%)	1 (1.4)	4 (8.2)	32 (41)	51 (53)
CR, *n* (%)	0 (0)	0 (0)	4 (5.1)	13 (14)
PR, *n* (%)	1 (1.4)	4 (8.2)	28 (36)	38 (40)
SD, *n* (%)	19 (27)	18 (37)	15 (19)	12 (13)
PD, *n* (%)	22 (31)	5 (10)	0 (0)	2 (2.1)
Non-CR/non-PD, *n* (%)	27 (38)	22 (45)	31 (40)	31 (32)

CI = confidence interval; CR = complete response; FAS = full analysis set; HR = hazard ratio; NE = not estimable; ORR = objective response rate (CR + PR); OS = overall survival; PD = progressive disease; PFS-FAS = progression-free survival-full analysis set; PR = partial response; PSA = prostate-specific antigen; PSMA = prostate-specific membrane antigen; rPFS = radiographic progression-free survival; SD = stable disease.

[a] Analysis in PFS-FAS for the ^{177}Lu-PSMA-617 arm only (n = 385).

[b] Analysis in FAS for the ^{177}Lu-PSMA-617 arm only (n = 551).

[c] Analysis in patients with evaluable response (n = 319).

[d] Versus PSA increase.

The authors would like to apologise for any inconvenience caused.