

## Problem Set 1

Student Name: Noah Reef

## Problem 6.1

To find the Fourier transform of  $f(x) = e^{-|x|}$  for  $x \in \mathbb{R}$ , we compute

$$\begin{aligned}
 \hat{f}(\xi) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-|x|} e^{-ix\xi} dx \\
 &= \frac{1}{\sqrt{2\pi}} \left( \int_{-\infty}^0 e^{x(1-i\xi)} dx + \int_0^{\infty} e^{-x(1+i\xi)} dx \right) \\
 &= \frac{1}{\sqrt{2\pi}} \left( \frac{1}{1-i\xi} e^{x(1-i\xi)} \Big|_{-\infty}^0 - \frac{1}{1+i\xi} e^{-x(1+i\xi)} \Big|_0^{\infty} \right) \\
 &= \frac{1}{\sqrt{2\pi}} \left( \frac{2}{(1-i\xi)(1+i\xi)} \right) \\
 &= \frac{1}{\sqrt{2\pi}} \left( \frac{2}{1+\xi^2} \right)
 \end{aligned}$$

## Problem 6.2

We will consider the parameteric curve defined as

$$\begin{aligned}
 \gamma_1 &= t & -R \leq t \leq R \\
 \gamma_2 &= R - it & 0 \leq t \leq \frac{\xi}{2a} \\
 \gamma_3 &= -t - i\frac{\xi}{2a} & -R \leq t \leq R \\
 \gamma_4 &= -R + it & -\frac{\xi}{2a} \leq t \leq 0
 \end{aligned}$$

and by Cauchy's Integral theorem we have that

$$\begin{aligned}
 0 &= \int_{-R}^R e^{-at^2 - i\xi t} dt + i \int_0^{\xi/2a} e^{-a(R-it)^2 - i\xi(R-it)} dt \\
 &\quad - \int_{-R}^R e^{-a(-t-i\xi/2a)^2 - i(-t-i\xi/2a)} dt - i \int_{-\xi/2a}^0 e^{-a(-R+it)^2 - i\xi(-R+it)} dt
 \end{aligned}$$

Then we see that the second and fourth terms go to zero since,

$$\left| i \int_0^{\xi/2a} e^{-a(R-it)^2 - i\xi(R-it)} dt \right| + \left| i \int_{-\xi/2a}^0 e^{-a(-R+it)^2 - i\xi(-R+it)} dt \right| \leq 2 \frac{\xi}{2a} e^{-aR^2} \rightarrow 0$$

and we see for the third term that

$$\lim_{R \rightarrow \infty} \int_{-R}^R e^{-a(-t-i\xi/2a)^2 - i(-t-i\xi/2a)} dt = \sqrt{\frac{\pi}{a}} e^{-\xi^2/(4a)}$$

and hence

$$\hat{f}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-a|x|^2} e^{-ix\xi} dx = \sqrt{\frac{\pi}{a}} \frac{e^{-\xi^2/(4a)}}{\sqrt{2\pi}}$$

## Problem 6.4

Suppose the  $f \in L^1(\mathbb{R}^d)$  and  $f(x) = g(|x|)$  for some  $g$ , then we see that

$$\begin{aligned} \hat{f}(\xi) &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x) e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} g(|x|) e^{-ix \cdot \xi} dx \\ &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} g(|x|) e^{-i|x||\xi| \cos(\theta)} dx \\ &= (2\pi)^{-d/2} \int_0^\infty \int_{\omega_d} g(r) e^{-ir|\xi| \cos(\theta)} r^d dr d\theta \\ &= h(|\xi|) \end{aligned}$$