

## Problem Set 4

*Student Name: Noah Reef***Problem 4.1****Part a**

Recall that the spin operator is given by  $\hat{S} = \frac{1}{2}\hbar\sigma$  where  $\sigma = \sigma_x u_x + \sigma_y u_y + \sigma_z u_z$ . Then we have that the expectation of the spin operator is

$$\langle \Psi | \hat{S} | \Psi \rangle = \frac{\hbar}{2} [\langle \Psi | \sigma_x | \Psi \rangle u_x + \langle \Psi | \sigma_y | \Psi \rangle u_y + \langle \Psi | \sigma_z | \Psi \rangle u_z]$$

and we have that

$$\begin{aligned} \langle \Psi | \sigma_x | \Psi \rangle &= [u^*(r) \quad d^*(r)] \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{bmatrix} u(r) \\ d(r) \end{bmatrix} = u^*(r)d(r) + d^*(r)u(r) \\ &= \operatorname{Re}(u^*d) + i \operatorname{Im}(u^*d) + \operatorname{Re}(u^*d) - i \operatorname{Im}(u^*d) \\ &= 2 \operatorname{Re}(u^*d) \end{aligned}$$

and

$$\begin{aligned} \langle \Psi | \sigma_y | \Psi \rangle &= [u^*(r) \quad d^*(r)] \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{bmatrix} u(r) \\ d(r) \end{bmatrix} = -iu^*(r)d(r) + id^*(r)u(r) \\ &= -i \operatorname{Re}(u^*d) + \operatorname{Im}(u^*d) + i \operatorname{Re}(u^*d) + \operatorname{Im}(u^*d) \\ &= 2 \operatorname{Im}(u^*d) \end{aligned}$$

lastly,

$$\begin{aligned} \langle \Psi | \sigma_z | \Psi \rangle &= [u^*(r) \quad d^*(r)] \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{bmatrix} u(r) \\ d(r) \end{bmatrix} = u^*(r)u(r) - d^*(r)d(r) \\ &= |u|^2 - |d|^2 \end{aligned}$$

and hence then expectation value of the spin operator on this spinsor is given by

$$\langle \Psi | \hat{S} | \Psi \rangle = \frac{\hbar}{2} [2 \operatorname{Re}(u^*d)u_x + 2 \operatorname{Im}(u^*d)u_y + (|u|^2 - |d|^2)u_z]$$

**Part b**

We compute the norm of  $\langle \Psi | \hat{S} | \Psi \rangle$  as

$$\begin{aligned}
 |\langle \Psi | \hat{S} | \Psi \rangle| &= \frac{\hbar}{2} \sqrt{4 \operatorname{Re}(u^* d)^2 + 4 \operatorname{Im}(u^* d)^2 + |u|^4 - 2|u|^2|d|^2 + |d|^4} \\
 &= \frac{\hbar}{2} \sqrt{4|u^* d|^2 + |u|^4 - 2|u|^2|d|^2 + |d|^4} \\
 &= \frac{\hbar}{2} \sqrt{4|u|^2|d|^2 + |u|^4 - 2|u|^2|d|^2 + |d|^4} \\
 &= \frac{\hbar}{2} \sqrt{|u|^4 + 2|u|^2|d|^2 + |d|^4} \\
 &= \frac{\hbar}{2} \sqrt{(|u|^2 + |d|^2)^2} \\
 &= \frac{\hbar}{2} \sqrt{|\Psi|^2} = \frac{\hbar}{2}
 \end{aligned}$$

**Part c**

Consider

$$\exp\left(i\varphi \frac{\hat{S}_z}{\hbar}\right) = \sum_{n=0}^{\infty} \frac{\left(i\varphi \frac{\hat{S}_z}{\hbar}\right)^n}{n!} = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{i\varphi}{2}\right)^n \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^n$$

note that for  $n$  even we have that

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^n = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and remains unchanged for  $n$  odd, thus we can write the above as

$$\exp\left(i\varphi \frac{\hat{S}_z}{\hbar}\right) = \begin{pmatrix} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\frac{i\varphi}{2}\right)^n & 0 \\ 0 & \sum_{n=0}^{\infty} \frac{1}{n!} \left(-\frac{i\varphi}{2}\right)^n \end{pmatrix} = \begin{pmatrix} \exp\left(\frac{i\varphi}{2}\right) & 0 \\ 0 & \exp\left(-\frac{i\varphi}{2}\right) \end{pmatrix}$$

**Part d**

Note that

$$\Psi' = \exp\left(i\varphi \frac{\hat{S}_z}{\hbar}\right) \Psi = \begin{pmatrix} \exp\left(\frac{i\varphi}{2}\right) & 0 \\ 0 & \exp\left(-\frac{i\varphi}{2}\right) \end{pmatrix} \begin{bmatrix} u \\ d \end{bmatrix} = \begin{bmatrix} e^{i\varphi/2} u \\ e^{-i\varphi/2} d \end{bmatrix}$$

then we get that

$$\begin{aligned}
 \langle \Psi | \sigma_x | \Psi \rangle &= \begin{bmatrix} e^{-i\varphi/2} u^*(r) & e^{i\varphi/2} d^*(r) \end{bmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{bmatrix} e^{i\varphi/2} u(r) \\ e^{-i\varphi/2} d(r) \end{bmatrix} = e^{-i\varphi} u^*(r) d(r) + e^{i\varphi} d^*(r) u(r) \\
 &= \operatorname{Re}(e^{-i\varphi} u^* d) + i \operatorname{Im}(e^{-i\varphi} u^* d) + \operatorname{Re}(e^{-i\varphi} u^* d) - i \operatorname{Im}(e^{-i\varphi} u^* d) \\
 &= 2 \operatorname{Re}(e^{-i\varphi} u^* d)
 \end{aligned}$$

and

$$\begin{aligned}\langle \Psi | \sigma_y | \Psi \rangle &= \begin{bmatrix} e^{-i\varphi/2} u^*(r) & e^{i\varphi/2} d^*(r) \end{bmatrix} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \begin{bmatrix} e^{i\varphi/2} u(r) \\ e^{-i\varphi/2} d(r) \end{bmatrix} = -ie^{i\varphi} u^*(r) d(r) + ie^{i\varphi} d^*(r) u(r) \\ &= -i \operatorname{Re}(e^{-i\varphi} u^* d) + \operatorname{Im}(e^{-i\varphi} u^* d) + i \operatorname{Re}(e^{-i\varphi} u^* d) + \operatorname{Im}(e^{-i\varphi} u^* d) \\ &= 2 \operatorname{Im}(e^{-i\varphi} u^* d)\end{aligned}$$

lastly,

$$\begin{aligned}\langle \Psi | \sigma_z | \Psi \rangle &= \begin{bmatrix} e^{-i\varphi/2} u^*(r) & e^{i\varphi/2} d^*(r) \end{bmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{bmatrix} e^{i\varphi/2} u(r) \\ e^{-i\varphi/2} d(r) \end{bmatrix} = u^*(r) u(r) - d^*(r) d(r) \\ &= |u|^2 - |d|^2\end{aligned}$$

and hence then expectation value of the spin operator on this spensor is given by

$$\langle \Psi' | \hat{S} | \Psi' \rangle = \frac{\hbar}{2} [2 \operatorname{Re}(e^{-i\varphi} u^* d) u_x + 2 \operatorname{Im}(e^{-i\varphi} u^* d) u_y + (|u|^2 - |d|^2) u_z]$$

## Part e

Suppose that  $u^* d = a + bi$  then  $a = \frac{1}{2} S_x$  and  $b = \frac{1}{2} S_y$  and we have that

$$e^{-i\varphi} u^* d = (a + bi)(\cos(\varphi) - i \sin(\varphi)) = a \cos(\varphi) + b \sin(\varphi) + i(b \cos(\varphi) - a \sin(\varphi))$$

thus

$$S'_x = 2 \operatorname{Re}(e^{-i\varphi} u^* d) = S_x \cos(\varphi) + S_y \sin(\varphi)$$

and

$$S'_y = 2 \operatorname{Im}(e^{-i\varphi} u^* d) = S_y \cos(\varphi) - S_x \sin(\varphi)$$

and clearly

$$S'_z = S_z$$

## Part f

From the change of coordinates in Part e, we have that  $\exp\left(i\varphi \frac{\hat{S}_z}{\hbar}\right)$  is a clockwise rotation about the  $z$ -axis by angle  $\varphi$ .

## Problem 4.2

### Part a

Recall that

$$\chi_{\uparrow} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \chi_{\downarrow} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

and note that

$$\begin{aligned}\chi_{\uparrow} \otimes \chi_{\uparrow} &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} \otimes \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \\ \chi_{\uparrow} \otimes \chi_{\downarrow} &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} \otimes \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} \\ \chi_{\downarrow} \otimes \chi_{\uparrow} &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} \otimes \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} \\ \chi_{\downarrow} \otimes \chi_{\downarrow} &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} \otimes \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}\end{aligned}$$

and hence we see that

$$\psi(\mathbf{r}_1, \mathbf{r}_2; 1)\chi_{\uparrow} \otimes \chi_{\uparrow} + \psi(\mathbf{r}_1, \mathbf{r}_2; 2)\chi_{\uparrow} \otimes \chi_{\downarrow} + \psi(\mathbf{r}_1, \mathbf{r}_2; 3)\chi_{\downarrow} \otimes \chi_{\uparrow} + \psi(\mathbf{r}_1, \mathbf{r}_2; 4)\chi_{\downarrow} \otimes \chi_{\downarrow}$$

becomes

$$\psi(\mathbf{r}_1, \mathbf{r}_2; 1) \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \psi(\mathbf{r}_1, \mathbf{r}_2; 2) \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \psi(\mathbf{r}_1, \mathbf{r}_2; 3) \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + \psi(\mathbf{r}_1, \mathbf{r}_2; 4) \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \psi(\mathbf{r}_1, \mathbf{r}_2; 1) \\ \psi(\mathbf{r}_1, \mathbf{r}_2; 2) \\ \psi(\mathbf{r}_1, \mathbf{r}_2; 3) \\ \psi(\mathbf{r}_1, \mathbf{r}_2; 4) \end{bmatrix}$$

## Part b

To compute  $(\hat{S}_{\text{tot}})_z$  we compute

$$(\hat{S}_{\text{tot}})_z = S_z \otimes I + I \otimes S_z = \frac{\hbar}{2} (\sigma_z \otimes I + I \otimes \sigma_z)$$

then we have that

$$\begin{aligned}\sigma_z \otimes I &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \\ I \otimes \sigma_z &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}\end{aligned}$$

then we have that

$$(\hat{S}_{\text{tot}})_z = \frac{\hbar}{2} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} = \hbar \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

Note that doing a similar process for  $(S_{\text{tot}})_x$  and  $(S_{\text{tot}})_y$  we get that

$$(\hat{S}_{\text{tot}})_x = \frac{\hbar}{2} \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$(\hat{S}_{\text{tot}})_y = \frac{\hbar}{2} \begin{pmatrix} 0 & -i & -i & 0 \\ i & 0 & 0 & -i \\ i & 0 & 0 & -i \\ 0 & i & i & 0 \end{pmatrix}$$

so then we have that

$$S_{\text{tot}}^2 = (\hat{S}_{\text{tot}})_x^2 + (\hat{S}_{\text{tot}})_y^2 + (\hat{S}_{\text{tot}})_z^2 = \begin{pmatrix} 2\hbar^2 & 0 & 0 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & 0 & 0 & 2\hbar^2 \end{pmatrix}$$

## Part c

Consider

$$\begin{aligned} \Psi_1 &= f(\mathbf{r}_1, \mathbf{r}_2) \chi_{\uparrow} \otimes \chi_{\uparrow} \\ \Psi_2 &= f(\mathbf{r}_1, \mathbf{r}_2) \chi_{\uparrow} \otimes \chi_{\downarrow} \\ \Psi_3 &= f(\mathbf{r}_1, \mathbf{r}_2) \chi_{\downarrow} \otimes \chi_{\uparrow} \\ \Psi_4 &= f(\mathbf{r}_1, \mathbf{r}_2) \chi_{\downarrow} \otimes \chi_{\downarrow} \end{aligned}$$

Then we have that

$$\begin{aligned} \langle \Psi_1 | S_{\text{tot}}^2 | \Psi_1 \rangle &= \int [f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0 \quad 0 \quad 0] \begin{pmatrix} 2\hbar^2 & 0 & 0 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & 0 & 0 & 2\hbar^2 \end{pmatrix} \begin{bmatrix} f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \\ 0 \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\ &= \int 2\hbar^2 |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = 2\hbar^2 \\ \langle \Psi_2 | S_{\text{tot}}^2 | \Psi_2 \rangle &= \int [0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0 \quad 0] \begin{pmatrix} 2\hbar^2 & 0 & 0 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & 0 & 0 & 2\hbar^2 \end{pmatrix} \begin{bmatrix} 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\ &= \int \hbar^2 |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = \hbar^2 \end{aligned}$$

$$\begin{aligned}
\langle \Psi_3 | S_{\text{tot}}^2 | \Psi_3 \rangle &= \int [0 \quad 0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0] \begin{pmatrix} 2\hbar^2 & 0 & 0 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & 0 & 0 & 2\hbar^2 \end{pmatrix} \begin{bmatrix} 0 \\ 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \int \hbar^2 |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = \hbar^2
\end{aligned}$$

$$\begin{aligned}
\langle \Psi_4 | S_{\text{tot}}^2 | \Psi_4 \rangle &= \int [0 \quad 0 \quad 0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2)] \begin{pmatrix} 2\hbar^2 & 0 & 0 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & \hbar^2 & \hbar^2 & 0 \\ 0 & 0 & 0 & 2\hbar^2 \end{pmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \int 2\hbar^2 |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = 2\hbar^2
\end{aligned}$$

### Part d

$$\begin{aligned}
\langle \Psi_1 | (S_{\text{tot}})_z | \Psi_1 \rangle &= \int [f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0 \quad 0 \quad 0] \begin{pmatrix} \hbar & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\hbar \end{pmatrix} \begin{bmatrix} f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \\ 0 \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \int \hbar |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = \hbar
\end{aligned}$$

$$\begin{aligned}
\langle \Psi_2 | (S_{\text{tot}})_z | \Psi_2 \rangle &= \int [0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0 \quad 0] \begin{pmatrix} \hbar & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\hbar \end{pmatrix} \begin{bmatrix} 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\langle \Psi_3 | (S_{\text{tot}})_z | \Psi_3 \rangle &= \int [0 \quad 0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2) \quad 0] \begin{pmatrix} \hbar & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\hbar \end{pmatrix} \begin{bmatrix} 0 \\ 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \\ 0 \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\langle \Psi_4 | (S_{\text{tot}})_z | \Psi_4 \rangle &= \int [0 \quad 0 \quad 0 \quad f^*(\mathbf{r}_1, \mathbf{r}_2)] \begin{pmatrix} \hbar & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\hbar \end{pmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ f(\mathbf{r}_1, \mathbf{r}_2) \end{bmatrix} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= - \int \hbar |f(\mathbf{r}_1, \mathbf{r}_2)|^2 d\mathbf{r}_1 d\mathbf{r}_2 = -\hbar
\end{aligned}$$

**Part e**

For  $\Psi(\mathbf{r}_1, \mathbf{r}_2) = f(\mathbf{r}_1, \mathbf{r}_2)\chi_1 \otimes \chi_2$  to be antisymmetric, we require that  $\Psi(\mathbf{r}_1, \mathbf{r}_2) = -\Psi(\mathbf{r}_2, \mathbf{r}_1)$ . This means that we require

$$f(\mathbf{r}_1, \mathbf{r}_2)\chi_1 \otimes \chi_2 = -f(\mathbf{r}_2, \mathbf{r}_1)\chi_2 \otimes \chi_1$$

that is if  $\chi_1 \otimes \chi_2 = \chi_2 \otimes \chi_1$  (i.e.  $\chi_1 = \chi_2$ ) then we have that we require  $f(\mathbf{r}_1, \mathbf{r}_2) = -f(\mathbf{r}_2, \mathbf{r}_1)$ , that is  $f$  must be antisymmetric. If however  $\chi_1 \neq \chi_2$  then

**Part f**