

Problem Set 1

*Student Name: Noah Reef***Problem 1****Problem 2****Part a**

$$L(x, \dot{x}, t) = T - V = \frac{1}{2}m\dot{x}^2 - \frac{1}{2}kx^2$$

Part b

Using the Euler-lagrange equation, we have:

$$\frac{\partial L}{\partial x} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = -kx - m\ddot{x} = 0$$

solving the differential equation above gives the general solution,

$$x(t) = c_2 \sin(\omega t) + c_1 \cos(\omega t)$$

where $\omega = \sqrt{\frac{k}{m}}$. Then we see with boundary equations that

$$x(0) = c_1 = x_1$$

$$x(\tau) = c_2 \sin(\omega\tau) + x_1 \cos(\omega\tau) = x_2$$

solving for c_2 gives

$$c_2 = \frac{x_2 - x_1 \cos(\omega\tau)}{\sin(\omega\tau)}$$

since $\tau \neq \frac{n\pi}{\omega}$, we have that $\sin(\omega\tau) \neq 0$.

Part c

To solve for the action S , we have that

$$\begin{aligned} S &= \int_0^\tau L(x, \dot{x}, t) dt \\ &= \int_0^\tau \left(\frac{1}{2}m\dot{x}^2 - \frac{1}{2}kx^2 \right) dt \\ &= \frac{1}{2}m \int_0^\tau \dot{x}^2 - \omega^2 x^2 dt \end{aligned}$$

Now notice that

$$\int_0^\tau \dot{x}^2 dt = x\dot{x}\Big|_0^\tau - \int_0^\tau x\ddot{x} dt$$

and since $\ddot{x} = -\omega^2 x$, we have that

$$\int_0^\tau \dot{x}^2 dt = x\dot{x}\Big|_0^\tau + \omega^2 \int_0^\tau x^2 dt$$

Thus, we have that

$$S = \frac{1}{2}m \left(x\dot{x}\Big|_0^\tau + \omega^2 \int_0^\tau x^2 dt - \omega^2 \int_0^\tau x^2 dt \right)$$

thus

$$S = \frac{1}{2}m \left(x\dot{x}\Big|_0^\tau \right)$$

to solve the inside term we have that

$$x\dot{x} = (c_2 \sin(\omega t) + x_1 \cos(\omega t)) (\omega c_2 \cos(\omega t) - x_1 \omega \sin(\omega t))$$

now evaluating at τ and 0 gives

$$\begin{aligned} x\dot{x}\Big|_0^\tau &= (c_2 \sin(\omega\tau) + x_1 \cos(\omega\tau)) (\omega c_2 \cos(\omega\tau) - x_1 \omega \sin(\omega\tau)) - x_1 \omega c_2 \\ &= \left(\frac{x_2 - x_1 \cos(\omega\tau)}{\sin(\omega\tau)} \sin(\omega\tau) + x_1 \cos(\omega\tau) \right) \left(\frac{\omega x_2 \cos(\omega\tau) - \omega x_1 \cos^2(\omega\tau)}{\sin(\omega\tau)} - x_1 \omega \sin(\omega\tau) \right) - \dots \\ &= \frac{\omega x_2^2 \cos(\omega\tau) - \omega x_2 x_1}{\sin(\omega\tau)} - \frac{\omega x_1 x_2 - \omega x_1^2 \cos(\omega\tau)}{\sin(\omega\tau)} \\ &= \frac{\omega}{\sin(\omega\tau)} (-2x_1 x_2 + (x_1^2 + x_2^2) \cos(\omega\tau)) \end{aligned}$$

thus

$$S = \frac{m\omega}{2 \sin(\omega\tau)} (-2x_1 x_2 + (x_1^2 + x_2^2) \cos(\omega\tau))$$

Part d

To write the action $S(x, \tau)$, we set $x_2 = x$ and $x_1 = 0$ to get

$$S(x, \tau) = \frac{m\omega}{2 \sin(\omega\tau)} (-2x + x^2 \cos(\omega\tau))$$

to solve for the action of a silicon atom, with mass $m_{\text{Si}} = 4.66 \times 10^{-26} \text{kg}$, vibrating at frequency $\omega = 15 \text{THz}$ and amplitude $x = a_0$ and a time $\tau = \pi/(4\omega)$