

## Analysis of Antarctic sectors

There are three sectors available:

1. SA := South Africa,
2. WS := Weddell Sea,
3. and EA := East Antarctica

### Table of contents

1. Preamble
2. Ice concentration (aice)
3. Pancake ice concentration
4. Pancake ice width
5. Pancake ice growth
6. SWH presence
7. SWH distance into the ice
8. Ice velocities
9. Wind velocities
10. Win and ice correlation
11. Ice thickness

Alberello et al. (2019) *Pancake ice FSD in the Antarctic MIZ.*:

Sector: SA

Measurements: Peak period, SWH, ice concentration, pancake FSD

Dates: 4 July 2017

### 1. Preamble

```
%% Read in the data.
clear
close all
addpath functions
addpath packages/quiverwcolorbar
%ncdisp(filename)

% Parameters
sector = "SA";
grid = 'gx1';
case_name = '3month';
```

```

filedir = 'cases/3month/history/iceh.2005-09-30.nc';
[lat,lon,row] = grid_read(grid);

% Make sector mask
[len,wid] = size(lat);


```

## 2. Ice area

```

variable = "aice";
sector_data = data_format_sector(filedir,variable,sector);
%data = data(sector_mask);
%color_map = seaicecolormap();
latitude = [-90,-60];
longitude = [10,50];
figure(1)
w = worldmap('world');
    axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
    setm(w, 'Origin', [0 0 0]);
    setm(w, 'maplatlimit', [-70,-40]);
    setm(w, 'maplonlimit', [10,50]);
    setm(w, 'meridianlabel', 'on')
    setm(w, 'parallellabel', 'on')
    setm(w, 'mlabellocation', 10);
    setm(w, 'plabellocation', 10);
    setm(w, 'mlabelparallel', 0);
    setm(w, 'grid', 'on');
    setm(w, 'frame', 'on');
    setm(w, 'labelrotation', 'on')
pcolorm(lat,lon,sector_data)
land = shaperead('landareas', 'UseGeoCoords', true);
geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
colorbar
caxis(colorlims(variable));
title(strcat("AICE in the ", sector_name(sector)), 'interpreter','latex','FontSize', 1

```



### 3. Pancake ice concentration

Floe size categories:

```
fsd_cats = ncread(filedir,'NFSD');
nfsd = length(fsd_cats);
fprintf('NFSD is: %g', nfsd)
```

NFSD is: 16

```
fprintf('The FSD categories are: [');
```

The FSD categories are: [

```
fprintf('%g ', fsd_cats);
```

2.6884 9.79845 21.6721 40.7349 70.1407 113.694 175.577 259.837 369.62 506.24 668.209 850.

```
fprintf(']\n');
```

```
]
```

Define floes with radii  $< 10\text{m}$  as pancakes (i.e., the smallest two FSD categories).


```
variable = "afsd";
% Grid
grid = "gx1";
[lat,lon,row] = grid_read(grid);
% Read in data
dim = 3;
data = data_format(filedir,variable,row,lat,lon,dim);
data = data(:,:,1) + data(:,:,2); % Smallest 2 categories combined (ie. pancake)
ocean_mask = data_format(filedir,'tmask',row,lat,lon);
% Coordinates
coords = sector_coords(sector); % (NW;SW,NE,SE) (lat,lon)
for i = 1:4
    [lat_out(i),lon_out(i)] = lat_lon_finder(coords(i,1),coords(i,2),lat,lon);
end
[len,wid] = size(data);
sector_data = zeros(len,wid);
sector_mask = false(len,wid);
sector_data = ~ocean_mask*NaN;
for i = 0:lat_out(1)-lat_out(2)
    sector_mask(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = true;
    sector_data(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = data(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3));
end

latitude = [-90,-60];
longitude = [10,50];
figure(3)
w = worldmap('world');
axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
setm(w, 'Origin', [0 0 0]);
setm(w, 'maplatlimit', [-70,-40]);
setm(w, 'maplonlimit', [10,50]);
setm(w, 'meridianlabel', 'on')
setm(w, 'parallellabel', 'on')
setm(w, 'mlabellocation', 10);
setm(w, 'plabellocation', 10);
setm(w, 'mlabelparallel', 0);
```

```

setm(w, 'grid', 'on');
setm(w, 'frame', 'on');
setm(w, 'labelrotation', 'on')
pcolorm(lat,lon,sector_data)
land = shaperead('landareas', 'UseGeoCoords', true);
geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
colorbar
caxis(colorlims(variable));
title(strcat("Pancakce ice concentration in the ", sector_name(sector)), 'interpreter

```



```

%colormap turbo

```

### 3. a) FSD into the ice pack

```

addpath packages/joyPlot
variable = "afsd";
transect_coords = [coords(1,1), (coords(1,2) + coords(3,2))/2];
% Grid
grid = "gx1";
[lat,lon,row] = grid_read(grid);
% Read in data

```

```

dim = 3;
data = data_format(filedir,variable,row,lat,lon,dim);
data_pancake = data(:,:,1) + data(:,:,2); % Smallest 2 categories combined (ie. pancake)
ocean_mask = data_format(filedir,'tmask',row,lat,lon);
% Coordinates
coords = sector_coords(sector); % (NW;NE;SW;SW) (lat,lon)
for i = 1:4
    [lat_out(i),lon_out(i)] = lat_lon_finder(coords(i,1),coords(i,2),lat,lon);
end
[len,wid] = size(data_pancake);
sector_data = zeros(len,wid);
sector_mask = false(len,wid);
sector_data = ~ocean_mask*NaN;
for i = 0:lat_out(1)-lat_out(2)
    sector_mask(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = true;
    sector_data(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = data_pancake(lon_out(1)
end

[lat_tran,lon_tran] = lat_lon_finder(transect_coords(1),transect_coords(2),lat,lon);

aice = data_format(filedir,'aice',row,lat,lon);
ice_idx = false(1,lat_tran);
for j = lat_tran:-1:1
    if aice(lon_tran,j)>eps
        ice_idx(j) = true;
    end
end
lat_subset = lat(lon_tran,lat_tran:-1:1);

lat_subset_ice = lat_subset(ice_idx);

for j = lat_tran:-1:1
    if ice_idx(j) == 1
        for i = 1:length(fsd_cats)
            transect_fsd(j,i) = data(lon_tran,j,i);
        end
    end
end

transect_fsd_subset = transect_fsd(ice_idx,:);
transect_fsd_data = transect_fsd_subset;

threshold = 0;
if threshold == 1
    transect_fsd_threshold = false(1,nfsd);


```

```

    for i = 1:nfsd
        if sum(transect_fsd(:,i))>0.05
            transect_fsd_threshold(i) = true;
        end
    end

    for i = 1:length(lat_subset_ice)
        mean_vec(i) = mean(transect_fsd_subset(i,:));
    end
    transect_fsd_subset_threshold = transect_fsd_subset(:,transect_fsd_threshold);
    transect_fsd_data = transect_fsd_subset_threshold;
end
Ndecimals = 1;
f = 10.^Ndecimals;
% Take first 3 categories
transect_fsd_data = transect_fsd_data(:,1:2);
[len,wid] = size(transect_fsd_data);
figure(31)
joyPlot(transect_fsd_data',1:wid,0.05,'FaceColor','position','FaceAlpha',0.6)
set(gca,'Color',[0.93 0.93 0.93])
set(gca,'XGrid','on')
colormap(spring)
xlabel("FSD categories")
ylabel(['Latitudes along 30' char(176), 'E'])
% Create a char array with alphabet
lat_lbs = round(f.*lat_subset_ice)./f;
n = size(transect_fsd_subset);
% Formatting
ylbls = sprintfc(['%g', char(176)],lat_lbs(n:-1:1));
% Apply
yticklabels(ylbls)

```



```
% title("Cell rest")
```

#### 4. Pancake ice width

#### 5. Pancake ice growth

```
variable = "fsdrad";
sector_data = data_format_sector(filedir,variable,sector);


%data = data(sector_mask);
%color_map = seaicecolormap();
%colormap turbo
latitude = [-90,-60];
longitude = [10,50];
figure(5)
w = worldmap('world');
axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
setm(w, 'Origin', [0 0 0]);
```



```

setm(w, 'maplatlimit', [-70,-40]);
setm(w, 'maplonlimit', [10,50]);
setm(w, 'meridianlabel', 'on')
setm(w, 'parallellabel', 'on')
setm(w, 'mlabellocation', 10);
setm(w, 'plabellocation', 10);
setm(w, 'mlabelparallel', 0);
setm(w, 'grid', 'on');
setm(w, 'frame', 'on');
setm(w, 'labelrotation', 'on')
pcolorm(lat,lon,sector_data)
land = shaperead('landareas', 'UseGeoCoords', true);
geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
colorbar
caxis(colorlims(variable));
title(strcat("FSDRAD in the ", sector_name(sector)), 'interpreter','latex','FontSize'

```



```

%colormap turbo
% variable = "fsdrad";
% sector_data = data_format_sector(filedir,variable,sector);
% %data = data(sector_mask);
% %color_map = seaicecolormap();

```

```

% %colormap turbo
% latitude = [-90,-60];
% longitude = [10,50];
% figure(51)
% w = worldmap('world');
% axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to e
% setm(w, 'Origin', [0 0 0]);
% setm(w, 'maplatlimit', [-70,-40]);
% setm(w, 'maplonlimit', [10,50]);
% setm(w, 'meridianlabel', 'on')
% setm(w, 'parallellabel', 'on')
% setm(w, 'mlabellocation', 10);
% setm(w, 'plabellocation', 10);
% setm(w, 'mlabelparallel', 0);
% setm(w, 'grid', 'on');
% setm(w, 'frame', 'on');
% setm(w, 'labelrotation', 'on')
% pcolorm(lat,lon,sector_data)
% land = shaperead('landareas', 'UseGeoCoords', true);
% geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
% colorbar
% caxis(colorlims(variable));
% title(strcat("FSDRAD in the ", sector_name(sector)), 'interpreter','latex','FontSize
% %colormap turbo

```

```

% variable = "afsd";
% data = data_format(filedir,variable,row,lat,lon,3);
% [len,wid,~] = size(data);
% nanmask = isnan(data(:, :, 1));
% for k = 1:16
%     data(nanmask,:) = 0;
% end
% for i = 1:len
%     for j = 1:wid
%         %mass = 0;
%         %for k = 1:16
%             % mass = mass + data(i,j,k)*fsd_cats(k);
%         %end
%         sector_data(i,j) = sum(data(i,j,:).*fsd_cats);
%     end
% end
% %data = data(sector_mask);
% %color_map = seaicecolormap();
% %colormap turbo

```

```

% latitude = [-90,90];
% longitude = [-180,180];
% figure(53)
% %variable = 'fsdrad';
% w = worldmap('world');
% axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eqaa
% setm(w, 'Origin', [0 0 0]);
% setm(w, 'maplatlimit', [-70,-40]);
% setm(w, 'maplonlimit', [10,180]);
% setm(w, 'meridianlabel', 'on')
% setm(w, 'parallellabel', 'on')
% setm(w, 'mlabellocation', 10);
% setm(w, 'plabellocation', 10);
% setm(w, 'mlabelparallel', 0);
% setm(w, 'grid', 'on');
% setm(w, 'frame', 'on');
% setm(w, 'labelrotation', 'on')
% pcolorm(lat,lon,sector_data)
% land = shaperead('landareas', 'UseGeoCoords', true);
% geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
% colorbar
% caxis([0,50]);
% title(strcat("FSDRAD (calculation) in the ", sector_name(sector)), 'interpreter','latex
% %colormap turbo

```

## 6. SWH presence

```

variable = "wave_sig_ht";
day = 1;
month = 9;
year = 2005;
date = sprintf('%d-0%d-0%d', year, month, day);
dim = 2;
datapoints = 24;

for i = 1:datapoints
    % Get the file name
    filename = strcat("cases/",case_name,"/history/iceh.",date,".nc");
    data = data_format(filedir,variable,row,lat,lon,dim);
    %dx = data_formatted > fsd_max;
    %swh_miz = data_formatted;
    %swh_miz(idx) = 0.0;


```

```

        %idx = swh_miz > wave_height;
        %swh_miz(idx) = 1.0;
        % only print SWH in areas where there is sea ice
        %mask = ice_mask(case_name,date,grid,SIC);
        total_data(:,:,i) = data;
        date = update_date(date);
    end
    % Proportion of days per season (datapoints)
    swh_data = sum(total_data,3)/datapoints;

    latitude = [-90,-60];
    longitude = [10,50];
    figure(6)
    w = worldmap('world');
        axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
        setm(w, 'Origin', [0 0 0]);
        setm(w, 'maplatlimit', [-70,-40]);
        setm(w, 'maplonlimit', [10,50]);
        setm(w, 'meridianlabel', 'on')
        setm(w, 'parallellabel', 'on')
        setm(w, 'mlabellocation', 10);
        setm(w, 'plabellocation', 10);
        setm(w, 'mlabelparallel', 0);
        setm(w, 'grid', 'on');
        setm(w, 'frame', 'on');
        setm(w, 'labelrotation', 'on')
    pcolorm(lat,lon,data)
    land = shaperead('landareas', 'UseGeoCoords', true);
    geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
    colorbar
    caxis(colorlims(variable));
    title(strcat("Mean SWH in the ", sector_name(sector), " over September"), 'interpreter

```



```
%colormap turbo
```

## 7. SWH distance into the ice

## 8. Ice velocities

```
%% ICE velocities
variable = "uvel"; % ice velocity (x)
ice_data_x_raw = data_format_sector(filedir,variable,sector);
variable = "vvel"; % ice velocity (y)
ice_data_y_raw = data_format_sector(filedir,"vvel",sector);
lat0 = lat(sector_mask);
lon0 = lon(sector_mask);
ice_data_x = ice_data_x_raw(sector_mask);
ice_data_y = ice_data_y_raw(sector_mask);


latitude = [-90,-60];
longitude = [10,50];
```

```

deltalat_ice = ice_data_x;
deltalon_ice = ice_data_y;
scale = 1;

figure(8)
w = worldmap('world');
    axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
    setm(w, 'Origin', [0 0 0]);
    setm(w, 'maplatlimit', [-70,-40]);
    setm(w, 'maplonlimit', [10,50]);
    setm(w, 'meridianlabel', 'on')
    setm(w, 'parallellabel', 'on')
    setm(w, 'mlabellocation', 10);
    setm(w, 'plabellocation', 10);
    setm(w, 'mlabelparallel', 0);
    setm(w, 'grid', 'on');
    setm(w, 'frame', 'on');
    setm(w, 'labelrotation', 'on')
    quiverm(lat0,lon0,deltalat_ice,deltalon_ice,'r')
    land = shaperead('landareas', 'UseGeoCoords', true);
    geoshow(w, land, 'FaceColor', [0.5 0.7 0.5]);
    title(strcat("Ice velocity (m/s)", sector_name(sector)), 'interpreter','latex','FontS

```



```

lat_min = lat_lon_finder(coords(2,1),coords(2,2),lat,lon);
lat_range = lat(lon_tran,lat_tran:-1:lat_min);

[wid,len] = size(lat);
ice_vec_x = reshape(ice_data_x,1,[]);
ice_vec_y = reshape(ice_data_y,1,[]);
lat1 = reshape(lat0,1,[]);
lon1 = reshape(lon0,1,[]);
deltalat_ice_vec = reshape(deltalat_ice,1,[]);
deltalon_ice_vec = reshape(deltalon_ice,1,[]);
mag_ice = sqrt(deltalat_ice_vec.^2+deltalon_ice_vec.^2);
scale = 2;
figure(80)
quiverwcolorbar(lon1',-lat1',deltalon_ice_vec',-deltalat_ice_vec',scale,'bounds',[min(mag


```

Warning: The colorbar TickLength is a scalar quantity. Only the first value will be used.

```

a = colorbar;
a.Label.String = 'Velocity (m/s)';
title('Ice velocity')
xlabel('Longitude (degrees East)');
ylabel('Latitude (degrees South)');

```



"The speed reached  $\approx 0.75m/s$  , which is the highest ice speed ever recorded in the Southern Ocean to our knowledge" Alberello et al. (2020) *Drift of pancake ice*.

## 9. Wind velocity

```
%% ATM velocities
variable = "uatm"; % atm velocity (x)
atm_data_x = data_format_sector(filedir,variable,sector);
atm_data_y = data_format_sector(filedir,"vatm",sector);
lat0 = lat(sector_mask);
lon0 = lon(sector_mask);
atm_data_x = atm_data_x(sector_mask);
atm_data_y = atm_data_y(sector_mask);

latitude = [-90,-60];
longitude = [10,50];
```




```

deltalat_atm = atm_data_x;
deltalon_atm = atm_data_y;

figure(9)
w = worldmap('world');
    axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
    setm(w, 'Origin', [0 0 0]);
    setm(w, 'maplatlimit', [-70,-40]);
    setm(w, 'maplonlimit', [10,50]);
    setm(w, 'meridianlabel', 'on')
    setm(w, 'parallellabel', 'on')
    setm(w, 'mlabellocation', 10);
    setm(w, 'plabellocation', 10);
    setm(w, 'mlabelparallel', 0);
    setm(w, 'grid', 'on');
    setm(w, 'frame', 'on');
    setm(w, 'labelrotation', 'on')
    quiverm(lat0,lon0,deltalat_atm,deltalon_atm,'r')
    land = shaperead('landareas', 'UseGeoCoords', true);
    geoshow(w, land, 'FaceColor', [0.5 0.7 0.5]);
    %a = colorbar;
    %a.Label.String = 'Velocity (m/s)';
    title(strcat("Atmos. velocity (m/s)", sector_name(sector)), 'interpreter','latex','For

```



```
lat_min = lat_lon_finder(coords(2,1),coords(2,2),lat,lon);
lat_range = lat(lon_tran,lat_tran:-1:lat_min);


[wid,len] = size(lat);
atm_vec_x = reshape(atm_data_x,1,[]);
atm_vec_y = reshape(atm_data_y,1,[]);
lat1 = reshape(lat0,1,[]);
lon1 = reshape(lon0,1,[]);

deltalat_atm_vec = reshape(deltalat_atm,1,[]);
deltalon_atm_vec = reshape(deltalon_atm,1,[]);
mag_atm = sqrt(deltalat_atm_vec.^2+deltalon_atm_vec.^2);
scale = 0.3;
figure(90)
quiverwcolorbar(lon1',-lat1',deltalon_atm_vec',-deltalat_atm_vec',scale,'bounds',[min(mag
```

Warning: The colorbar TickLength is a scalar quantity. Only the first value will be used.

```
a = colorbar;
a.Label.String = 'Velocity (m/s)';
xlabel('Longitude (degrees East)');
```

```
ylabel('Latitude (degrees South)');
```



## 10. Wind and ice velocity correlaion


```
ice_lat_mask = deltalat_ice>eps;
ice_lon_mask = deltalon_ice>eps;
% Apply ice mask, so we only take wind data where there is ice
deltalat_atm_mask = deltalat_atm.*ice_lat_mask;
deltalon_atm_mask = deltalon_atm.*ice_lat_mask;
% Do we need this?
atm_vec = [deltalat_atm_mask, deltalon_atm_mask]; %(lat,lon)
ice_vec = [deltalat_ice, deltalon_ice]; %(lat,lon)
%% How far the did winds move the ice?
% Project ice onto atmosphere
proj = dot(ice_vec,atm_vec)/dot(atm_vec,atm_vec)*atm_vec;

%deltalat_corr = deltalat_atm.*ice_lat_mask + deltalat_ice;
%deltalon_corr = deltalon_atm.*ice_lon_mask + deltalon_ice;
```

```

figure(10)
w = worldmap('world');
    axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
    setm(w, 'Origin', [0 0 0]);
    setm(w, 'maplatlimit', [-70,-40]);
    setm(w, 'maplonlimit', [10,50]);
    setm(w, 'meridianlabel', 'on')
    setm(w, 'parallellabel', 'on')
    setm(w, 'mlabellocation', 10);
    setm(w, 'plabellocation', 10);
    setm(w, 'mlabelparallel', 0);
    setm(w, 'grid', 'on');
    setm(w, 'frame', 'on');
    setm(w, 'labelrotation', 'on')
    quiverm(lat0,lon0,proj(:,1),proj(:,2),'r')
    land = shaperead('landareas', 'UseGeoCoords', true);
    geoshow(w, land, 'FaceColor', [0.5 0.7 0.5]);
    %a = colorbar;
    %a.Label.String = 'Velocity (m/s)';
    title(strcat("Joint ice and atmos. velocity (m/s)", sector_name(sector)), 'interpreter

```



```

%% What influence does wind have on the ice velocity?
[len, ~] = size(proj);
for i = 1:len
    mag_proj(i) = sqrt(proj(i,1)^2+proj(i,2)^2);
end
scale = 2;
figure(99)
quiverwcolorbar(lon1',-lat1',proj(:,2),-proj(:,1),scale,'bounds',[min(mag_proj) max(mag_p


```

Warning: The colorbar TickLength is a scalar quantity. Only the first value will be used.

```

a = colorbar;
a.Label.String = 'Velocity (m/s)';
title('Winds effect on ice velocity')
xlabel('Longitude (degrees East)');
ylabel('Latitude (degrees South)');

```



```

% Calculate percentage ice velocity is of wind velocity in that direction
%cos(proj,atm_vec)
for i = 1:length(proj)

```

```

        norm_ratio(i) = norm(proj(i,:))/norm(ice_vec(i,:));
    end

    nan_idx = isnan(norm_ratio);
    error_idx = norm_ratio > 1;

    norm_ratio_adjusted = norm_ratio;
    norm_ratio_adjusted(nan_idx) = 0;
    norm_ratio_adjusted(error_idx) = 0;
    zero_idx = norm_ratio_adjusted > eps;
    mean(norm_ratio_adjusted(zero_idx));

```

## 11. Ocean current

```

% % uocn and vocn
% variable = "uocn"; % ocn velocity (x)
% ocn_data_x = data_format_sector(filedir,variable,sector);
% ocn_data_y = data_format_sector(filedir,"vocn",sector);
% lat0 = lat(sector_mask);
% lon0 = lon(sector_mask);
% ocn_data_x = ocn_data_x(sector_mask);
% ocn_data_y = ocn_data_y(sector_mask);
%
% latitude = [-90,-60];
% longitude = [10,50];
%
% deltalat_ocn = ocn_data_x;
% deltalon_ocn = ocn_data_y;
%
%
% figure(11)
% w = worldmap('world');
%     axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to e
%     setm(w, 'Origin', [0 0 0]);
%     setm(w, 'maplatlimit', [-70,-40]);
%     setm(w, 'maplonlimit', [10,50]);
%     setm(w, 'meridianlabel', 'on')
%     setm(w, 'parallellabel', 'on')
%     setm(w, 'mlabellocation', 10);
%     setm(w, 'plabellocation', 10);
%     setm(w, 'mlabelparallel', 0);
%     setm(w, 'grid', 'on');
%     setm(w, 'frame', 'on');
%     setm(w, 'labelrotation', 'on')

```

```

%     quiverm(lat0,lon0,deltalat_ocn,deltalon_ocn,'r')
%     land = shaperead('landareas', 'UseGeoCoords', true);
%     geoshow(w, land, 'FaceColor', [0.5 0.7 0.5]);
%     %a = colorbar;
%     %a.Label.String = 'Velocity (m/s)';
%     title(strcat("Ocean velocity (m/s)", sector_name(sector)), 'interpreter','latex','F
%
% lat_min = lat_lon_finder(coords(2,1),coords(2,2),lat,lon);
% lat_range = lat(lon_tran,lat_tran:-1:lat_min);
%
% [wid,len] = size(lat);
% ocn_vec_x = reshape(ocn_data_x,1,[]);
% ocn_vec_y = reshape(ocn_data_y,1,[]);
% lat1 = reshape(lat0,1,[]);
% lon1 = reshape(lon0,1,[]);
%
% deltalat_ocn_vec = reshape(deltalat_ocn,1,[]);
% deltalon_ocn_vec = reshape(deltalon_ocn,1,[]);
% mag_ocn = sqrt(deltalat_ocn_vec.^2+deltalon_ocn_vec.^2);
% scale = 0.3;
% figure(41)
% quiverwcolorbar(lon1',-lat1',deltalon_ocn_vec',-deltalat_ocn_vec',scale);
%     a = colorbar;
%     a.Label.String = 'Velocity (m/s)';
%     xlabel('Longitude (degrees East)');
%     ylabel('Latitude (degrees South)');

```

## 12. Atm/ice stress

```

% strairx and strairy
variable = "strairx"; % (x)
ocn_data_x = data_format_sector(filedir,variable,sector);
ocn_data_y = data_format_sector(filedir,"strairy",sector);
lat0 = lat(sector_mask);
lon0 = lon(sector_mask);
ocn_data_x = ocn_data_x(sector_mask);
ocn_data_y = ocn_data_y(sector_mask);

latitude = [-90,-60];
longitude = [10,50];

deltalat_ocn = ocn_data_x;


```

```

deltalon_ocn = ocn_data_y;

figure(12)
w = worldmap('world');
axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
setm(w, 'Origin', [0 0 0]);
setm(w, 'maplatlimit', [-70,-40]);
setm(w, 'maplonlimit', [10,50]);
setm(w, 'meridianlabel', 'on')
setm(w, 'parallellabel', 'on')
setm(w, 'mlabellocation', 10);
setm(w, 'plabellocation', 10);
setm(w, 'mlabelparallel', 0);
setm(w, 'grid', 'on');
setm(w, 'frame', 'on');
setm(w, 'labelrotation', 'on')
quiverm(lat0,lon0,deltalat_ocn,deltalon_ocn,'r')
land = shaperead('landareas', 'UseGeoCoords', true);
geoshow(w, land, 'FaceColor', [0.5 0.7 0.5]);
%a = colorbar;
%a.Label.String = 'Velocity (m/s)';
title(strcat("Ocean velocity (m/s) in the x direction", sector_name(sector)), 'interp

```





```

lat_min = lat_lon_finder(coords(2,1),coords(2,2),lat,lon);
lat_range = lat(lon_tran,lat_tran:-1:lat_min);

[wid,len] = size(lat);
ocn_vec_x = reshape(ocn_data_x,1,[]);
ocn_vec_y = reshape(ocn_data_y,1,[]);
lat1 = reshape(lat0,1,[]);
lon1 = reshape(lon0,1,[]);

deltalat_ocn_vec = reshape(deltalat_ocn,1,[]);
deltalon_ocn_vec = reshape(deltalon_ocn,1,[]);
mag_ocn = sqrt(deltalat_ocn_vec.^2+deltalon_ocn_vec.^2);
scale = 1.5;
figure(42)
quiverwcolorbar(lon1',-lat1',deltalon_ocn_vec',-deltalat_ocn_vec',scale,'bounds',[min(mag


```

Warning: The colorbar TickLength is a scalar quantity. Only the first value will be used.

```


a = colorbar;
a.Label.String = 'Velocity (m/s)';
xlabel('Longitude (degrees East)');
ylabel('Latitude (degrees South)');

```



### 13. Ice thickness

```
variable = "hi";
sector_data = data_format_sector(filedir,variable,sector);
%data = data(sector_mask);
%color_map = seaicecolormap();
%colormap turbo
latitude = [-90,-60];
longitude = [10,50];
figure(13)
w = worldmap('world');
    axesm miller; %, eqaazim eqdazim vperspec, eqdazim flips the x-axis, and y-axis to eq
    setm(w, 'Origin', [0 0 0]);
    setm(w, 'maplatlimit', [-70,-40]);
    setm(w, 'maplonlimit', [10,50]);
    setm(w, 'meridianlabel', 'on')
    setm(w, 'parallellabel', 'on')
    setm(w, 'mlabellocation', 10);
    setm(w, 'plabellocation', 10);
    setm(w, 'mlabelparallel', 0);
    setm(w, 'grid', 'on');
    setm(w, 'frame', 'on');
    setm(w, 'labelrotation', 'on')
    pcolorm(lat,lon,sector_data)
    land = shaperead('landareas', 'UseGeoCoords', true);
    geoshow(w, land, 'FaceColor', [0.5 0.7 0.5])
    colorbar
    caxis(colorlims(variable));
    title(strcat("Ice thickness in the ", sector_name(sector)), 'interpreter','latex','For
```



```
%colormap turbo
```

## Functions

```
function coords = sector_coords(sector)
% Coordinates of sector
% There are three sectors available:
%     SA := South Africa,
%     WS := Weddell Sea,
%     and EA := East Antarctica
if sector == "SA"
    coords = [-45,20;-65,20;-45,40;-65,40]; %(NW;SW,NE,SE)
elseif sector == "EA"
    coords = [];
elseif sector == "WS"
    coords = [];
end
end
```

```

function sector_data = data_format_sector(filedir,variable,sector,dim)
    if ~exist('dim', 'var')
        dim = 2;
    end

    % Grid
    grid = "gx1";
    [lat,lon,row] = grid_read(grid);
    % Read in data
    data = data_format(filedir,variable,row,lat,lon,dim);
    ocean_mask = data_format(filedir,'tmask',row,lat,lon);
    % Coordinates
    coords = sector_coords(sector); % (NW;NE;SW;SW) (lat,lon)
    for i = 1:4
        [lat_out(i),lon_out(i)] = lat_lon_finder(coords(i,1),coords(i,2),lat,lon);
    end
    [len,wid] = size(data);
    sector_data = zeros(len,wid);
    sector_mask = false(len,wid);
    sector_data = ~ocean_mask*NaN;
    for i = 0:lat_out(1)-lat_out(2)
        sector_mask(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = true;
        sector_data(lon_out(1):lon_out(3), lat_out(1)-i:lat_out(3)) = data(lon_out(1):lon.
    end

end

function name = sector_name(sector)
    if sector == "SA"
        name = "South African Antarctic";
    elseif sector == "WS"
        name = "Weddell Sea";
    elseif sector == "EA"
        name = "East Antarctic";
    else
        name = "";
    end

end
end

```