

Physics of Nuclear Reactor - Numerical Exercises

Noah Rotunno

December 27, 2024

To run an exercise, open CMD and run the main file. The scripts that run a simulation is always named as main :

```
1 preject_path/exercice_i> python main.py
```

Some scripts can create other files:

1. pdf : Figure used in the report
2. npy : file to save a numpy array. This is used for saving data for big convergence study.

WARNING : In certain convergence studies, the plots generated by the Python script differ from those presented in the report. This discrepancy arises because the report plots were produced using a higher number of simulations to obtain better plots. However, for efficiency and to reduce computation time, the number of simulations was lowered in the submission scripts.

1 Exercice #1 - Modeling a planar source of neutrons

1.1 Question #1




Figure 1.1: Analytical Solution for the neutron flux as a function of depth.

Flux values at the following locations (4 significant digits):

Flux value at x_0 : $13.5896 \text{ n cm}^{-2} \text{ s}^{-1}$

Flux value at 0: $12276.8979 \text{ n cm}^{-2} \text{ s}^{-1}$

1.2 Question #2

Relationship between Φ_i , Φ_{i+1} , Φ_{i-1} at any point within the material:

$$A\Phi_i + B\Phi_{i-1} + C\Phi_{i+1} = D \quad (1)$$

Coefficients of the matrix A (4 significant digits), for a mesh size of 0.1cm:

Coef $A_{i,i}$: -16.6037

Coef $A_{i-1,i}$: 8.2919

Coef $A_{i+1,i}$: 8.2919

Relationship between Φ_i , Φ_{i-1} , Φ_{i+1} ,
at the source:

$$A\Phi_1 + B\Phi_2 = C \quad (2)$$

at the RHS of the problem:

$$A\Phi_n + B\Phi_{n-1} = C \quad (3)$$

Associated coefficients of the matrix A (4 significant digits), for a mesh size of 0.1cm: at the source:

Coef $A_{1,i}$: -8.3119

Coef $A_{2,i}$: 8.2919

at the RHS:

Coef $A_{n-1,i}$: 8.2919

Coef $A_{n,i}$: -12.1536

1.3 Question #3

Flux values from the numerical solver at the following locations (4 significant digits):

Flux value at x_0 : 13.6244

Flux value at 0: 12280.5980

Add two sentences describing what you see in Figure 5.5 and a possible explanation:
: The error decrease as the mesh increase, meaning that the simulation converge. The order of convergence has been evaluated to ~ 1 .




Figure 1.2: Distance between the solutions at each mesh point for a mesh size of 0.1 cm.




Figure 1.3: Evolution with mesh size of the absolute error of $\Phi(x_0)$.

2 Exercice #2 - Modeling a planar reactor (1 group)

2.1 Question #1




Figure 2.1: Flux in the bare reactor for a mesh size of 0.1 cm.

Numerical solution for the bare system

keff (scientific format with 5 significant digits): 0.97305

Net current at the core boundary (scientific format with 5 significant digits): 0.075436

2.2 Question #2




Figure 2.2: Distance between the solutions at each mesh point for a mesh size of 0.1 cm.

Analytical solution for the bare system

keff (scientific format with 5 significant digits): 0.97356

Net current at the core boundary (scientific format with 5 significant digits): 0.075367

2.3 Question #3

Numerical solution for the reflected system

keff (scientific format with 5 significant digits): 1.12038

Net current at the core boundary (scientific format with 5 significant digits): -0.054402




Figure 2.3: Flux in the reflected reactor for a mesh size of 0.1 cm.

3 Exercice #3 - Modeling a planar reactor (2 groups)

3.1 Question #1 Numerical formulation

Relationship between Φ_i , Φ_{i+1} , Φ_{i-1} for group g at any point within the material:

$$A\Phi_i + B\Phi_{i-1} + C\Phi_{i+1} = D \quad (4)$$

Expression of the source term for the fast and thermal energy group:

$$\begin{aligned} b_1 &= \nu\Sigma_{f1}\Phi_1 + \nu\Sigma_{f2}\Phi_2 \\ b_2 &= \Sigma_{s1\rightarrow2}\Phi_1 \end{aligned} \quad (5)$$

3.2 Question #2 Analytical solution for the bare system

$$k_{eff} = \frac{\nu\Sigma_{f1}(B^2D_2 + \Sigma_{t2}) + \Sigma_s^{1\rightarrow2}\nu\Sigma_{f2}}{(B^2D_1 + \Sigma_{t1})(B^2D_2 + \Sigma_{t2}) - \Sigma_s^{1\rightarrow2}\Sigma_s^{2\rightarrow1}} \quad (6)$$

keff (scientific format with 5 significant digits): 1.08440

3.3 Question #3 Numerical solution of the bare homogeneous reactor

keff (scientific format with 5 significant digits): 1.08677

fast buckling (scientific format with 5 significant digits): 0.030229

thermal buckling (scientific format with 5 significant digits): 0.030348




Figure 3.1: Fast and thermal fluxes in the bare reactor for a mesh size of 0.1 cm.

3.4 Question #4




Figure 3.2: Difference between numerical and analytical solutions for the fast and thermal fluxes in the bare reactor for a mesh size of 0.1 cm.

3.5 Question #5 Numerical solution of the reflected reactor

keff (scientific format with 5 significant digits): 1.09199

fast net current (scientific format with 5 significant digits) at the core/reflector interface: -0.039511

thermal net current (scientific format with 5 significant digits) at the core/reflector interface: 0.0058747




Figure 3.3: Fast and thermal fundamental fluxes in the reflected reactor for a mesh size of 0.1 cm.




Figure 3.4: Fast and thermal first harmonic of the fluxes in the reflected reactor for a mesh size of 0.1 cm.

first harmonic eigenvalue (scientific format with 5 significant digits): 1.02113

4 Exercice #4 - Modeling a more realistic planar reactor in two groups

4.1 Question #1 Initial Loading Pattern

keff (scientific format with 5 significant digits): 1.21882

peak power (scientific format with 5 significant digits): $7.37503 \cdot 10^5$ W

fast flux at the core boundary (scientific format with 5 significant digits): $1.87308 \cdot 10^{15}$ n cm⁻² s⁻¹




Figure 4.1: Normalized fast and thermal fluxes

Explanation of the main features of the power distribution: The newest batch are placed at the right boundary, meaning that more fission product are present in the middle of the reactor. This means that the neutrons are more absorbed in the middle, leading to a greater flux at the boundary. Moreover, the newest batch create more fission meaning more power and the reflector reflect a part of the neutrons at the boundary leading to a small increase in the power at the boundary.




Figure 4.2: Relative power distribution

4.2 Question #2 Optimized Loading Pattern

2	1	2	3	1	3	4
---	---	---	---	---	---	---

Table 1: Loading pattern from the core center (left to right)

keff (scientific format with 5 significant digits): 1.19399

peak power (scientific format with 2 significant digits): $6.53340 \cdot 10^5$ W

fast flux at the core boundary (scientific format with 2 significant digits): $6.28 \cdot 10^{13}$
n cm⁻² s⁻¹




Figure 4.3: Normalized fast and thermal fluxes




Figure 4.4: Relative power distribution

5 Exercice #5 - Fuel Evolution with exposure in a homogeneous media

5.1 Question #1 Analytical solution

U235	2.30941
U238	7.46708
Pu239	0.00000
X	0.00000
Y	0.00000

Table 2: Initial atomic concentrations in at/cm^{-3}

Governing differential equations for each isotope (U235,U238,Pu239, X and Y):

$$\begin{aligned}
\frac{dN_{U235}}{dt} &= -\Phi(\sigma_{c,U235} + \sigma_{f,U235})N_{U235} \\
\frac{dN_{U238}}{dt} &= -\Phi(\sigma_{c,U238} + \sigma_{f,U238})N_{U238} \\
\frac{dN_{Pu239}}{dt} &= -\Phi(\sigma_{c,Pu239} + \sigma_{f,Pu239})N_{Pu239} + \Phi\sigma_{c,U238}N_{U238} \\
\frac{dN_X}{dt} &= \Phi\sigma_{f,U235}FY_XN_{U235} - (\Phi\sigma_{c,X} + \lambda_X)N_X \\
\frac{dN_Y}{dt} &= \Phi\sigma_{f,U235}FY_YN_{U235} - \Phi\sigma_{c,Y}N_Y
\end{aligned} \tag{7}$$




Figure 5.1: Evolution of the various isotopes concentrations in a semilog-Y scale between 0 and 365 days




Figure 5.2: Evolution of the various isotopes concentrations in a semilog-Y scale between 0 and 1 day

5.2 Question #2 Numerical solution




Figure 5.3: Evolution of the error with respect to the analytical solution; for the various isotopes concentrations in a semilog-Y scale between 0 and 365 days with the Forward Euler method and a time-step of 1min




Figure 5.4: Illustration of the convergence rate of Euler and Matrix exponential methods, varying the time step between 1min and 6h. The quantity of interest is the concentration of X after 1 day.

5.3 Question #3 Advanced Problem




Figure 5.5: Evolution of the flux level ($\phi[cm^{-2}.s^{-1}]$) as a function of time.

Explain the observed trends in Figure 5.5 : Initially the flux decrease because of the production of ^{239}Pu that also produce power, leading to a smaller neutron flux to keep the power constant. The neutron flux decrease between 0 s and $0.87 \cdot 10^6$ s. Then the Y fission product accumulate and the amount of ^{235}U decrease, leading to a greater flux to keep the power constant. The flux increase after $0.87 \cdot 10^6$ s.




Figure 5.6: Evolution of the various isotopes concentrations in a semilog-Y scale between 0 and 180 days.