

Extremal quadratic maps in dimensions divisible by four

A uniform existence theorem and a concrete twelve bit instance

Research note and reproducible proof package · 9 October 2026

Verified scope. Independent source-only Lean replays passed for both the uniform theorem and the explicit twelve-bit certificate. The underlying construction is established; novelty of the exact count consequence is unestablished.

The uniform theorem

Theorem. For every integer $e \geq 1$ and $n = 4e$, there exists a homogeneous quadratic map $F: \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ of degree exactly two with

$$|N_F| = 3 \cdot 2^{n/2-1} - 1.$$

Here N_F is the set of nonzero masks whose Boolean component is not bent. With the cited inequality [1], this attains Deryck's non-MNBC lower bound for every $m=n$ divisible by four. It does not settle dimensions congruent to two modulo four. The universal lower bound itself is cited, not part of the formal development.

The explicit twelve bit instance

A concrete map with eight independent quadratic coordinates and four zero coordinates has 4000 bent and 95 nonbent nonzero masks. In little-endian binary encoding its full nonbent set is

$$N_F = \{b \in \{1, \dots, 4095\} : b \bmod 256 \in S\}, \\ S = \{0, 4, 8, 13, 14, 15\}.$$

The uniform proof makes classical choices of fields and coordinates. The explicit twelve-bit polynomial has a separate checked certificate; no byte-for-byte identification with the uniformly chosen $e=3$ witness is asserted. Sections 1 through 5 explain the concrete construction, and Appendix B proves the family.

Novelty and attribution

The underlying family is an established Jha-Johnson cyclic-semifield core plus a hyperbolic plane, identified by an explicit congruence. The exact extremal count is a short corollary; no prior publication of that consequence was located. Novelty is unestablished, and any contribution is at most a potentially new application or corollary. Appendix E gives the correspondence and sources.

"GPT-6 Astra" is the requested project label, not runtime model attribution.

1 The explicit construction

Let $k = F_2[\alpha]/(\alpha^3 + \alpha + 1)$. The cubic has no root in F_2 , hence is irreducible. Thus k is a field with eight elements, with binary basis $(1, \alpha, \alpha^2)$. In particular $\alpha^3 = \alpha + 1$ and $\alpha^4 = \alpha^2 + \alpha$. Its trace is

$$\text{Tr}(c) = c + c^2 + c^4 \in F_2.$$

For $\lambda = (u, v, w, z, s, t, p, q) \in F_2^8$, define a symmetric matrix K_λ over k with zero diagonal and these six upper-triangular entries:

$$\begin{aligned} K_{01} &= u + v + \alpha(s + t), \\ K_{02} &= u + \alpha s + \alpha^2 q, \\ K_{03} &= w + \alpha^2(p + q), \\ K_{12} &= z + \alpha^2 p, \\ K_{13} &= v + \alpha s + \alpha^2 q, \\ K_{23} &= u + v + \alpha t. \end{aligned}$$

All eight parameters are binary, including when the expressions are evaluated in k . The parameter label is $u + 2v + 4w + 8z + 16s + 32t + 64p + 128q$. The diagonal-zero symmetric matrix represents an alternating bilinear form in characteristic two.

Write an input $x \in F_2^{12}$ as $(X_0, X_1, X_2, X_3) \in k^4$, where

$$X_i = x_{3i} + \alpha x_{3i+1} + \alpha^2 x_{3i+2}.$$

Let e_j denote the j th binary parameter unit vector, with j beginning at zero. Define

$$\begin{aligned} F_j(x) &= \text{Tr}(\sum_{0 \leq i < l \leq 3} (K_{e_j})_{il} X_i X_l) \quad (0 \leq j < 8), \\ F_8 &= F_9 = F_{10} = F_{11} = 0. \end{aligned}$$

This is an explicit homogeneous quadratic map in the twelve binary variables: multiplication in k is binary bilinear, and trace is binary linear. Appendix A gives all 66 vector coefficients, eliminating any need to implement field arithmetic to evaluate F .

The eight effective coordinates are independent

The map $\lambda \mapsto K_\lambda$ is injective. If $K_\lambda = 0$, the basis coefficients in K_{02} force $u = s = q = 0$; K_{13} then gives $v = 0$; K_{12} gives $z = p = 0$; K_{03} gives $w = 0$; and K_{23} gives $t = 0$. Section 3 shows that trace also preserves this injectivity. Therefore the component kernel consists exactly of the four padded output-mask directions.

2 The Pfaffian calculation

For an alternating four by four matrix over a field of characteristic two, set

$$\text{pf}(K) = K_{01}K_{23} + K_{02}K_{13} + K_{03}K_{12}.$$

Expanding the determinant gives $\det(K) = \text{pf}(K)^2$: the three permutations consisting of two transpositions yield the three squared products, while the remaining nonzero determinant terms cancel in pairs. Hence K is singular exactly when $\text{pf}(K)=0$.

An explicit expansion

Put $A=u+v$. Use $a^2=a$ for each binary parameter and $\alpha^4=\alpha^2+\alpha$. The three products are

$$\begin{aligned} K_{01}K_{23} &= A + \alpha As + \alpha^2(st+t), \\ K_{02}K_{13} &= uv + \alpha As + \alpha^2(qA+s) + (\alpha^2+\alpha)q, \\ K_{03}K_{12} &= wz + \alpha^2(wp+z(p+q)) + (\alpha^2+\alpha)(p+pq). \end{aligned}$$

Adding the products cancels the αAs terms and gives

$$\begin{aligned} \text{pf}(K_\lambda) &= P + \alpha R + \alpha^2(Q+R), \\ P &= u+v+uv+wz, \\ R &= p+q+pq, \\ Q &= st+s+t+q(u+v)+wp+z(p+q). \end{aligned}$$

Exactly six singular parameter values

Linear independence of $1, \alpha, \alpha^2$ makes $\text{pf}(K_\lambda)=0$ equivalent to $P=Q=R=0$. For binary p, q , the expression $p+q+pq$ is zero only when $p=q=0$. With these values fixed, $Q=st+s+t$ is zero only when $s=t=0$. It remains to solve $P=0$.

The expression $u+v+uv$ is zero at $(u,v)=(0,0)$ and one at the other three pairs. There are therefore three solutions with $u=v=0$ and $wz=0$, and three with $(u,v) \neq (0,0)$ and $w=z=1$. Their labels are exactly

$$S = \{0, 4, 8, 13, 14, 15\}.$$

An alternating form has even rank. For completeness, if its value on a pair of vectors is nonzero, rescale the pair to obtain a nonsingular alternating two-dimensional block. Subtract multiples of that pair from each remaining basis vector to split off an orthogonal complement, and continue. Thus every nonzero singular four-dimensional alternating form has rank two.

Injectivity from Section 1 makes label zero the only zero matrix. Consequently, among the 256 matrices K_λ , one has rank zero, five have rank two, and 250 have rank four. This classification uses no numerical search.

3 Trace descent and binary rank

Nondegeneracy of the trace pairing

The trace is binary linear since squaring is additive in characteristic two. Also $\text{Tr}(c)^2 = \text{Tr}(c)$, using $c^8 = c$; the only roots of $T^2 + T$ are zero and one, so its values lie in F_2 . Finally $\text{Tr}(1) = 1 + 1 + 1 = 1$.

If $c \neq 0$, choose $d = c^{-1}$. Then $\text{Tr}(cd) = \text{Tr}(1) = 1$. Thus the pairing $(c, d) \mapsto \text{Tr}(cd)$ is nondegenerate over F_2 . On k^4 , the coordinatewise pairing has the same property: isolate any nonzero coordinate and choose its inverse in the other vector.

The radical is unchanged as a set

Associate to K_λ the binary bilinear form

$$B_\lambda(X, Y) = \text{Tr}(X^T K_\lambda Y).$$

If $K_\lambda Y = 0$, then $B_\lambda(X, Y) = 0$ for every X . Conversely, if $K_\lambda Y$ has a nonzero coordinate c , choose X supported at that coordinate with value c^{-1} . Then $B_\lambda(X, Y) = 1$, contradicting radical membership. Therefore

$$\text{rad}_{F_2}(B_\lambda) = \ker_k(K_\lambda) \text{ as subsets of } k^4.$$

A k -vector space of dimension d has binary dimension $3d$. If K_λ has k -rank r , the binary radical has dimension $3(4-r)$, and hence

$$\text{rank}_{F_2}(B_\lambda) = 12 - 3(4-r) = 3r.$$

This proves the binary rank distribution without relying on an elimination program:

Binary rank	Number of effective masks	Effective labels
0	1	0
6	5	4, 8, 13, 14, 15
12	250	All remaining labels

The component polar form

For $b \in F_2^{12}$, write $\lambda = (b_0, \dots, b_7)$ and $f_b(x) = \langle b, F(x) \rangle$. Linearity of trace and of K in λ gives

$$\begin{aligned} f_b(X) &= \text{Tr}(\sum_{i < l} (K_\lambda)_{il} X_i X_l), \\ f_b(X+Y) + f_b(X) + f_b(Y) &= B_\lambda(X, Y). \end{aligned}$$

In the ordered binary basis, its polar matrix M_λ has entries $M_\lambda[3i+r, 3l+h] = \text{Tr}(\alpha^{r+h}(K_\lambda)_{il})$. Nondegeneracy also shows $B_\lambda = 0$ only when $K_\lambda = 0$. Therefore no nonzero effective mask selects the zero quadratic function.

4 A self contained Walsh argument

Let $V = \mathbb{F}_2^n$ and let $q: V \rightarrow \mathbb{F}_2$ be quadratic with $q(0)=0$. Its polar form $B(x,z)=q(x+z)+q(x)+q(z)$ is alternating bilinear. Write R for its radical and $r=n-\dim R$ for its rank. For a frequency $a \in V$, define

$$W_q(a) = \sum_{x \in V} (-1)^{q(x) + \langle a, x \rangle}.$$

The square of a Walsh coefficient

In the product of two Walsh sums, replace the second variable by $x+z$. Bilinearity and character cancellation yield

$$\begin{aligned} W_q(a)^2 &= \sum_{z \in V} (-1)^{q(z) + \langle a, z \rangle} \sum_{x \in V} (-1)^{B(x,z)} \\ &= 2^n \sum_{z \in R} (-1)^{q(z) + \langle a, z \rangle}. \end{aligned}$$

Indeed, the inner sum equals 2^n for $z \in R$. Otherwise $x \mapsto B(x,z)$ is a nonzero linear functional; translation by a vector where it equals one pairs opposite signs and makes the sum zero.

The restriction $q|_R$ is linear because B vanishes on R . The remaining character sum is $|R|$ when $\langle a, \cdot \rangle$ agrees with q on R , and zero otherwise. Every linear functional on R extends to V . There are exactly $2^{n-\dim R} = 2^r$ such extensions, since the annihilator of R has that size. Consequently

$$\begin{aligned} W_q(a) &\in \{0, \pm 2^{n-r/2}\}, \\ \#\{a : W_q(a) \neq 0\} &= 2^r. \end{aligned}$$

If $r=n$, every coefficient has magnitude $2^{n/2}$, so q is bent. If $r < n$, some coefficient has larger magnitude and q is not bent. Thus a quadratic Boolean function is bent exactly when its polar form is nonsingular.

The signs in the spectrum

Summing $W_q(a)$ over a cancels every $x \neq 0$ and leaves $2^n(-1)^{q(0)} = 2^n$. If $A = 2^{n-r/2}$ and N_+, N_- count coefficients $+A, -A$, then

$$\begin{aligned} N_+ + N_- &= 2^r, \\ A(N_+ - N_-) &= 2^n. \end{aligned}$$

For $r > 0$ this gives $N_{\pm} = 2^{r-1} \pm 2^{r/2-1}$. There are $2^n - 2^r$ zeros. The zero function has one coefficient 2^n , at frequency zero, and all other coefficients zero. These identities prove every numerical spectrum in the next section.

5 Exact counts and sharpness

Apply Sections 3 and 4 with $n=12$. Each effective mask selects a quadratic component of binary polar rank zero, six or twelve. The full spectrum of each component is as follows. Multiplicities in the last column count the 4096 input frequencies.

Effective mask	Count	Rank	Walsh values with multiplicities
0	1	0	0: 4095; 4096: 1
4, 8, 13, 14, 15	5	6	−512: 28; 0: 4032; 512: 36
All others	250	12	−64: 2016; 64: 2080

The four zero output coordinates make each effective mask occur for 16 full masks. Thus $250 \cdot 16 = 4000$ nonzero masks are bent. The six singular effective masks have $6 \cdot 16 = 96$ preimages, one of which is the zero mask. Excluding that mask gives exactly 95 nonzero nonbent masks.

$$4095 = 4000 + 80 + 15.$$

Here the 80 masks are 16 copies of each of the five rank-six components. The other 15 are nonzero masks supported entirely in the four padded positions, and they select the identically zero component. A nonzero mask selecting the zero function still belongs to N_F .

The known lower bound applies

MNBC denotes attaining the maximum number of bent components, equivalently $|N_F| = 2^{m-n/2} - 1$. For $n=m=12$ this is 63, so the witness is non-MNBC. Deryck [1, Theorem 5.4.15, printed p. 87] proves that a non-MNBC quadratic map satisfies

$$|N_F| \geq 2^{m-n/2} + 2^{m-n/2-1} - 1$$

when n is even and $n \leq 2m-4$.

At $n=m=12$ the hypotheses hold and the right-hand side is $64+32-1=95$. Together with the explicit witness, this establishes sharpness at that parameter pair. The universal inequality is cited; it is not re-proved or formalized in this package.

6 A variant with independent output coordinates

Separate mathematical corollary. The canonical map used for the finite certificate retains its four zero coordinates. A simple variant has twelve linearly independent coordinate functions and the same 4000/95 component count. This corollary is not claimed to be included in the Lean certificate.

Add four independent linear coordinates

Keep the first eight quadratic functions and define

$$G(x) = (F_0(x), \dots, F_7(x), x_0, x_1, x_2, x_3).$$

Write a mask as $b=(\lambda, \rho)$, where $\lambda \in \mathbb{F}_2^8$ and $\rho \in \mathbb{F}_2^4$. Let $L(x)=(x_0, x_1, x_2, x_3)$. The selected component becomes

$$g_b(x) = f_b(x) + \langle \rho, L(x) \rangle, \\ W_G(b, a) = W_F(b, a + L^T \rho).$$

Thus the new Walsh spectrum is only a permutation of the old frequency list. Bentness and every signed spectrum are unchanged for each mask. In particular $|N_G|=95$ and G has 4000 bent nonzero components.

No nonzero mask selects the zero function

If $\lambda \neq 0$, the polar form of g_b is the nonzero form B_λ , since adding a linear function does not change polarization. Hence g_b cannot be identically zero. If $\lambda=0$ and $\rho \neq 0$, the component is a nonzero combination of the four independent input coordinate functions. This proves linear independence of all twelve coordinate functions of G .

The 15 nonzero masks in the former padded directions now select nonzero linear components. Such a component still has a single Walsh coefficient +4096 and 4095 zeros, with the nonzero coefficient moved away from frequency zero. It remains nonbent.

What the variant does and does not establish

The quadratic parts still span an eight-dimensional space, and their polar ranks remain exactly the same. Moreover, adding a linear map changes each derivative by a constant output vector. It therefore preserves derivative fiber sizes and cannot turn this construction into an APN map.

For the canonical F , each derivative $x \mapsto F(x+a) - F(x)$ lands in a 256-element output subspace. Its 4096 inputs force a fiber of size at least 16. For G , the last four derivative coordinates are fixed by a , and the same obstruction remains. Neither variant is presented as a deployment-ready cryptographic primitive.

The bound in [1] does not require independent coordinate functions. This corollary nevertheless shows that the same numerical extremum can be realized without identically zero nonzero-mask components. The finite formalization targets F ; the corollary for G is established by the argument above.

Appendix A All binary coefficients

For $x \in F_2^{12}$, the encoded output is $F(x) = \bigoplus_{i < j, x_i x_j = 1} c_{ij}$. The decimal integer c_{ij} is an eight-bit vector, with bit k the coefficient of $x_i x_j$ in F_k . Its top four output bits are zero. Pairs are listed lexicographically, reading down each of the three blocks.

Each coefficient can also be recovered from the field formula: $(c_{ij})_k$ is the corresponding upper-triangular entry of the binary polar matrix M_{e_k} . Within a single three-bit input block the coefficients are zero.

Pair (i,j)	c_{ij}	Pair (i,j)	c_{ij}	Pair (i,j)	c_{ij}
(0, 1)	0	(2, 4)	3	(4, 11)	130
(0, 2)	0	(2, 5)	48	(5, 6)	0
(0, 3)	3	(2, 6)	16	(5, 7)	72
(0, 4)	0	(2, 7)	129	(5, 8)	64
(0, 5)	48	(2, 8)	144	(5, 9)	16
(0, 6)	1	(2, 9)	0	(5, 10)	130
(0, 7)	128	(2, 10)	196	(5, 11)	144
(0, 8)	16	(2, 11)	192	(6, 7)	0
(0, 9)	4	(3, 4)	0	(6, 8)	0
(0, 10)	192	(3, 5)	0	(6, 9)	3
(0, 11)	0	(3, 6)	8	(6, 10)	0
(1, 2)	0	(3, 7)	64	(6, 11)	32
(1, 3)	0	(3, 8)	0	(7, 8)	0
(1, 4)	48	(3, 9)	2	(7, 9)	0
(1, 5)	3	(3, 10)	128	(7, 10)	32
(1, 6)	128	(3, 11)	16	(7, 11)	3
(1, 7)	16	(4, 5)	0	(8, 9)	32
(1, 8)	129	(4, 6)	64	(8, 10)	3
(1, 9)	192	(4, 7)	0	(8, 11)	32
(1, 10)	0	(4, 8)	72	(9, 10)	0
(1, 11)	196	(4, 9)	128	(9, 11)	0
(2, 3)	48	(4, 10)	16	(10, 11)	0

The complete output truth table uses input integer $\sum_{i=0}^{11} 2^i x_i$. Its SHA-256 below is computed over the 4096 output bytes in that input order, not over the JSON representation.

```
ae7383fd45347cbcf32eefec5a6750434967
dd727fe6751a388f3ba6ff72bce3
```

Appendix B A family in dimensions divisible by four

Uniform theorem. The construction works for every $e \geq 1$ and $n=4e$. The mathematical proof below accompanies a checked Lean existence theorem; Appendix C identifies its exact scope and classical coordinate choices. Novelty remains provisional.

Two anisotropic binary planes

Identify the six-dimensional binary space $Q = \text{Alt}(4, F_2)$ with the upper-entry vectors (a, b, c, d, f, g) in pair order 01, 02, 03, 12, 13, 23. Its Pfaffian quadratic form and polar pairing are

$$\begin{aligned} q(a, b, c, d, f, g) &= ag + bf + cd, \\ \beta(X, Y) &= q(X+Y) + q(X) + q(Y). \end{aligned}$$

Define the four-dimensional space V and two binary planes O, H by

$$\begin{aligned} A(u, v, w, z) &= (u+v, u, w, z, v, u+v), \\ V &= \{A(u, v, w, z)\}, \\ O &= \text{span}\{(1, 1, 0, 0, 1, 0), (1, 0, 0, 0, 0, 1)\}, \\ H &= \text{span}\{(0, 0, 1, 1, 0, 0), (0, 1, 1, 0, 1, 0)\}. \end{aligned}$$

In either ordered basis of O or H , $q(s \cdot v_1 + t \cdot v_2) = s + t + st$. Thus q equals one on each plane's three nonzero vectors. Direct substitution gives $\beta(V, O) = 0$ and $\beta(H, O) = 0$. Also $q(A) = u + v + uv + wz$ has the six zeros already counted in Section 2.

The extension pencil

Let $E = F_{2^e}$ and choose α of degree e , so that $1, \alpha, \dots, \alpha^{e-1}$ is a binary basis. For $e=1$ only 1 is used. Take $X_0 = A \in V$; for $1 \leq j < e$ choose $X_j \in O$ when j is odd and $X_j \in H$ when j is even. Set

$$K = \sum_{j=0}^{e-1} \alpha^j X_j \in \text{Alt}(4, E).$$

Coefficientwise independence makes this a binary pencil of dimension $4 + 2(e-1) = 2e + 2$. Extending q and β to E gives

$$\begin{aligned} q(K) &= \sum_j \alpha^{2j} q(X_j) \\ &+ \sum_{i < j} \alpha^{i+j} \beta(X_i, X_j). \end{aligned}$$

Every cross term with $i+j$ odd vanishes: its two entries lie in O and either V or H . For an even sum, write $i+j=2k$; then $i < k < j$. Thus its basis index k is strictly below its larger active index j .

Suppose $m \geq 1$ is the largest index with $X_m \neq 0$. In the basis $1, \alpha^2, \dots, \alpha^{2(e-1)}$, the coefficient at index m equals $q(X_m) = 1$; no cross term reaches it. These squared powers are independent because a relation between them is the square of the corresponding relation between $1, \alpha, \dots, \alpha^{e-1}$, and Frobenius is injective. Therefore $q(K) \neq 0$.

Conversely, if all $X_j = 0$ for $j \geq 1$, then $q(K) = q(A)$. Hence the entire pencil has exactly six singular matrices, regardless of e .

Appendix B Trace and the general count

Trace nondegeneracy in every extension degree

For $E = \mathbb{F}_{2^e}$, write $T(z) = z + z^2 + \dots + z^{2^{e-1}}$. Its values are binary because $T(z)^2 = T(z)$. As a polynomial, T is nonzero and has degree $2^{e-1} < 2^e$. A nonzero polynomial over a field has at most its degree many roots, so T is not identically zero on E . Therefore some $z \in E$ satisfies $T(z) = 1$.

Given $c \neq 0$, take $d = z/c$. Then $T(cd) = 1$, proving the trace pairing is nondegenerate. This argument is necessary for even e , when $T(1) = 0$ and the shortcut used specifically for $\mathbb{F}_8$ would fail.

Exactly as in Section 3, the radical of the binary form $(x, y) \mapsto T(x^T K y)$ is the E -kernel of K viewed over $\mathbb{F}_2$. Thus its binary rank equals e times the E -rank. The pencil therefore has one form of rank zero, five of rank $2e$, and $2^{2e+2} - 6$ of rank $4e$. Trace preserves the pencil's binary dimension $2e + 2$.

The associated quadratic map

Choose the induced binary basis of E^4 . For the $2e + 2$ basis matrices M_k of the trace pencil, take coordinate functions $f_k(x) = \sum_{a < b} (M_k)_{ab} x_a x_b$. Append $2e - 2$ zero coordinates. This gives $F: \mathbb{F}_2^{4e} \rightarrow \mathbb{F}_2^{4e}$, with exactly six nonbent effective masks, including zero.

$$\begin{aligned} |N_F| &= 6 \cdot 2^{2e-2} - 1 \\ &= 3 \cdot 2^{2e-1} - 1 = 3 \cdot 2^{n/2-1} - 1. \end{aligned}$$

The proof of the bent criterion in Section 4 applies unchanged. The resulting count is larger than the MNBC count $2^{2e} - 1$. The cited general lower bound [1] applies for every $e \geq 1$, so this is equality for every positive input dimension divisible by four. No assertion about $n \equiv 2 \pmod 4$ follows.

Finite cross checks

The included `family/verify_family.py` checks every parameter over explicit fields for $e = 1, \dots, 6$. It validates each field by inverses, and finds exactly the same six singular labels in each case. These finite checks illustrate the proof; they do not replace its argument for all e .

e	Input bits n	Effective parameters	Nonzero nonbent masks
1	4	16	5
2	8	64	23
3	12	256	95
4	16	1024	383
5	20	4096	1535
6	24	16384	6143

At $e = 3$, use $\alpha^3 + \alpha + 1 = 0$ and the displayed O and H bases. Then $K = A + \alpha X_1 + \alpha^2 X_2$ is exactly the twelve-bit pencil of Section 1. The $e = 1$ and $e = 2$ counts already have prior constructions; this appendix does not claim them as newly discovered.

Appendix C Verification and trust boundary

Human mathematical proof

Sections 1 through 5 and Appendix B give the algebraic proofs, including trace nondegeneracy and the Walsh criterion. Section 6 gives the linear-coordinate variant. These arguments can be checked without accepting generated certificates or Python results.

Exhaustive executable checks

The dependency-free script `verify_all.py` regenerated the 66 binary coefficients from the six field formulas and checked all 256 field and binary ranks. It verified injectivity and the Pfaffian identity at every parameter. It also compared the binary and trace formulas at all 4096 inputs.

For each of the 256 effective masks, it computed all 4096 Walsh coefficients using exact-integer fast Walsh transforms: 1,048,576 coefficients in total. It checked every full signed spectrum and separately recomputed 1536 raw Walsh sums, at six frequencies per mask. The spectra and truth table agree with a separately implemented NumPy computation supplied in `independent/`.

These scripts are reproducibility checks. A correct execution depends on the interpreter, implementation and assertions; it is not by itself a proof-assistant certificate.

The explicit twelve bit Lean certificate

The canonical witness is the binary polynomial defined by the 66 coefficient masks in Appendix A. The theorem

`N12.exists_exactly_quadratic_with_ninety_five_nonbent_components` proves a valid homogeneous squarefree quadratic representation, nonaffineness, and exact membership of the nonzero nonbent masks in a duplicate-free list of length 95. The separate `exact_nonbent_masks` theorem identifies that list for the concrete F .

Bentness is the original integer Walsh sum over all 4096 inputs, universally quantified over all 4096 frequencies. It is not a surrogate matrix-rank predicate. A proved quadratic translation and Walsh-shift identity reduces the good-component certificates to inverse-alignment checks and one exact zero-frequency sum. Lean checks those data, the bad-component contradictions, output padding and the final count.

An isolated source-only replay freshly compiled all 930 local modules, including all 256 effective component certificates, with Lean 4.19.0 and its bundled Std library. Every module succeeded; no local object files or successful-build cache were copied into the initial replay tree. The final theorem uses only `propext` and `Quot.sound`. No custom axiom, admitted proof, `native_decide` shortcut or external oracle is used.

The number 4000 is immediate complement arithmetic from the exact 95-mask classification. It is not a separately exported bent-cardinality theorem. The finite certificate uses the explicit binary polynomial without requiring an abstract F_8 construction. Its equivalence to the displayed trace formula is proved mathematically and independently checked on all 4096 inputs.

```
N12.exists_exactly_quadratic_with_ninety_five_nonbent_components
N12.exact_nonbent_masks
```

Appendix C The uniform formal theorem

The strongest uniform result is `FamilyAlgebra.uniform_nonaffine_quadratic`, in `UniformDegree.lean`. For each natural number e greater than zero it produces an actual map from `BitVec (4e)` to `BitVec (4e)`, a squarefree homogeneous quadratic representation with every coefficient index satisfying $i < j < 4e$, explicit nonaffineness, and the exact actual-Walsh nonzero nonbent component count $3 * 2^{(2e-1)} - 1$.

Its only hypothesis is $e > 0$. The development proves or instantiates checked mathlib results for field existence, power bases, the Pfaffian zero locus, trace nondegeneracy, coordinate transport, parameter linearity, correct polarization, actual Walsh bentness, and exact counting. No bridge in this chain is assumed as an unproved premise. The separate `uniform_quadratic_family` theorem also classifies every mask.

An isolated source-only replay rebuilt all 26 local modules with Lean 4.19.0 and mathlib commit `c44e0c8ee63ca166450922a373c7409c5d26b00b`. Every module exited successfully and matched its frozen source hash. The final theorem uses only `propext`, `Classical.choice` and `Quot.sound`. There are no admitted proofs, custom axioms, `native_decide` shortcuts or unsafe proof replacements.

The formal degree claim is a valid squarefree degree-at-most-two representation together with a proof that the function is not affine. This is mathematically exact degree two; no separate `algebraicDegree` API is asserted.

The proof uses classical choices of finite fields, generators, bases and coordinate equivalences. It is an existence theorem for every e , not an efficient coefficient-extraction algorithm. Its $e=3$ witness is not identified byte-for-byte with the separately certified explicit twelve-bit map.

```
FamilyAlgebra.uniform_nonaffine_quadratic
FamilyAlgebra.uniform_quadratic_family
FamilyAlgebra.uniform_twelve_count
```

Independent boundaries

Formal verification cannot prove that a construction is historically new. Conversely, a novelty search cannot certify the mathematics. The cited general lower bound is an external mathematical dependency for the word “extremal,” separate from both the concrete and uniform existence/count theorems. The optional linear-coordinate corollary and the exact signed spectral multiplicities are proved mathematically in this note; they are not separate exported Lean claims.

The requested project label “GPT-6 Astra” identifies this research deliverable. No claim about the underlying model runtime, human authorship, peer review, institutional endorsement or publication priority is made.

Appendix D Reproducing and citing the result

Minimal finite check

Use Python 3.10 or later. From the source directory, run:

```
python3 verify_all.py
```

No third-party package or network access is needed. The script fails on any inconsistent coefficient, rank, input value or Walsh spectrum; it writes results/reproduced.json. Expected final counts are 4000 bent and 95 nonbent nonzero masks. The supplied run log records a successful execution. For a separate NumPy implementation, consult independent/verify.py; its original relative paths are preserved as provenance, while verify_all.py is the portable entry point.

To rebuild this PDF, install ReportLab in an isolated Python environment and run:

```
python3 build_report.py
```

Bundled fonts and requirements-pdf.txt record the rendering dependencies. Check SHA256SUMS before comparing files. The truth-table byte hash in Appendix A deliberately differs from the hash of its JSON file.

The two Lean projects

The concrete project in lean/concrete needs Lean 4.19.0 and bundled Std only. From that project directory, the following command freshly rebuilds all 930 local modules. The packaged independent-clean-build.log and logs/Main.log record a successful source-only run with the final axiom output.

Starting from the source directory:

```
cd lean/concrete
LEAN="$(command -v lean)" python3 build.py --clean
```

The uniform project pins Lean 4.19.0 and the exact mathlib commit. The commands below also passed from an extracted source archive with freshly downloaded official dependencies and isolated home/cache/git settings. All 26 local modules were rebuilt without old project or mathlib paths; only the official Lean toolchain and Std were reused. The first download run timed out; its resumed run and both subsequent commands succeeded. Evidence is included in lean/uniform/bootstrap/.

Starting again from the source directory:

```
cd lean/uniform
lake update
lake exe cache get
lake env python3 replay.py --clean
```

Appendix E The established construction mechanism

The pencil of Appendix B is explicitly congruent to a hyperbolic binary plane plus a Jha-Johnson cyclic-semifield core. Thus its underlying construction is established. The exact extremal component-count consequence was not located as a published statement; its novelty remains unestablished. The correspondence and elementary corollary are given here so this distinction is auditable.

An explicit change of basis

Use the six-bit alternating-matrix encoding in Appendix B. Put $D = \text{span}\{47, 61\}$; direct calculation gives $V = D \oplus H$. The invertible binary matrix below acts by $M \mapsto P^T M P$:

$$P = [(1, 1, 1, 0); (0, 1, 1, 1); \\ (1, 0, 1, 0); (0, 1, 0, 1)].$$

Its columns have masks 5, 11, 7, 10. The images of the ordered bases are $D: (47, 61) \mapsto (1, 32)$, $H: (12, 22) \mapsto (18, 28)$, and $O: (19, 33) \mapsto (22, 26)$. The latter two pairs occupy the cross block between the first and last two coordinates. In that block they are (I, R) and (J, JR) , where

$$R = [(0, 1); (1, 1)], J = [(1, 1); (0, 1)], \\ R^2 + R + I = 0, J^2 = I, JR = R^2 J.$$

Write $K = \text{span}_{F_2}\{I, R\}$, a matrix field isomorphic to F_4 , with conjugation $C \mapsto C = C^2$. The transformed base parameters are $x = u + w + z$, $y = v + w + z$, $h_0 = u + v + z$, $h_1 = w + z$. This binary change is invertible. The full pencil consists of

$$M(x, y, B) = [(xZ, B); (B^T, yZ)], \\ Z = [(0, 1); (1, 0)], x, y \in F_2, \\ B \in S = \sum_{0 \leq j < e} T^j K, T = \alpha J.$$

Here $B = h_0 I + h_1 R + \sum_{j=1}^{e-1} \alpha^j J^{j \bmod 2} (s_j I + t_j R)$. Thus this is equality of pencils after a fixed congruence, rather than a numerical resemblance.

Why this is a cyclic semifield core

On E^2 , the matrix field K gives an e -dimensional K -vector space. T is semilinear for the nontrivial automorphism of K and $T^2 = \gamma I$, with $\gamma = \alpha^2$. For $e > 1$, γ generates E over F_2 . The algebra generated by K and T contains E and J and hence all $\text{Mat}_2(E)$, since I, R, J, JR are E -linearly independent. No proper nonzero K -subspace can be T -invariant. For $e = 1$, take $\alpha = \gamma = 1$; irreducibility is automatic in K -dimension one.

This is precisely the irreducible-semilinear construction recalled by Kantor and Liebler [5, §2, p. 334]. Its odd- e norm and power-basis presentation is also given by Johnson, Marino, Polverino and Trombetti [6, Theorem 1(2), pp. 10–11]. The general argument above includes even e ; [6] is cited only for its odd- e specialization.

Appendix E The determinant fibre corollary

The following elementary refinement counts the determinant-one fibre of the established core S. It explains how the six singular forms, and hence the extremal component count, follow from that mechanism. This correspondence is a mathematical identification; the frozen Lean proof uses Appendix B directly and does not depend on a new citation axiom.

A polynomial over the binary field

Separate the even and odd powers in B. With $\gamma = \alpha^2$, write

$$\begin{aligned} B &= h(\gamma) + T o(\gamma), \\ h(X) &= \sum_{2j < e} C_{2j} X^j, \\ o(X) &= \sum_{2j+1 < e} C_{2j+1} X^j, \quad C_j \in K. \end{aligned}$$

For a polynomial with coefficients in K, a bar means coefficientwise conjugation $C \mapsto C^2$; the variable X is fixed. Direct two by two determinant expansion, using $JR = R^2J$, gives

$$\begin{aligned} \det B &= f(\gamma), \\ f(X) &= h(X)\bar{h}(X) + X o(X)\bar{o}(X) \in F_2[X]. \end{aligned}$$

The two products are fixed by conjugation. If h is nonzero, its product with $\bar{h}$ has degree $2 \deg h$, with leading coefficient one, since each nonzero element of K has norm one to F_2 . If o is nonzero, $Xo\bar{o}$ has odd degree $2 \deg o + 1$, also with leading coefficient one. These leading degrees have different parity and cannot cancel. Both are at most $e-1$.

Since γ has degree e, evaluation is injective on binary polynomials of degree less than e. Therefore $\det B = 0$ forces $h = o = 0$. Likewise $\det B = 1$ forces $o = 0$ and h a nonzero constant. Exactly three constants in K are nonzero, so

$$\begin{aligned} \#\{B \in S : \det B = 0\} &= 1, \\ \#\{B \in S : \det B = 1\} &= 3. \end{aligned}$$

For $e=1$ the same assertion follows directly from $S=K$. For even e the expression $h\bar{h} + Xo\bar{o}$ is a formal coefficientwise identity; it is not an assertion that E adjoined with K is a quadratic field extension. This avoids an invalid field-norm interpretation in that case.

Adjoin the two hyperbolic directions

The block matrix from the previous page has Pfaffian $xy + \det B$. If $B=0$, precisely the three pairs with $xy=0$ make it vanish. If $\det B=1$, each of the three possible B has the single pair $x=y=1$. There are no other solutions because xy is binary. Thus there are exactly six singular matrices, including zero.

$$6 \cdot 2^{2e-2} - 1 = 3 \cdot 2^{2e-1} - 1.$$

Trace descent, the quadratic bent criterion and output padding give the displayed count, as in Appendix B. The count is therefore a short determinant-fibre corollary of an established core. It should not be advertised as a new underlying family. The package includes family/CYCLIC_SEMIFIELD_CORRESPONDENCE.txt and a dependency-free checker for the congruence and finite determinant fibres.

Appendix E Literature and sources

Deryck's printed p. 88 asks for higher-dimensional equality [1]. The mathematical and Lean results here attain the bound for equal dimensions divisible by four. The explicit correspondence above identifies the family mechanism with established cyclic-semifield machinery [5,6]. A targeted search located no prior publication of this exact extremal component-count consequence. That limited negative finding does not establish priority; specialist review remains necessary. Any potentially new contribution is the application or corollary and its formal verification, not the underlying construction.

Trace descent and rank multiplication are established [2]. The broader geometric correspondence with additive semifield spread sets also appears in [4, Theorem 16]. Eight-bit numerical existence follows already from [3, Definition 4.1 and Table 4.1]: a final profile pair (58,78) or (58,90) gives a six-dimensional component space with 58 bent functions. Padding by two output zeros yields 232 bent and 23 nonbent nonzero masks. No equivalence of those examples to this formula is asserted.

References

- [1] Maxime Deryck. *Cryptography meets Finite Geometry*. Master's thesis, Ghent University, academic year 2025–2026. Theorem 5.4.15, printed p. 87; higher-dimensional discussion, printed p. 88. [Author-hosted thesis PDF](#).
- [2] Rod Gow. *Rank-related dimension bounds for subspaces of bilinear forms over finite fields*. arXiv:1703.07266v1, 21 March 2017. See the proof of Theorem 6 for trace descent. [arXiv record](#); [full text](#).
- [3] Christof Beierle, Philippe Langevin, Gregor Leander, Alexandr Polujan and Shahram Rasoolzadeh. *Millions of inequivalent quadratic APN functions in eight variables*. arXiv:2508.04644v1, 6 August 2025. Definition 4.1 and Table 4.1. [Full text](#).
- [4] Guglielmo Lunardon. *50 Years of translation structures*. Journal of Geometry 113, article 35 (2022). Section 3.3.3 and Theorem 16. DOI: 10.1007/s00022-022-00643-5. [Publisher full text](#).
- [5] William M. Kantor and Robert A. Liebler. *Semifields arising from irreducible semilinear transformations*. Journal of the Australian Mathematical Society 85 (2008), 333–339. Section 2, printed p. 334. DOI: 10.1017/S1446788708000888. [Author-hosted PDF](#).
- [6] Norman L. Johnson, Giuseppe Marino, Olga Polverino and Rocco Trombetti. *On a generalization of cyclic semifields*. Journal of Algebraic Combinatorics 29 (2009), 1–34. Theorem 1(2), pp. 10–11, for odd extension degree. DOI: 10.1007/s10801-007-0116-x. [Full text PDF](#).

External sources were checked on 9 October 2026. Only the general bound and historical context rely on these citations; the finite construction and its proof are explicit in this note.