

1 Lecture 3: January 27, 2026

Lecture Overview: We develop the set-theoretic foundations needed for analysis. We define **functions** as special relations and introduce key properties: surjectivity, injectivity, and bijectivity. Using bijections, we define when two sets have the same **cardinality**, leading to the notions of **finite**, **infinite**, and **countable** sets.

1.1 Functions

Section Overview: We define functions as relations with a uniqueness property, introduce images and preimages, and classify functions as injective (one-to-one), surjective (onto), or bijective (both).

Definition 1.1. A **function** $f : A \rightarrow B$ is a relation $R \subseteq A \times B$ such that for all $x \in A$, there exists a unique $y \in B$ with $(x, y) \in R$. This y is denoted $f(x)$. The set A is called the **domain** of f , and B is the **codomain** (or **range**).

Remark 1.2. Not all relations are functions. A relation fails to be a function if some element of A maps to multiple elements of B , or to no element at all.

Definition 1.3. Let $f : A \rightarrow B$ be a function.

- If $E \subseteq A$, the **image** of E under f is $f(E) = \{f(x) : x \in E\}$.
- If $F \subseteq B$, the **preimage** of F under f is $f^{-1}(F) = \{x \in A : f(x) \in F\}$.

Definition 1.4. Let $f : A \rightarrow B$ be a function.

- f is **surjective** (or **onto**) if $f(A) = B$, i.e., for every $y \in B$, there exists $x \in A$ with $f(x) = y$.
- f is **injective** (or **one-to-one**) if $f(x_1) = f(x_2)$ implies $x_1 = x_2$, i.e., distinct inputs map to distinct outputs.
- f is **bijective** if it is both injective and surjective.

1.2 Equivalence Relations

Section Overview: We review equivalence relations and observe how function properties relate to composition and inverses.

Remark 1.5. Recall from Lecture 1 that an equivalence relation satisfies:

- **Reflexivity:** $x \sim x$ (like the identity function)
- **Symmetry:** $x \sim y \Rightarrow y \sim x$ (like invertible functions: if $f : A \rightarrow B$, then $f^{-1} : B \rightarrow A$)
- **Transitivity:** $x \sim y$ and $y \sim z \Rightarrow x \sim z$ (like composition: $f : A \rightarrow B$ and $g : B \rightarrow C$ give $g \circ f : A \rightarrow C$)

1.3 Cardinality

Section Overview: We define cardinality using bijections, then classify sets as finite, infinite, or countable.

Definition 1.6. Two sets A and B have the same **cardinality**, written $A \sim B$, if there exists a bijection $f : A \rightarrow B$.

Definition 1.7. Let $[n] = \{1, 2, 3, \dots, n\}$. We say that a set A is:

1. **finite** if there exists $n \in \mathbb{N}$ such that $A \sim [n]$,
2. **infinite** if A is not finite,
3. **countable** if $A \sim \mathbb{N}$.

Example. \mathbb{Z} is countable. We can list the integers as:

$$0, 1, -1, 2, -2, 3, -3, \dots$$

This defines a bijection $f : \mathbb{N} \rightarrow \mathbb{Z}$ given by

$$f(n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ -\frac{n-1}{2} & \text{if } n \text{ is odd} \end{cases}$$

(where we start \mathbb{N} at 0). Thus $\mathbb{Z} \sim \mathbb{N}$, so \mathbb{Z} is countable.

Definition 1.8. A **sequence** is a function $f : \mathbb{N} \rightarrow X$ where X is some set. If we denote $f(n)$ by x_n , then we write the sequence as $(x_n)_{n=1}^{\infty}$.

Remark 1.9. If A is a countable set, then there exists a surjection $f : \mathbb{N} \rightarrow A$. We say that A can be **arranged in a sequence**: the elements of A can be listed as $f(1), f(2), f(3), \dots$

Note: The **set difference** $A \setminus B$ (read “ A minus B ”) is defined as

$$A \setminus B = \{x \in A : x \notin B\}.$$

Theorem 1.10. *Every infinite subset of a countable set is countable.*

Proof. Let A be a countable set and let $B \subseteq A$ be infinite. Since A is countable, we can arrange its elements as a sequence $(x_n)_{n=1}^{\infty}$.

We construct a bijection $f : \mathbb{N} \rightarrow B$ inductively. Let $B_0 = \emptyset$. For each $k \geq 1$:

- Let $n_k = \min\{n \in \mathbb{N} : x_n \in B \setminus B_{k-1}\}$
- Define $f(k) = x_{n_k}$
- Set $B_k = B_{k-1} \cup \{x_{n_k}\}$


At each step, x_{n_k} is assigned position $k = |B_{k-1}| + 1$ in our enumeration of B . Since B is infinite, we never exhaust it, so each n_k exists.

This f is a bijection: it is injective since each element is added exactly once, and surjective since every element of B appears at some position n in the sequence (x_n) and will eventually be enumerated.


Thus $B \sim \mathbb{N}$, so B is countable. □

1.4 Unions and Intersections of Sets

Section Overview: We define unions and intersections of collections of sets, including arbitrary (possibly infinite) unions and intersections.



$A \cup B$ (union)



$A \cap B$ (intersection)

Definition 1.11. Let A and B be sets, and let $f : A \rightarrow \mathcal{P}(B)$ be a function assigning to each $\alpha \in A$ a subset $f(\alpha) \subseteq B$. We define:

- The **union** of the family is

$$\bigcup_{\alpha \in A} f(\alpha) = \{x \in B : x \in f(\alpha) \text{ for some } \alpha \in A\}.$$

- The **intersection** of the family is

$$\bigcap_{\alpha \in A} f(\alpha) = \{x \in B : x \in f(\alpha) \text{ for all } \alpha \in A\}.$$

Theorem 1.12 (De Morgan's Laws). *Let A and B be subsets of a universal set U . Then:*

1. $(A \cup B)^c = A^c \cap B^c$
2. $(A \cap B)^c = A^c \cup B^c$

Proof. We prove (1); the proof of (2) is similar.

(\subseteq) Let $x \in (A \cup B)^c$. Then $x \notin A \cup B$, so $x \notin A$ and $x \notin B$. Thus $x \in A^c$ and $x \in B^c$, so $x \in A^c \cap B^c$.

(\supseteq) Let $x \in A^c \cap B^c$. Then $x \in A^c$ and $x \in B^c$, so $x \notin A$ and $x \notin B$. Thus $x \notin A \cup B$, so $x \in (A \cup B)^c$. \square

Note: To prove two sets are equal, $X = Y$, a standard technique is the **mutual subset argument**: show $X \subseteq Y$ and $Y \subseteq X$. For each direction, take an arbitrary element of one set and show it belongs to the other.

Theorem 1.13 (Distributive Law). *Let A , B , and C be sets. Then*

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$$

Proof. (\subseteq) Let $x \in A \cap (B \cup C)$. Then $x \in A$ and $x \in B \cup C$. Since $x \in B \cup C$, either $x \in B$ or $x \in C$.

- If $x \in B$, then $x \in A \cap B$, so $x \in (A \cap B) \cup (A \cap C)$.

- If $x \in C$, then $x \in A \cap C$, so $x \in (A \cap B) \cup (A \cap C)$.
- (\supseteq) Let $x \in (A \cap B) \cup (A \cap C)$. Then $x \in A \cap B$ or $x \in A \cap C$.
- If $x \in A \cap B$, then $x \in A$ and $x \in B \subseteq B \cup C$, so $x \in A \cap (B \cup C)$.
- If $x \in A \cap C$, then $x \in A$ and $x \in C \subseteq B \cup C$, so $x \in A \cap (B \cup C)$.

□

1.5 The (Un)countability of Number Systems

Section Overview: We apply our results to the classical number systems. We show \mathbb{Q} is countable (as a countable union of countable sets), but \mathbb{R} is **uncountable** using Cantor's diagonal argument on binary sequences. This reveals a hierarchy of infinities: $|\mathbb{N}| = |\mathbb{Q}| < |\mathbb{R}|$.

Theorem 1.14. Let $\{E_n\}_{n=1}^{\infty}$ be a sequence of countable sets. Then

$$\bigcup_{n=1}^{\infty} E_n$$

is countable.

Proof. Since each E_n is countable, we can enumerate its elements as

$$E_n = \{x_{n,1}, x_{n,2}, x_{n,3}, \dots\}.$$

Arrange all elements in an infinite grid:

| | | | | | |
|--------|-----------|-----------|-----------|-----------|---------|
| | 1 | 2 | 4 | 7 | |
| $E_1:$ | $x_{1,1}$ | $x_{1,2}$ | $x_{1,3}$ | $x_{1,4}$ | \dots |
| | 3 | 5 | 8 | | |
| $E_2:$ | $x_{2,1}$ | $x_{2,2}$ | $x_{2,3}$ | $x_{2,4}$ | \dots |
| | 6 | 9 | | | |
| $E_3:$ | $x_{3,1}$ | $x_{3,2}$ | $x_{3,3}$ | $x_{3,4}$ | \dots |
| | 10 | | | | |
| $E_4:$ | $x_{4,1}$ | $x_{4,2}$ | $x_{4,3}$ | $x_{4,4}$ | \dots |
| | \vdots | \vdots | \vdots | \vdots | |

We enumerate the union by traversing diagonals: first $x_{1,1}$, then $x_{1,2}, x_{2,1}$, then $x_{1,3}, x_{2,2}, x_{3,1}$, and so on. The k -th diagonal contains all $x_{n,m}$ with $n + m = k + 1$.

This gives a surjection from \mathbb{N} onto $\bigcup_{n=1}^{\infty} E_n$ (skipping repeats if sets overlap). Thus the union is countable. □

Note: The **diagonal argument** is a powerful technique for enumerating countable unions. By arranging elements in a grid and traversing along diagonals, we reduce a “two-dimensional” infinite collection to a “one-dimensional” sequence.

Theorem 1.15. *If A is countable, then A^n is countable for all $n \in \mathbb{N}$.*

Proof. By induction on n .

Base case ($n = 1$): $A^1 = A$ is countable by assumption.

Inductive hypothesis: Assume A^n is countable for some $n \geq 1$.

Inductive step: We show A^{n+1} is countable. Observe that

$$A^{n+1} = A^n \times A.$$

By the inductive hypothesis, A^n is countable, so we can enumerate it as $A^n = \{b_1, b_2, b_3, \dots\}$. Since A is countable by assumption, we can write $A = \{a_1, a_2, a_3, \dots\}$. The Cartesian product $A^n \times A$ can then be arranged in a grid:

$$\begin{array}{cccc} (b_1, a_1) & (b_1, a_2) & (b_1, a_3) & \cdots \\ (b_2, a_1) & (b_2, a_2) & (b_2, a_3) & \cdots \\ (b_3, a_1) & (b_3, a_2) & (b_3, a_3) & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array}$$

By the diagonal argument, $A^n \times A$ is countable.

Conclusion: By the principle of mathematical induction, A^n is countable for all $n \in \mathbb{N}$. □

Corollary 1.16. \mathbb{Q} is countable.

Proof. For each $n \in \mathbb{N}$, let $E_n = \{\frac{m}{n} : m \in \mathbb{Z}\}$ be the set of rationals with denominator n . Each E_n is countable (since $E_n \sim \mathbb{Z}$ and \mathbb{Z} is countable). Then

$$\mathbb{Q} = \bigcup_{n=1}^{\infty} E_n$$

is a countable union of countable sets, hence countable. □

Theorem 1.17 (\mathbb{R} is uncountable). *The set of binary sequences $\{0, 1\}^{\mathbb{N}}$ is uncountable.*

Proof. By contradiction. Suppose $\{0, 1\}^{\mathbb{N}}$ is countable. Then we can list all binary sequences as s_1, s_2, s_3, \dots where each $s_n = (s_{n,1}, s_{n,2}, s_{n,3}, \dots)$:

| | pos 1 | pos 2 | pos 3 | pos 4 | pos 5 | |
|----------|----------|----------|----------|----------|----------|----------|
| s_1 : | 0 | 1 | 0 | 1 | 1 | \cdots |
| s_2 : | 1 | 1 | 0 | 0 | 1 | \cdots |
| s_3 : | 0 | 0 | 1 | 1 | 0 | \cdots |
| s_4 : | 1 | 0 | 1 | 0 | 0 | \cdots |
| s_5 : | 0 | 1 | 1 | 1 | 1 | \cdots |
| \vdots | \vdots | \vdots | \vdots | \vdots | \vdots | |
| d : | 1 | 0 | 0 | 1 | 0 | \cdots |

flip diagonal

Construct a new sequence $d = (d_1, d_2, d_3, \dots)$ by flipping the diagonal entries:

$$d_n = \begin{cases} 1 & \text{if } s_{n,n} = 0 \\ 0 & \text{if } s_{n,n} = 1 \end{cases}$$

Then d differs from s_n in the n -th position for every $n \in \mathbb{N}$. Thus $d \neq s_n$ for all n , so d is not in our list. But $d \in \{0, 1\}^{\mathbb{N}}$, contradicting that our list contains all binary sequences.

Therefore $\{0, 1\}^{\mathbb{N}}$ is uncountable. □

Note: This is **Cantor's diagonal argument**. Since there is a bijection between $\{0, 1\}^{\mathbb{N}}$ and \mathbb{R} (via binary expansions), this proves \mathbb{R} is uncountable. The key insight is that any proposed enumeration can be “diagonalized” to produce a missing element.