TitrateRL: A Simulation-Based Reinforcement Learning Approach for Personalized Basal Insulin Titration in Type 2 Diabetes

Boburmirzo Uchqunov (240447), Nodirjon Khabibullaev (230019),
Ibrohim Abralov (230129), and Abdurashid Abdusattarov (221229)
School of Engineering, BSc in Computer Science
Central Asian University, Tashkent, Uzbekistan

***Abstract*—TitrateRL proposes an open, simulation-based research framework for personalized basal-insulin titration in Type 2 diabetes (T2D). A published T2D glucose–insulin model is extended with long-acting glargine absorption, meal variability and sensor noise, calibrated against a landmark titration trial, and exposed as a Gymnasium environment with 150 heterogeneous virtual patients. Proximal Policy Optimization (PPO) agents with a rule-based safety shield will be compared with three clinical titration algorithms over 24-week simulated courses. The implemented simulator passes integrator, equilibrium, cross-implementation and glargine-profile checks, and one 168-day episode runs in about 0.15 s; calibration against the Treat-to-Target trial is still in progress. Evaluation of reward design, comparison with clinical algorithms and analysis of personalization on held-out virtual patients is planned. All results are obtained in silico and are not intended for clinical use.**

***Index Terms*—reinforcement learning, Type 2 diabetes, basal insulin titration, in silico simulation, safe reinforcement learning, Gymnasium**

I. Introduction

Type 2 diabetes (T2D) affects most of the 589 million adults living with diabetes [1]. When non-insulin agents no longer reach the glycemic target, a once-daily basal insulin such as glargine is added and its dose is titrated, for example by +2 U every three days, until the fasting plasma glucose (FPG) reaches 80–130 mg/dL [2], [3]. Clinical titration algorithms apply the same thresholds to every patient, and in routine care up-titration is slow: 81% of patients remained above their HbA1c target six months after starting basal insulin [4]. This project investigates whether a reinforcement learning (RL) policy, trained entirely in simulation, can titrate basal insulin more effectively than clinical rules without increasing hypoglycemia, and builds the open infrastructure needed to answer this question reproducibly.

A. Background and Related Work

The treat-to-target, AT.LANTUS and INSIGHT trials established weekly, three-day and daily fasting-glucose titration rules [5]–[7]. The meal simulation model of Dalla Man *et al.* [8] is the basis of the UVA/Padova simulator, accepted for preclinical testing of insulin-dosing algorithms [9]; its open re-implementation `simglucose` contains only Type 1 patients [10]. Bonet *et al.* showed in silico that CGM-based rules reach the optimal T2D dose faster than guideline titration [11]. RL improves glucose control in Type 1 simulations but can fail catastrophically when applied naively [12]; 24 of 29 RL glucose-control studies address Type 1 diabetes [13]. RL for T2D has used non-public hospital records [14] or a commercial simulator [15], and a recent open safe-RL testbed keeps the basal dose fixed [16]. Table I summarizes the design choices these studies motivate.

B. Problem Statements

P1: Formulation. Daily T2D basal titration has not been posed as an RL problem in an open environment, and it is unclear how clinical objectives with strongly asymmetric risks should be encoded in a scalar reward.

P2: Performance. Clinical algorithms are population-level and use only fasting glucose. Whether a policy learned in simulation keeps glucose in range more effectively without increasing hypoglycemia is unknown for T2D.

P3: Heterogeneity. Insulin sensitivity varies widely between patients and is not observed directly [17]. How the benefit and the dosing behavior of a learned policy depend on insulin sensitivity has not been characterized.

C. Research Questions

1) **RQ1.** How does the reward formulation (R1 time in range, R2 risk index, R3 time in range with hypoglycemia penalties) affect time in range (TIR) and time below range (TBR) of PPO titration policies?
2) **RQ2.** Does a shielded PPO policy achieve a higher maintenance-phase TIR than the AT.LANTUS, Treat-to-Target and INSIGHT algorithms on held-out patients while remaining non-inferior in $\mathrm{TBR}_{<70}$?
3) **RQ3.** How does the benefit vary with insulin sensitivity, and does the policy choose different dose changes for insulin-resistant and insulin-sensitive patients with similar fasting glucose?

D. Research Objectives

General objective. Design, implement and evaluate within 14 weeks an open-source, simulation-based RL framework that learns daily basal-insulin titration policies for virtual patients

TABLE I
RELATED STUDIES AND IMPLICATIONS FOR TITRATERL

Study	Method or setting	Implication for TitrateRL
Dalla Man *et al.* [8]	Meal simulation model with published T2D parameters	Implement the T2D model openly in Python and add subcutaneous basal insulin.
Kovatchev *et al.* [9]	In silico preclinical trials (UVA/Padova, T1D)	Verify and calibrate the simulator before any policy is trained.
Xie, `simglucose` [10]	Open Python T1D simulator with a Gym interface	Reuse the shared subsystems and test against them (cross-implementation).
Riddle, Davies, Gerstein [5]–[7]	Clinical fasting-glucose titration algorithms	Implement them as baselines B1–B3 and calibrate against Treat-to-Target.
Bonet *et al.* [11]	CGM-based titration rules, 300 virtual T2D subjects	Use CGM features and compare titration speed (time-to-target).
Fox *et al.* [12]	Deep RL for closed-loop control in T1D	Use validation-based checkpoint selection and multi-seed reporting.
Wang *et al.* [14]	Model-based RL from hospital records (T2D)	Provide an open benchmark that does not depend on non-public data.
Panagiotou *et al.* [15]	Actor-critic advisor in a commercial T2D simulator	Keep the simulator, cohort and code fully open and reproducible.
Kwon *et al.* [16]	Safe-RL testbed; safety fails on unseen patients	Evaluate on held-out patients and add a rule-based safety shield.

with T2D and benchmarks them against clinical titration algorithms using internationally recommended glycemic metrics.

Specific objectives. Train PPO under R1, R2 and R3 with five seeds and select the reward on validation patients (RQ1). Evaluate the shielded PPO policy against B1–B3 on 30 test patients with ten scenarios each (RQ2). Quantify the association of the benefit with insulin sensitivity and the dose behavior by insulin-sensitivity tertile (RQ3). Deliver the validated environment as an installable package with an evaluation pipeline and an interactive dashboard.

II. METHODOLOGY

A. Development Process and Project Scope

The team follows an iterative, Scrum-inspired process with two-week sprints, a backlog in GitHub Projects and supervisor reviews at milestones M1–M4. The simulator, wrappers and metrics are developed test-first; GitHub Flow with a protected `main` branch, reviewed pull requests and continuous integration (linting and tests on Python 3.10–3.12) separates development from stable revisions. Experiments are version-controlled YAML files with fixed seeds, and models and result tables are stored separately from the source code. The project covers in silico research software only: real patient data, dosing recommendations for real people, Type 1 diabetes, bolus insulin and regulatory validation are outside its scope, and every output is labelled as a simulation result [18].

B. System Architecture and Deployment

TitrateRL follows a layered architecture in which each layer depends only on the layers below it: configuration and data, simulation core, environment, decision, experiment and evaluation, and presentation. Fig. 1 shows where each part runs and how artifacts move between the team's workstations, GitHub (repository, CI and releases), the training machine and the users of the package and the dashboard.

C. Repository Structure and Class Design

The repository (Fig. 2) follows the `src` layout and mirrors the architecture layers. Fig. 3 shows the main classes: `T2DTitrationEnv` composes a `VirtualPatient`, which owns the ODE state and three random streams (meals, CGM, fingerstick) so that all policies are evaluated with common random numbers. Rule-based baselines and RL agents share one abstract `Policy` interface, and `SafetyShield` is a Gymnasium wrapper applied only to RL agents.

D. Simulation Environment and Virtual Cohort

The patient model is the Dalla Man meal model with published T2D parameters [8], [19]. The glucose, insulin and gut subsystems follow `simglucose` [10] and are extended with β-cell secretion, time-varying hepatic extraction and a two-compartment glargine model [20], [21]. The 15-state system is integrated with a 1-min fourth-order Runge–Kutta scheme compiled with Numba [22]. Each day contains three meals with random timing and size; a CGM samples interstitial glucose every 5 min with the Breton–Kovatchev noise model [23], and a fasting fingerstick is taken at 07:00 [24]. The cohort of 150 virtual adults varies pre-insulin FPG, body weight, insulin sensitivity (ρ_{SI}) and β-cell function (ρ_{β}), is calibrated against the Treat-to-Target trial [5], and is split 90/30/30 into training, validation and test patients by ρ_{SI} tertile.

Titration is a partially observable decision process exposed through the Gymnasium API [25]: one step is one day and one episode lasts 168 days. The observation stacks 11 daily features (fasting SMBG, dose per kg, last dose change, CGM TIR, TBR, TAR, mean and variability, recent hypoglycemic events, body weight and time) over three days. The action is a dose change in $\{-4, -2, 0, +2, +4\}$ U; the cumulative increase over any seven days is limited to 8 U and the dose to $[0, 1.0$ U/kg$]$ for every policy. The rewards are R1 $=$ TIR, R2 $=$ the negative mean of the Kovatchev risk function [26], and R3 $=$ TIR $- 5\,\mathrm{TBR}_{<70} - 10\,\mathrm{TBR}_{<54}$.

E. Agents, Safety Shield and Baselines

PPO [27] is the primary algorithm and DQN [28] a secondary one, both from Stable-Baselines3 [29]. Each configuration is trained for 10^6 simulated days with five seeds [30], a new training patient in every episode, and checkpoint selection on validation patients only. A safety shield inspired by shielding in safe RL [31] blocks dose increases after level-1 hypoglycemia in the last 72 h and reduces the dose by 20% after level-2 hypoglycemia in the last 24 h. The baselines are a fixed dose of 0.2 U/kg (B0), AT.LANTUS (B1), Treat-to-Target (B2), INSIGHT (B3) and a simulator-tuned rule (B4).




Fig. 1. Deployment map. Code flows from the four workstations through pull requests and CI to tagged releases; experiments run on a multi-core CPU machine from a pinned commit; trained models and result tables are published as release assets (never committed to git) and consumed by the dashboard and by researchers who install the package.

```
titraterl/
├─ .github/  — CI workflow, PR and issue templates
├─ configs/
│  ├─ cohort.yaml
│  ├─ train/  — ppo_r1/r2/r3, ppo_rstar_shield, dqn, fpg, smoke
│  └─ eval/  — test.yaml, baselines_quick.yaml
├─ src/titraterl/
│  ├─ simulation/  — parameters, model (ODE, RK4), scenario, sensors, patient
│  ├─ cohort/  — generator, split, Parquet I/O
│  ├─ envs/  — titration_env, observation, rewards, wrappers
│  ├─ policies/  — base, rules (B0–B4), sb3
│  ├─ training/  — train, callbacks
│  ├─ metrics/  — glycemic, episode
│  ├─ evaluation/  — runner, statistics, logging, pipeline
│  ├─ validation/  — checks V1–V5
│  └─ config.py, cli.py
├─ dashboard/app.py
├─ tests/
│  ├─ unit/  — model, cross-impl., env, shield, rules, ...
│  └─ integration/  — pipeline, CLI, training, dashboard
├─ docs/  — architecture, model, environment, evaluation, ...
├─ examples/  — quickstart.py, quickstart.ipynb
└─ README.md, MODEL_CARD.md, pyproject.toml, Makefile, LICENSE
```

Fig. 2. Repository file structure (86 tracked files, ≈3 800 lines of package code and ≈800 lines of tests).

F. Preliminary Implementation Evidence

The environment, baselines, training pipeline, evaluation pipeline and dashboard are implemented. Table II records the simulator checks and a preliminary run of the clinical baselines. Calibration (check V4) does not yet meet all trial tolerances: the virtual patients reach a lower fasting glucose than the trial participants at a comparable final dose. Trained-agent results and held-out evaluation remain planned.

G. Evaluation Protocol

Every policy is evaluated on the 30 test patients, each under ten 168-day scenarios with common random numbers. Endpoints are computed from noise-free glucose following the international CGM consensus [32], [33] (Table III). The primary efficacy endpoint is TIR in weeks 21–24 and the primary safety endpoint is $\text{TBR}_{<70}$ over 24 weeks. H2 is tested with Holm-adjusted one-sided paired Wilcoxon tests, the Hodges–Lehmann estimate of the TIR gain ($\geq$5 pp) and 95% bootstrap confidence intervals (non-inferiority margin 1 pp); RQ3 uses Spearman's correlation. Overbasalization [34] and the glucose management indicator [35] are secondary outcomes.

TABLE II
PRELIMINARY SIMULATOR CHECKS AND BASELINE RUN

Check	Result	Criterion
RK4 vs. SciPy integrator	max error 6×10^{-8} mg/dL	<0.5 mg/dL
Right-hand side vs. simglucose	identical (10^{-9})	match
Equilibrium; dose–response	drift 3×10^{-14}; monotone	pass
Glargine flatness, 2–24 h	23.7% from mean	≤25%
Glargine steady state, dose 4	90.3%	≥90%
B2 week-24 FPG (90 patients)	99.6 mg/dL	110–125
B2 final dose	0.57 U/kg	0.4–0.6
B2 patients with FPG ≤100	56%	25–50%
Runtime, 168-day episode	0.15 s	<2 s
Maintenance TIR, B0/B1/B2/B3†	39 / 60 / 63 / 61%	—

† Validation patients, one scenario each; preliminary software check.

H. MVP and Implementation Roadmap

The MVP demonstrates the complete pipeline: virtual-patient generation, training and shielded titration, statistical comparison and interactive visualization. Table IV prioritizes the features with the MoSCoW method, and the team uses Table V to track implementation and evaluation milestones.




Fig. 3. Class diagram of the implemented TitrateRL package (simplified UML). Open triangles: inheritance; filled diamonds: composition; solid arrows: association; dashed arrows: dependency. B1, B3 and B4 inherit from `_IntervalRule`; B2 from `Policy`.

TABLE III
PLANNED MODEL AND SYSTEM EVALUATION METRICS

Metric	Definition	Target / role
Product (acceptance)		
Test suite	unit + integration tests in CI	100% pass
Episode runtime	one 168-day episode	$<$2 s
Dashboard re-run	24-week simulation	$<$5 s
Simulator checks	V1–V4 (gate G1)	all pass by Wk 5
Reproducibility	same seed $\Rightarrow$ same trajectory	exact
Research (glycemic)		
TIR	% time 70–180 mg/dL, weeks 21–24	$>$70%; primary efficacy
$\text{TBR}_{<70}$, $\text{TBR}_{<54}$	% time below 70 / 54 mg/dL	$<$4%, $<$1%; primary safety
$\text{TAR}_{>180}$, $\text{TAR}_{>250}$	% time above 180 / 250	$<$25%, $<$5%
Mean, CV, GMI	mean, SD/mean, $3.31+0.02392\,\bar{G}$	CV $\leq$36%
LBGI, HBGI	Kovatchev risk indices	risk
Time-to-target	3 consecutive fasting values 80–130	titration speed
Nocturnal TBR	00:00–06:00	safety
Overbasalization	bedtime$-$morning $\geq$50; dose $>$0.5 U/kg	appropriateness
Dose burden	final U/kg; changes per week	burden
Shield rate	% days with intervention	safety
Statistical decision rules		
H2 superiority	Holm-adj. one-sided Wilcoxon	$p < 0.05$
H2 effect size	Hodges–Lehmann TIR gain	$\geq$5 pp
H2 non-inferiority	upper 95% CI of $\text{TBR}_{<70}$ diff.	$<$1 pp
H3	Spearman(ρ_{SI}, ΔTIR)	negative, $p < 0.05$

TABLE IV
MVP PRIORITIZATION WITH THE MOSCOW METHOD

Priority	Feature	Acceptance criterion	Status
Must	F1 T2D simulator and calibrated 150-patient cohort	generated reproducibly from one seed; checks V1–V4 pass	implemented; V4 open
Must	F2 Gymnasium environment `TitrateRL/T2D-v0`	passes `check_env`; 168-day episode <2 s; rate-limit and shield tests pass	done (0.15 s)
Must	F3 Trained agents: PPO with R1–R3 (shield off) and shielded PPO with $R^\star$, 5 seeds each	checkpoints and learning curves stored; selection on validation patients	pipeline done; runs Wk 7–10
Must	F4 Rule-based policies B0–B3 behind the `Policy` interface	unit tests reproduce the published titration steps	done
Must	F5 Evaluation pipeline	one command regenerates metric tables, statistical tests and figures	done
Must	F6 Interactive dashboard	select a test patient and ≤ 3 policies; traces, doses, metrics, disclaimer; 24-week re-run <5 s; CSV export	done
Must	F7 Installable package	`pip install`; environment registered; documented API and example notebook	done
Must	RQ1–RQ3 analyses	H1–H3 decision rules applied to test-cohort data	Wk 10–13
Should	DQN agent; FPG-only PPO; λ grid for R3; simulator-tuned rule B4; robustness scenario set	completed and reported as exploratory	code ready
Could	Oracle per-patient fine-tuning; 2013 hypoglycemia model extensions; LSTM policy; SAC	started only if M3 is on time	—
Won't (this semester)	Real patient data or clinical deployment; Type 1 diabetes, pumps and bolus insulin; exercise, illness, missed injections; offline RL from EHRs; mobile app	out of scope; stated in README and model card	—

TABLE V
FOURTEEN-WEEK IMPLEMENTATION ROADMAP

Weeks	Planned milestone	Responsible member(s)
1–3	Requirements and architecture; repository, CI and packaging; T2D patient model, glargine model, integrator and cross-implementation tests	N. Khabibullaev, B. Uchqunov, I. Abralov, A. Abdusattarov
4–6	Cohort calibration; Gymnasium environment with reward, rate limit and shield; baselines B0–B3; gate G1; **M1** `v0.1-env`	I. Abralov, A. Abdusattarov, B. Uchqunov, N. Khabibullaev
7–8	PPO training with R1–R3; reward selection and power check (gate G2); **M2** `v0.2-agents`	A. Abdusattarov, N. Khabibullaev, B. Uchqunov
9–10	Evaluation pipeline, statistics and dashboard; shielded PPO-$R^\star$; RQ1 test evaluation	N. Khabibullaev, A. Abdusattarov, I. Abralov, B. Uchqunov
11–12	Full test-cohort evaluation (RQ2); packaging; buffer week; **M3** `v0.3-eval`	N. Khabibullaev, B. Uchqunov, A. Abdusattarov, I. Abralov
13	RQ3 analyses: correlation, dose behavior, generalization gap	N. Khabibullaev, A. Abdusattarov
14	Final report, documentation, model card and demonstration; **M4** `v1.0-mvp`	All members

III. Results

A. Overall Prototype Technical Evaluation (General Objective)

[Insert Results for General Objective here: High-level overview of the integrated titration framework, simulator verification and calibration checks (gate G1), runtime per episode, training throughput, and end-to-end pipeline testing.]

B. Reward Design (SO1)

[Insert Results for Reward Design here: TIR, $\text{TBR}_{<70}$, and $\text{TBR}_{<54}$ of PPO trained with R1, R2, and R3 on the held-out test cohort; median and 95% bootstrap CI; share of patients meeting the consensus targets; H1 test.]

C. Comparison with Clinical Algorithms (SO2)

[Insert Results for Comparison with Clinical Algorithms here: Maintenance-phase TIR and full-period $\text{TBR}_{<70}$ of the shielded PPO policy versus AT.LANTUS, Treat-to-Target, and INSIGHT; Hodges–Lehmann gains, Holm-adjusted p-values, non-inferiority intervals, time-to-target, and dose burden.]

D. Personalization by Insulin Sensitivity (SO3)

[Insert Results for Personalization here: Spearman correlation between insulin sensitivity and the TIR benefit; dose changes at matched fasting glucose by insulin-sensitivity tertile; train–test generalization gap.]

IV. Discussion

A. Overall Prototype Evaluation and Technical Feasibility

[Insert Discussion for General Objective here: Interpretation of simulator validity and calibration deviations, reproducibility of the pipeline, computational feasibility, and in silico scope boundaries.]

B. Reward Design (RQ1 / SO1)

[Insert Discussion for Reward Design here: Interpretation of RQ1, the efficacy–safety trade-off linked to Problem 1, overbasalization risk, and implications for encoding asymmetric clinical objectives.]

C. Comparison with Clinical Algorithms (RQ2 / SO2)

[Insert Discussion for Comparison with Clinical Algorithms here: Interpretation of RQ2 linked to Problem 2, the value of CGM features versus fasting-glucose-only titration, decision frequency, and simulator-specific tuning relative to related work.]

D. Personalization by Insulin Sensitivity (RQ3 / SO3)

[Insert Discussion for Personalization here: Interpretation of RQ3 linked to Problem 3, adaptation to unobserved insulin sensitivity, generalization to unseen patients, and limitations of in silico evidence.]

REFERENCES

[1] International Diabetes Federation, *IDF Diabetes Atlas*, 11th ed. Brussels, Belgium, 2025. [Online]. Available: https://diabetesatlas.org

[2] American Diabetes Association Professional Practice Committee for Diabetes, "9. Pharmacologic approaches to glycemic treatment: Standards of Care in Diabetes—2026," *Diabetes Care*, vol. 49, suppl. 1, pp. S183–S215, 2026, doi: 10.2337/dc26-S009.

[3] American Diabetes Association Professional Practice Committee for Diabetes, "6. Glycemic goals, hypoglycemia, and hyperglycemic crises: Standards of Care in Diabetes—2026," *Diabetes Care*, vol. 49, suppl. 1, pp. S132–S149, 2026, doi: 10.2337/dc26-S006.

[4] M. Mocarski *et al.*, "Slow titration and delayed intensification of basal insulin among patients with type 2 diabetes," *J. Manag. Care Spec. Pharm.*, vol. 24, no. 4, pp. 390–400, 2018, doi: 10.18553/jmcp.2017.17218.

[5] M. C. Riddle, J. Rosenstock, and J. Gerich, "The treat-to-target trial: Randomized addition of glargine or human NPH insulin to oral therapy of type 2 diabetic patients," *Diabetes Care*, vol. 26, no. 11, pp. 3080–3086, 2003, doi: 10.2337/diacare.26.11.3080.

[6] M. Davies, F. Storms, S. Shutler, M. Bianchi-Biscay, and R. Gomis, "Improvement of glycemic control in subjects with poorly controlled type 2 diabetes: Comparison of two treatment algorithms using insulin glargine," *Diabetes Care*, vol. 28, no. 6, pp. 1282–1288, 2005, doi: 10.2337/diacare.28.6.1282.

[7] H. C. Gerstein *et al.*, "A randomized trial of adding insulin glargine vs. avoidance of insulin in people with type 2 diabetes on either no oral glucose-lowering agents or submaximal doses of metformin and/or sulphonylureas. The Canadian INSIGHT study," *Diabet. Med.*, vol. 23, no. 7, pp. 736–742, 2006, doi: 10.1111/j.1464-5491.2006.01881.x.

[8] C. Dalla Man, R. A. Rizza, and C. Cobelli, "Meal simulation model of the glucose-insulin system," *IEEE Trans. Biomed. Eng.*, vol. 54, no. 10, pp. 1740–1749, 2007, doi: 10.1109/TBME.2007.893506.

[9] B. P. Kovatchev, M. Breton, C. Dalla Man, and C. Cobelli, "In silico preclinical trials: A proof of concept in closed-loop control of type 1 diabetes," *J. Diabetes Sci. Technol.*, vol. 3, no. 1, pp. 44–55, 2009, doi: 10.1177/193229680900300106.

[10] J. Xie, "Simglucose v0.2.1," GitHub repository, 2018. [Online]. Available: https://github.com/jxx123/simglucose

[11] J. Bonet, R. Visentin, and C. Dalla Man, "Smart algorithms for efficient insulin therapy initiation in individuals with type 2 diabetes: An in silico study," *J. Diabetes Sci. Technol.*, vol. 19, no. 5, pp. 1271–1279, 2025, doi: 10.1177/19322968241245930.

[12] I. Fox, J. Lee, R. Pop-Busui, and J. Wiens, "Deep reinforcement learning for closed-loop blood glucose control," in *Proc. Mach. Learn. Healthcare Conf.*, PMLR, vol. 126, 2020, pp. 508–536.

[13] M. Tejedor, A. Z. Woldaregay, and F. Godtliebsen, "Reinforcement learning application in diabetes blood glucose control: A systematic review," *Artif. Intell. Med.*, vol. 104, Art. no. 101836, 2020, doi: 10.1016/j.artmed.2020.101836.

[14] G. Wang *et al.*, "Optimized glycemic control of type 2 diabetes with reinforcement learning: A proof-of-concept trial," *Nat. Med.*, vol. 29, no. 10, pp. 2633–2642, 2023, doi: 10.1038/s41591-023-02552-9.

[15] M. Panagiotou *et al.*, "Personalized insulin adjustment with reinforcement learning: An in-silico validation for people with diabetes on intensive insulin treatment," *IEEE Access*, vol. 13, pp. 148436–148455, 2025, doi: 10.1109/ACCESS.2025.3600738.

[16] M. Kwon, J. Lamp, and L. Feng, "Safety generalization under distribution shift in safe reinforcement learning: A diabetes testbed," in *Proc. 43rd Int. Conf. Mach. Learn.*, PMLR, vol. 306, 2026, pp. 61952–61985.

[17] D. M. Karl, J. Gill, R. Zhou, and M. C. Riddle, "Clinical predictors of risk of hypoglycaemia during addition and titration of insulin glargine for type 2 diabetes mellitus," *Diabetes Obes. Metab.*, vol. 15, no. 7, pp. 622–628, 2013, doi: 10.1111/dom.12072.

[18] O. Gottesman *et al.*, "Guidelines for reinforcement learning in healthcare," *Nat. Med.*, vol. 25, no. 1, pp. 16–18, 2019, doi: 10.1038/s41591-018-0310-5.

[19] C. Dalla Man, D. M. Raimondo, R. A. Rizza, and C. Cobelli, "GIM, simulation software of meal glucose–insulin model," *J. Diabetes Sci. Technol.*, vol. 1, no. 3, pp. 323–330, 2007, doi: 10.1177/193229680700100303.

[20] M. Schiavon *et al.*, "Modeling subcutaneous absorption of long-acting insulin glargine in type 1 diabetes," *IEEE Trans. Biomed. Eng.*, vol. 67, no. 2, pp. 624–631, 2020, doi: 10.1109/TBME.2019.2919250.

[21] M. Lepore *et al.*, "Pharmacokinetics and pharmacodynamics of subcutaneous injection of long-acting human insulin analog glargine, NPH insulin, and ultralente human insulin and continuous subcutaneous infusion of insulin lispro," *Diabetes*, vol. 49, no. 12, pp. 2142–2148, 2000, doi: 10.2337/diabetes.49.12.2142.

[22] S. K. Lam, A. Pitrou, and S. Seibert, "Numba: A LLVM-based Python JIT compiler," in *Proc. 2nd Workshop LLVM Compiler Infrastructure HPC*, 2015, pp. 1–6, doi: 10.1145/2833157.2833162.

[23] M. Breton and B. Kovatchev, "Analysis, modeling, and simulation of the accuracy of continuous glucose sensors," *J. Diabetes Sci. Technol.*, vol. 2, no. 5, pp. 853–862, 2008, doi: 10.1177/193229680800200517.

[24] *In Vitro Diagnostic Test Systems—Requirements for Blood-Glucose Monitoring Systems for Self-Testing in Managing Diabetes Mellitus*, ISO 15197:2013, International Organization for Standardization, Geneva, Switzerland, 2013.

[25] M. Towers *et al.*, "Gymnasium: A standard interface for reinforcement learning environments," in *Adv. Neural Inf. Process. Syst., Datasets and Benchmarks Track*, vol. 38, 2025, arXiv: 2407.17032.

[26] B. P. Kovatchev, D. J. Cox, L. A. Gonder-Frederick, and W. Clarke, "Symmetrization of the blood glucose measurement scale and its applications," *Diabetes Care*, vol. 20, no. 11, pp. 1655–1658, 1997, doi: 10.2337/diacare.20.11.1655.

[27] J. Schulman, F. Wolski, P. Dhariwal, A. Radford, and O. Klimov, "Proximal policy optimization algorithms," arXiv preprint arXiv:1707.06347, 2017.

[28] V. Mnih *et al.*, "Human-level control through deep reinforcement learning," *Nature*, vol. 518, no. 7540, pp. 529–533, 2015, doi: 10.1038/nature14236.

[29] A. Raffin, A. Hill, A. Gleave, A. Kanervisto, M. Ernestus, and N. Dormann, "Stable-Baselines3: Reliable reinforcement learning implementations," *J. Mach. Learn. Res.*, vol. 22, no. 268, pp. 1–8, 2021.

[30] P. Henderson, R. Islam, P. Bachman, J. Pineau, D. Precup, and D. Meger, "Deep reinforcement learning that matters," in *Proc. AAAI Conf. Artif. Intell.*, vol. 32, no. 1, 2018, pp. 3207–3214, doi: 10.1609/aaai.v32i1.11694.

[31] M. Alshiekh, R. Bloem, R. Ehlers, B. Könighofer, S. Niekum, and U. Topcu, "Safe reinforcement learning via shielding," in *Proc. AAAI Conf. Artif. Intell.*, vol. 32, no. 1, 2018, pp. 2669–2678, doi: 10.1609/aaai.v32i1.11797.

[32] T. Battelino *et al.*, "Clinical targets for continuous glucose monitoring data interpretation: Recommendations from the international consensus on time in range," *Diabetes Care*, vol. 42, no. 8, pp. 1593–1603, 2019, doi: 10.2337/dci19-0028.

[33] T. Battelino *et al.*, "Continuous glucose monitoring and metrics for clinical trials: An international consensus statement," *Lancet Diabetes Endocrinol.*, vol. 11, no. 1, pp. 42–57, 2023, doi: 10.1016/S2213-8587(22)00319-9.

[34] K. Cowart, "Overbasalization: Addressing hesitancy in treatment intensification beyond basal insulin," *Clin. Diabetes*, vol. 38, no. 3, pp. 304–310, 2020, doi: 10.2337/cd19-0061.

[35] R. M. Bergenstal *et al.*, "Glucose management indicator (GMI): A new term for estimating A1C from continuous glucose monitoring," *Diabetes Care*, vol. 41, no. 11, pp. 2275–2280, 2018, doi: 10.2337/dc18-1581.