

Let's simulate some Lévy processes :

- Keep in mind that $X_t = \log(S_t/S_0)$.

I) Let's start with jump-diffusion :

We simulate here "jump diffusion" processes, which can be written as :

$X_t = \mu t + \sigma W_t + \sum_{s \in [0,t)} \Delta X_s = \mu t + \sigma W_t + \sum_{i=1}^{N(t)} Y_i$, N being a Poisson Process of intensity λ and $(Y_i)_i$ i.i.d random variables.

I-1) Merton Model :

- $Y_i \sim N(\mu_J, \delta_J^2)$
- Defined by $(\sigma, \lambda, \mu_J, \delta_J)$

Process parameters :

```
S0 = 1;           % Initial value
mu = 0.05;        % Drift (as in GBM)
sigma = 0.4;      % Volatility (as in GBM)
lambda = 2;       % Poisson intensity/rate
muJ = 0.01;       % Jumpsize mean
deltaJ = 0.2;     % Jumpsize std
```

Simulation parameters :

```
T = 2;           % Maturity
M = 100;         % Number of steps in time
dt = T/M;        % Time step
```

Conditional simulation of jump times (the easiest way) :

(cf ALGORITHM 6.2 from Simulate_Jump_Diffusion.pdf)

```
X = zeros(M + 1, 1); % X = (X0, X1, ..., XM)
Z = randn(M, 1);     % Z = (Z1, Z2, ..., ZM) [BM]

NT = poissrnd(lambda*T); % Nb of jumps in [0,T]


% Conditional to NT, we simulate when jumps occur
% via an Uniform Distribution :
jumpT = sort(rand(1,NT)*T);

% In Merton, jumpSize ~ Normal(muJ, deltaJ^2) :
jumpSize = muJ + deltaJ * randn(NT,1);
for i=1:M
    % Continuous part of the process :
    X(i+1) = X(i) + mu * dt + sigma * dt * Z(i);
    % Jump part :
    for j=1:NT
```

```

    % Did a jump occur in ](i-1)dt, idt] ?
    if jumpT(j) > (i-1)*dt && jumpT(j) <= i*dt
        X(i+1) = X(i+1) + jumpSize(j);
    end
end
end
figure; % Plot
plot(S0 * exp(X));
title("One simulated path via Merton model :")

```



Another simulation of jump times : (cf ALGO 6.1 from Simulate_Jump_Diffusion.pdf) we simulate jumps inside the loop, at each time step.

```

X = zeros(M + 1, 1); % X = (X0, X1, ..., XM)
Z = randn(M, 1);      % Z = (Z1, Z2, ..., ZM) [BM]

for i=1:M
    % Continuous part of the process :
    X(i+1) = X(i) + mu * dt + sigma * dt * Z(i);
    % Jump part :
    Ndt = poissrnd(lambda * dt);


    if Ndt == 0
        J = 0; % No jump !
    else
        J = sum(muJ + deltaJ * randn(Ndt,1)); % Sum of the jump(s)
        % that happened during this time step.
    end
end

```

```

end
X(i+1) = X(i+1) + J; % @ each time step : add the jump(s).
end
figure; % Plot
plot(S0 * exp(X));
title("One simulated path via Merton model, V2 :")

```



I-2) Kou Model :

- Positive jumpsize $\sim \exp(\lambda^+)$;
- Negative jumpsize $\sim \text{Exp}(\lambda^-)$;