

How can vector calculus be used to model
and analyze incompressible fluid flow in
two-dimensional spaces, and what insights
can this provide about real-world fluid
systems?

Mathematics AA HL

Word Count: 86

Contents

1 Introduction 2

1.1 Scope 2

1.2 Background 2

1.3 Fluid dynamics 2

1.4 The Navier-Stokes equations 3

2 References 4

1 Introduction

1.1 Scope

This essay will for simplicity's sake only cover fluid flow in \mathbb{R}^2 spaces; an analysis of fluid flow in \mathbb{R}^3 spaces would be orders of magnitude more complex. Furthermore, only incompressible fluids ($\nabla \cdot \mathbf{F} = 0$), such that there are no sinks nor wells, will be analyzed. Test

1.2 Background

Vector calculus is the mathematical study of applying multi-variable calculus to vector valued functions, often for the spaces \mathbb{R}^2 and \mathbb{R}^3 .

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n \ni n > 1, n \in \mathbb{Z}$$
$$\rightsquigarrow \nabla \triangleq \left. \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \\ \vdots \end{bmatrix} \right\} n \text{ times} \quad (1)$$

1.3 Fluid dynamics

An incompressible fluid is any fluid such that $\nabla \cdot \mathbf{F} = 0$, which is to say that the divergence of the fluid is 0.

1.4 The Navier-Stokes equations

$$\frac{\mathbf{D}f}{\mathbf{D}t} = \iiint_V \left(\frac{\mathbf{D}\rho}{\mathbf{D}t} + \rho(\nabla \cdot u) \right) dV \quad (2)$$

Lorem ipsum dolor sit amet [Peyret and Taylor, 2012]

2 References

[Peyret and Taylor, 2012] Peyret, R. and Taylor, T. D. (2012). *Computational methods for fluid flow*. Springer Science & Business Media.