How can vector calculus be used to model and analyze incompressible fluid flow in two-dimensional spaces, and what insights can this provide about real-world fluid systems?

Mathematics AA HL

Word Count: 150

# Contents

| 1 Introduction |     | roduction                   | 2 |
|----------------|-----|-----------------------------|---|
|                | 1.1 | Scope                       | 2 |
|                | 1.2 | Background                  | 2 |
|                | 1.3 | Fluid dynamics              | 2 |
|                | 1.4 | The Navier-Stokes equations | 2 |

## 1 Introduction

#### 1.1 Scope

This essay will for simplicity's sake only cover fluid flow in  $\mathbb{R}^2$  spaces; an alaysis of fluid flow in  $\mathbb{R}^3$  spaces would be orders of magnitude more complex. Furthermore, only incompressible fluids  $(\nabla \cdot \mathbf{F} = 0)$ , such that there are no sinks nore wells, will be analyzed. Test

#### 1.2 Background

Vector calculus is the mathematical study of applying multi-variable calculus to vector valued functions, often for the spaces  $\mathbb{R}^2$  and  $\mathbb{R}^3$ .

$$f: \mathbb{R}^n \to \mathbb{R}^n \ni n > 1, n \in \mathbb{Z}$$

$$\leadsto \nabla \stackrel{\Delta}{=} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \\ \vdots \end{bmatrix}$$
  $n \text{ times}$  (1)

### 1.3 Fluid dynamics

An incompressible fluid is any fluid such that  $\nabla \cdot \mathbf{F} = 0$ , which is to say that the divergence of the fluid is 0.

#### 1.4 The Navier-Stokes equations

$$\frac{\mathbf{D}f}{\mathbf{D}\mathbf{t}} = \iiint_{V} (\frac{\mathbf{D}\rho}{\mathbf{D}\mathbf{t}} + \rho(\nabla \cdot u))dV$$
 (2)

Lorem ipsum dolor sit amet