

# Discrete Math

---

# Contents

## Part I Discrete Math: Logic 1

### Chapter I Propositional Logic 1

§ 1.1	Connectives and Truth	
	Assingments . . . . .	1
§ 1.2	Consequence and	
	Equivalent . . . . .	1
1	The definition of	
	consequence and	
	logically equivalent	2
2	Important properties	2
3	Prove Logical	
	Equivalence . . . . .	3

4	Relation among	
	tautologies, con-	
	tradictions, satis-	
	fiabile asser-tions,	
	consequence re-	
	lations and logic	
	equivalence . . . . .	4

§ 1.3	Normal Forms . . . . .	4
-------	------------------------	---

### Chapter II First Order Logic, FOL 6

§ 2.1	The syntax of first or-	
	der language . . . . .	6
§ 2.2	The semantics of first	
	order language . . . . .	6
1	Structure . . . . .	7
2	Interpretation . . . . .	7

## Part I Discrete Math: Logic

### Chapter I Propositional Logic

#### § 1.1 Connectives and Truth Assignments

**Define 1.1.1** (Truth table of Connectives) (Omitted)

**Define 1.1.2** (Truth Assignments) Suppose  $\Sigma$  is the set of propositional variables. A mapping from  $\Sigma$  to  $\{\mathbf{T}, \mathbf{F}\}$  called a truth assignment.

**Define 1.1.3** Suppose  $\Sigma$  is the set of propositional variables and  $\mathcal{J} : \Sigma \rightarrow \{\mathbf{T}, \mathbf{F}\}$  is a truth assignment. The truth value of the compound proposition on  $\mathcal{J}$  ...  
(Omitted)

**Define 1.1.4** (Tautology, contradiction) (Omitted)

**Define 1.1.5** (Contingency, Satisfiable) A contingency is a compound proposition that is neither a tautology nor a contradiction.  
A compound proposition is **satisfiable** if it is true under some truth assignment.

#### § 1.2 Consequence and Equivalent

## 1 The definition of consequence and logically equivalent

**Define 1.2.1 (Consequence)** Suppose  $\Phi$  is a set of propositions and  $\psi$  is one single proposition. We say that  $\psi$  is a consequence of  $\Phi$ , written as  $\Phi \models \psi$ , if  $\Phi$ 's being all true implies that  $\psi$  is also true.

In other words,  $\Phi \models \psi$  if for any truth assignment  $\mathcal{J}$ ,  $\llbracket \phi \rrbracket_{\mathcal{J}} = \mathbf{T}$  for any  $\phi \in \Phi$  implies  $\llbracket \psi \rrbracket_{\mathcal{J}} = \mathbf{T}$ .

**Define 1.2.2 (Logically Equivalent)**  $\phi$  is a logically equivalent to  $\psi$ , written as  $\phi \equiv \psi$ , if  $\phi$ 's truth value and  $\psi$ 's truth value are the same under any situation. In other words,  $\phi \equiv \psi$  if  $\llbracket \phi \rrbracket_{\mathcal{J}} = \llbracket \psi \rrbracket_{\mathcal{J}}$  for any truth assignment  $\mathcal{J}$ .

**Example 1.2.1**  $\Phi = \{ \}$ ,  $\psi = p \vee \neg p$ ,  $\Phi \models \psi$

## 2 Important properties

### Theorem 1.2.1

- $\phi \vee \neg \phi$  is an tautology
- $\phi \wedge \neg \phi$  is a contradiction
- $\phi, \psi \models \phi \wedge \psi$  ( $\wedge$ -Introduction)
- $\phi \wedge \psi \models \phi$  ( $\wedge$ -Elimination)
- $\phi \models \phi \vee \psi$  ( $\vee$ -Introduction)
- If  $\Phi, \phi_1 \models \psi$ ,  $\Phi, \phi_2 \models \psi$ , then  $\Phi, \phi_1 \vee \phi_2 \models \psi$  ( $\vee$ -Elimination)

**Proof (Proof of the last one)** Suppose  $\llbracket \phi \rrbracket_{\mathcal{J}} = \mathbf{T}$ ,  $\llbracket \phi_1 \vee \phi_2 \rrbracket_{\mathcal{J}} = \mathbf{T}$ .

Then at least one of the following holds:  $\llbracket \phi_1 \rrbracket_{\mathcal{J}} = \mathbf{T}$ ,  $\llbracket \phi_2 \rrbracket_{\mathcal{J}} = \mathbf{T}$ . □

**Theorem 1.2.2 (Contrapositive)** If  $\Phi, \neg \phi \models \psi$ , then  $\Phi, \neg \psi \models \phi$

**Theorem 1.2.3**

- $\neg(\neg q) \equiv q$  (Double Negation)
- $\phi \wedge \phi \equiv \phi, \quad \phi \vee \phi \equiv \phi$  (Idempotent Laws)
- $\phi \wedge \psi \equiv \psi \wedge \phi, \quad \phi \vee \psi \equiv \psi \vee \phi$  (Commutative Laws)
- $\phi \vee (\psi \wedge \chi) \equiv (\phi \vee \psi) \wedge (\phi \vee \chi), \quad \phi \wedge (\psi \vee \chi) \equiv (\phi \wedge \psi) \vee (\phi \wedge \chi)$   
(Distributive Laws)
- $\neg(q \wedge p) \equiv \neg q \vee \neg p, \quad \neg(q \vee p) \equiv \neg q \wedge \neg p$  (De Morgan's Laws)
- $\phi \wedge (\neg \phi) \equiv \mathbf{F}, \quad \phi \vee (\neg \phi) \equiv \mathbf{T}$  (Negation Laws)
- $\phi \wedge \mathbf{T} \equiv \phi, \quad \phi \vee \mathbf{F} \equiv \phi, \quad \phi \wedge \mathbf{F} \equiv \mathbf{F}, \quad \phi \vee \mathbf{T} \equiv \mathbf{T}$  (Laws of logical constants)
- $\phi \vee (\phi \wedge \psi) \equiv \phi, \quad \phi \wedge (\phi \vee \psi) \equiv \phi$  (Absorption Laws)

**3 Prove Logical Equivalence**

**Theorem 1.2.4** (Transitivity) If  $\phi \equiv \psi$  and  $\psi \equiv \chi$ , then  $\phi \equiv \chi$ .

**Theorem 1.2.5** (Congruence Property)

- If  $\phi \equiv \psi$ , then  $\neg \phi \equiv \neg \psi$
- If  $\phi_1 \equiv \phi_2, \psi_1 \equiv \psi_2$ , then  $\phi_1 \wedge \psi_1 \equiv \phi_2 \wedge \psi_2$
- If  $\phi_1 \equiv \phi_2, \psi_1 \equiv \psi_2$ , then  $\phi_1 \vee \psi_1 \equiv \phi_2 \vee \psi_2$

**Theorem 1.2.6** (Reflexivity)  $\phi \equiv \phi$

## 4 Relation among tautologies, contradictions, satisfiable assertions, consequence relations and logic equivalence

### Theorem 1.2.7

- $\phi_1, \phi_2, \dots, \phi_n \models \psi$  iff.  $\left( \bigwedge_{k=1}^n \phi_k \right) \wedge \neg \psi$  is not satisfiable.
- $\{ \} \models \phi$  iff.  $\phi$  is an tautology.
- $\phi \equiv \psi$  iff.  $\phi \models \psi$  and  $\psi \models \phi$ .

**Theorem 1.2.8** If  $\phi \models \psi$  and  $\psi \models \chi$ , then  $\phi \models \chi$ .

## § 1.3 Normal Forms

### Define 1.3.1 (Disjunctive Normal Form, DNF)

- A **literal** is a propositional variable or its negation.
- A **conjunctive clause** is a conjunctions of literals.
- A **compound proposition** is in disjunctive normal form if it is a disjunction of conjunctive clauses.

### Define 1.3.2 (Conjunctive Normal Form, CNF)

(Similar as above)

### Example 1.3.1

- literals  $x, y, z, p, q, r, \neg q$
- conjunctive clauses  $p, p \wedge q, \neg p \wedge q$
- DNF  $p, p \vee (\neg q \wedge r), \neg p \vee (q \wedge p \wedge r)$

**Theorem 1.3.1** Every compound proposition is logically equivalent to some compound proposition in DNF.

**Proof (Proof 1)** Suppose that the compound proposition  $\phi$  consists of the literals  $p_1, p_2, \dots, p_n$ .

For all  $\mathcal{J}$  as a interpretation, we only need to prove that

$$\phi \equiv \bigvee_{\llbracket \phi \rrbracket_{\mathcal{J}} = \mathbf{T}} \left( \bigwedge_{\mathcal{J}(p_i) = \mathbf{T}} p_i \wedge \bigwedge_{\mathcal{J}(p_i) = \mathbf{F}} \neg p_i \right) = \mathbf{T}$$

Consider a specific interpretation  $\mathcal{J}_0$ , if  $\llbracket \phi \rrbracket_{\mathcal{J}} = \mathbf{T}$ , then

$$\left[ \bigvee_{\llbracket \phi \rrbracket_{\mathcal{J}} = \mathbf{T}} \left( \bigwedge_{\mathcal{J}(p_i) = \mathbf{T}} p_i \wedge \bigwedge_{\mathcal{J}(p_i) = \mathbf{F}} \neg p_i \right) \right]_{\mathcal{J}_0} = \left[ \left( \bigwedge_{\mathcal{J}_0(p_i) = \mathbf{T}} p_i \wedge \bigwedge_{\mathcal{J}_0(p_i) = \mathbf{F}} \neg p_i \right) \right]_{\mathcal{J}_0}$$

If  $\mathcal{J}_0(p_i) = \mathbf{T}$ , then  $\llbracket p_i \rrbracket_{\mathcal{J}_0} = \mathbf{T}$ ,

if  $\mathcal{J}_0(p_i) = \mathbf{F}$ , then  $\llbracket \neg p_i \rrbracket_{\mathcal{J}_0} = \mathbf{T}$ .

So

$$\left[ \left( \bigwedge_{\mathcal{J}_0(p_i) = \mathbf{T}} p_i \wedge \bigwedge_{\mathcal{J}_0(p_i) = \mathbf{F}} \neg p_i \right) \right]_{\mathcal{J}_0} = \mathbf{T}$$

□

**Proof (Proof 2)** Define  $DNF(\phi)$  as follow and prove that  $DNF(\phi) \equiv \phi$ .

**Define 1.3.3** •  $DNF(\phi) \triangleq DNF_2(DNF_1(\phi))$

•  $DNF_1(\neg\neg\phi) = DNF_1(\phi)$ .

(The De Morgan's law)

$DNF_1(\phi \wedge \psi) = DNF_1(\phi) \wedge DNF_1(\psi)$  ( $\vee$  is the same)

$DNF_1(l) = l$   $l$  is a literal.

•  $DNF_2(l) = l$   $l$  is a literal,

$DNF_2(\phi \vee \psi) = DNF_2(\phi) \vee DNF_2(\psi)$

If  $\phi = \bigvee_{i=1}^n \phi_i, \psi = \bigvee_{j=1}^m \psi_j$ , then

$$DNF_2(\phi \wedge \psi) = \bigvee_{i=1}^n \bigvee_{j=1}^m (\phi_i \wedge \psi_j)$$

Then it's obvious that  $\phi \equiv DNF(\phi)$  and  $DNF(\phi)$  is a DNF.

□

**Theorem 1.3.2** Every compound proposition is logically equivalent to some compound proposition in CNF.

**Proof** (Similar as above)

**Example 1.3.2 (\*)** The CDCL algorithm.  
(Suspended now)

## Chapter II First Order Logic, FOL

### § 2.1 The syntax of first order language

#### Define 2.1.1

- Predicate Logic's Language
  - Variables  $x, y, z, \dots$
  - Constants  $c_1, c_2, \dots$
  - Prelicates  $P, Q, R, \dots$
  - Functions  $f, g, h, \dots$
  - Logic patterns  $\exists, \forall, \wedge, \vee, \neg$
- Terms  $x, y, c_1, c_2, f(x), g(x, y), \dots$
- propositions  $P(x), Q(f(x, g(x, y))), \exists x \forall y R(x, g(y)), \dots$

### § 2.2 The semantics of first order language



## 1 Structure

**Define 2.2.1** ( $S$ -structure)

Given a symbol set  $S$ , an  $S$ -structure  $\mathcal{A} = (A, \alpha)$  contains

- a domain  $A$ , which is a non-empty set.
- an interpretation of every predicate symbol.

**Example 2.2.1** if  $P$  is a symbol of binary predicate, then  $\alpha(P)$  is a mapping from  $A \times A$  to  $\{\mathbf{T}, \mathbf{F}\}$ .

- an interpretation of every function symbol.

**Example 2.2.2** if  $f$  is a symbol of unary function, then  $\alpha(f)$  is a mapping from  $A$  to  $A$ .

- an interpretation of every constant symbol.

**Example 2.2.3** if  $s$  is a constant symbol,  $\alpha(c)$  is an element in domain  $A$ .

With a structure, we can determine the truth of an closed proposition.

## 2 Interpretation