

Assignment 3

Problem 1. Gauss elimination and LU decomposition (50 marks)

Consider a system of three linear equations

$$2x + y + z = 4$$

$$x + 3y + z = 0$$

$$x - y + 3z = 6$$

- a) Write the above equation in the form of $Ap = b$.
- b) Find the solution p of the above equations using Gauss Elimination.
- c) Decompose the coefficient matrix A into a lower triangular matrix L and upper triangular U matrix using Gauss Elimination. **Hint:** You already have the elements of L and U after solving step (b).
- d) Find the solution p of the above equations using forward and back substitution with L and U , respectively, i.e., by solving

$$Lm = b$$

$$Up = m$$

- e) Determinant of a triangular matrix (e.g., L and U) can be easily calculated by multiplying the main diagonal elements. For example,

$$\det(L) = L_{11} L_{22} \dots L_{nn} = \prod_{i=1}^n L_{ii}$$

$$\det(U) = U_{11} U_{22} \dots U_{nn} = \prod_{i=1}^n U_{ii}$$

where n is the size of matrices L and U . Therefore, Gauss elimination provides a convenient way to calculate the determinant of a matrix. Check if the determinant of A is same as the determinant of U .

- f) Calculate the inverse of matrix A using the triangular matrices L and U .

Problem 2. Thomas Algorithm/Tridiagonal Matrix Algorithm (20 marks)

$$0.8x - 0.4y = 41$$

$$-0.4x + 0.8y - 0.4z = 25$$

$$-0.4y + 0.8z = 105$$

- a) Write the above equations in the form of $Ap = b$.
- b) Find the solution p of the above equations using the Thomas algorithm.
- c) Comment on the number of calculations performed in part (b), compared to Problem 1, where you used Gauss Elimination.

Problem 3. Gauss Seidel and relaxation (30 marks)

Consider a system of two linear equations

$$x + 2y = 1 \quad (1)$$

$$x - y = 4 \quad (2)$$

- a) Use **four iterations** of the Gauss Seidel method to solve for x and y . Use Eq. (1) to solve for x and Eq. (2) to solve for y . Report the values of x and y and the approximate percent relative error in x and y at the end of each iteration. What do you observe? Is the method converging?
- b) Use underrelaxation with $\lambda = 0.6$ and repeat part (a). Does underrelaxation help?
- c) Make the system of equations diagonally dominant. Repeat step (a). What do you observe? Is the method converging?