

Gödel's incompleteness theorems

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' *extends* S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. *If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.*

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' *extends* S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. *If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.*

def: soundness/correctness Proof system S is *sound/correct* if all provable statements are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' *extends* S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. *If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.*

def: soundness/correctness Proof system S is *sound/correct* if all provable statements are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

def: consistency an approximation to correctness. Proof system is *consistent* if for no statement w

$$S \vdash w \text{ and } S \vdash (\sim w)$$

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' *extends* S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. *If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.*

def: soundness/correctness Proof system S is *sound/correct* if all provable statements are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

def: consistency an approximation to correctness. Proof system is *consistent* if for no statement w

$$S \vdash w \text{ and } S \vdash (\sim w)$$

def: completeness Proof system S is *complete* if for all statements $w \in L$ either w or $\sim w$ is provable

$$S \vdash w \quad \text{or} \quad S \vdash (\sim w)$$

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' extends S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.

def: soundness/correctness Proof system S is *sound/correct* if all proven predicates are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

def: consistency an approximation to correctness. Proof system is *consistent* if for no predicate w

$$S \vdash w \text{ and } S \vdash (\sim w)$$

def: completeness Proof system S is *complete* if for all predicates $w \in L$ either w or $\sim w$ is provable

$$S \vdash w \quad \text{or} \quad S \vdash (\sim w)$$

Lemma 2. If S is complete, then the language

$$\{w \in L : S \vdash w\}$$

is decidable

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' extends S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.

def: soundness/correctness Proof system S is *sound/correct* if all proven predicates are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

def: consistency an approximation to correctness. Proof system is *consistent* if for no predicate w

$$S \vdash w \text{ and } S \vdash (\sim w)$$

def: completeness Proof system S is *complete* if for all predicates $w \in L$ either w or $\sim w$ is provable

$$S \vdash w \quad \text{or} \quad S \vdash (\sim w)$$

Lemma 2. If S is complete, then the language

$$\{w \in L : S \vdash w\}$$

is decidable

Proof. Enumerate all proofs. If proof of w appears, accept. If proof of $\sim w$ appears, reject. \square

1 Properties of proof systems

$$S = (\Sigma, L, A, R) \quad , S' (= \Sigma, L', A', R')$$

def: extension We say S' extends S and write $S \subseteq S'$ if

$$\Sigma \subseteq \Sigma' \quad , \quad L \subseteq L' \quad , \quad A \subseteq A' \quad , \quad R \subseteq R'$$

Lemma 1. If $S \subseteq S'$ and $S \vdash w$ then $S' \vdash w$.

def: soundness/correctness Proof system S is *sound/correct* if all proven predicates are true, i.e. if for all $w \in L$:

$$S \vdash w \Rightarrow w \text{ is true}$$

def: consistency an approximation to correctness. Proof system is *consistent* if for no predicate w

$$S \vdash w \text{ and } S \vdash (\sim w)$$

def: completeness Proof system S is *complete* if for all predicates $w \in L$ either w or $\sim w$ is provable

$$S \vdash w \quad \text{or} \quad S \vdash (\sim w)$$

Lemma 2. If S is complete, then the language

$$\{w \in L : S \vdash w\}$$

is decidable

Proof. Enumerate all proofs. If proof of w appears, accept. If proof of $\sim w$ appears, reject. \square

Lemma 3. If S is correct and complete, then the set

$$T = \{w \in L : w \text{ is true}\}$$

why?

is decidable

2 First Incompleteness Theorem

Lemma 4. *Let S be any proof system whose language is as expressive as L_Z*

$$S = (\Sigma_Z, L_Z, A, R)$$

If S is sound, then it is incomplete.

- reduce Halting problem H to T with lemma 5
- if Z_E would be complete, then by lemma 3 T would be decidable

Lemma 5. *There is a total computable function*

$$H : \{0, 1, \#\} \rightarrow L_E$$

such that

$$H(u\#v) \text{ true} \Leftrightarrow \text{TM } M_u \text{ started with } v \text{ halts}$$

2 First Incompleteness Theorem

Lemma 4. *Let S be any proof system whose language is as expressive as L_Z*

$$S = (\Sigma_Z, L_Z, A, R)$$

If S is sound, then it is incomplete.

- reduce Halting problem H to T with lemma 5
- if Z_E would be complete, then by lemma 3 T would be decidable

Lemma 5. *There is a total computable function*

$$H : \{0, 1, \#\} \rightarrow L_E$$

such that

$$H(u\#v) \text{ true} \Leftrightarrow \text{TM } M_u \text{ started with } v \text{ halts}$$

from Sipser (MIT): Introduction to the Theory of Computation. A GREAT book. 'the actual construction of ... is too complicated to present here'.

- in contrast (KIU):

2 First Incompleteness Theorem

Lemma 5. *There is a total computable function*

$$H : \{0, 1, \#\} \rightarrow L_E$$

such that

$$H(u\#v) \text{ true} \Leftrightarrow \text{TM } M_u \text{ started with } v \text{ halts}$$

Consider 1-tape TM

$$M_u = (Z, A, \delta, z_0, E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

1. $k_0 = B \dots B z_0 v B \dots B$
2. $k_i \vdash k_{i+1}$ for $i < t$
3. $|k_i| = |k_j|$ for all i, j
4. in no k_i if a $z \in Z$ first or last element
5. k_t is endconfiguraion

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

-

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

-

$$\psi(\varepsilon) = 0$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$, \})^*$ as number representation to base p .

-

$$\psi : A \cup Z \cup \{\$, \} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

-

$$\psi(\varepsilon) = 0$$

- extend to

$$\psi : (A \cup Z \cup \{\$, \})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

- expression whose value codes $a \in A \cup Z \cup \{\$, \}$

$$\hat{a} = \overline{\psi(a)} \in T$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

-

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

-

$$\psi(\varepsilon) = 0$$

- extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

- expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

$$v=xy$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

-

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

-

$$\psi(\varepsilon) = 0$$

- extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

- expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

obviously using decomposition lemma for base p numbers. Top down:

-

$$v = [x, y] \quad :\equiv \quad (y = 0 \wedge v = x) \vee (\sim y = 0 \wedge \exists u (v = y + u \cdot x \wedge u = p^{|y|}))$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

-

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

-

$$\psi(\varepsilon) = 0$$

- extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

- expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

obviously using decomposition lemma for base p numbers. Top down:

-

$$v = [x, y] \quad :\equiv \quad (y = 0 \wedge v = x) \vee (\sim y = 0 \wedge \exists u (v = y + u \cdot x \wedge u = p^{|y|}))$$

-

$$u = p^{|y|} \quad :\equiv \quad u \text{ is power of } p \\ \wedge y < u \wedge u \leq p \cdot u$$

example:

$$456 < 1000 < 10 \cdot 456$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

•

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

•

$$\psi(\varepsilon) = 0$$

• extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

• expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

obviously using decomposition lemma for base p numbers. Top down:

•

$$v = [x, y] \quad :\equiv \quad (y = 0 \wedge v = x) \vee (\sim y = 0 \wedge \exists u (v = y + u \cdot x \wedge u = p^{|y|}))$$

•

$$u = p^{|y|} \quad :\equiv \quad u \text{ is power of } p \\ \wedge y < u \wedge u \leq p \cdot u$$

example:

$$456 < 1000 < 10 \cdot 456$$

•

$$u \leq y \quad :\equiv \quad \exists r \, u = y + r$$

$$u < y \quad :\equiv \quad u \leq y \wedge \sim u = y$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

•

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

•

$$\psi(\varepsilon) = 0$$

• extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

• expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

obviously using decomposition lemma for base p numbers. Top down:

•

$$v = [x, y] \quad :\equiv \quad (y = 0 \wedge v = x) \vee (\sim y = 0 \wedge \exists u (v = y + u \cdot x \wedge u = p^{|y|}))$$

•

$$u = p^{|y|} \quad :\equiv \quad u \text{ is power of } p \\ \wedge y < u \wedge u \leq p \cdot u$$

example:

$$456 < 1000 < 10 \cdot 456$$

•

$$u \leq y \quad :\equiv \quad \exists r \ u = y + r$$

$$u < v \quad :\equiv \quad u < v \wedge \sim u = v$$

•

$$u \text{ is power of } p \quad :\equiv \quad \forall r \ \forall s \ (r \cdot s = u \rightarrow r = 1 \vee p|r)$$

$$p|r \quad :\equiv \quad \exists s \ r = \bar{p} \cdot s$$

coding strings in numbers Let

$$p > \#A + \#Z + 2 \quad \text{prime number}$$

Interpret $w \in (A \cup Z \cup \{\$\})^*$ as number representation to base p .

•

$$\psi : A \cup Z \cup \{\$\} \rightarrow [1 : p - 1]$$

$$\psi(a) \in \mathbb{N} \quad \text{injective}$$

•

$$\psi(\varepsilon) = 0$$

• extend to

$$\psi : (A \cup Z \cup \{\$\})^* \rightarrow [1 : p - 1]$$

$$\psi(w[s - 1 : 0]) = \sum_{i=1}^{s-1} \psi(w_i) \cdot p^i$$

• expression whose value codes $a \in A \cup Z \cup \{\$\}$

$$\hat{a} = \overline{\psi(a)} \in T$$

expressing concatenation

Lemma 6. *There is an arithmetic predicate $v = [x, y]$ such that for all $V, X, Y \in (A \cup Z \cup \{\$\})^*$ holds*

$$\psi(V) = [\psi(X), \psi(Y)] \Leftrightarrow V = XY$$

obviously using decomposition lemma for base p numbers. Top down:

•

$$v = [x, y] \quad :\equiv \quad (y = 0 \wedge v = x) \vee (\sim y = 0 \wedge \exists u (v = y + u \cdot x \wedge u = p^{|y|}))$$

•

$$u = p^{|y|} \quad :\equiv \quad u \text{ is power of } p \\ \wedge y < u \wedge u \leq p \cdot u$$

example:

$$456 < 1000 < 10 \cdot 456$$

•

$$u \leq y \quad :\equiv \quad \exists r \, u = y + r \\ u < v \quad :\equiv \quad u < v \wedge \sim u = v$$

•

$$u \text{ is power of } p \quad :\equiv \quad \forall r \, \forall s (r \cdot s = u \rightarrow r = 1 \vee p|r) \\ p|r \quad :\equiv \quad \exists s \, r = \bar{p} \cdot s$$

concatenating several strings

$$u = [xyv] \quad :\equiv \quad \exists w (w = [x, y] \wedge u = [w, v])$$

etc.

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots Bz_0vB \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(1): start configuration of computation

•

$$\begin{aligned} & \exists k \exists m \exists a \exists b \exists c \\ (w &= [\hat{\$}, k, \hat{\$}, m] \\ & \wedge k = [a, b, c] \\ & \wedge a = \psi(B \dots B) \\ & \wedge c = \psi(B \dots B) \\ & b = \psi(z_0v_{n-1} \dots v_0)) \end{aligned}$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots Bz_0vB \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(1): start configuration of computation

•

$$\begin{aligned} & \exists k \exists m \exists a \exists b \exists c \\ (w &= [\hat{\$}, k, \hat{\$}, m] \\ & \wedge k = [a, b, c] \\ & \wedge a = \psi(B \dots B) \\ & \wedge c = \psi(B \dots B) \\ & b = \psi(z_0 v_{n-1} \dots v_0)) \end{aligned}$$

- $\sigma(d,a)$: d codes a single symbol in $\psi^{-1}(a)$

$$\sigma(d,a) :\equiv \exists e \exists f (a = [e,d,f] \wedge d < \overline{p} \wedge \sim d = 0)$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots B z_0 v B \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(1): start configuration of computation

•

$$\begin{aligned} & \exists k \exists m \exists a \exists b \exists c \\ (w &= [\hat{\$}, k, \hat{\$}, m] \\ & \wedge k = [a, b, c] \\ & \wedge a = \psi(B \dots B) \\ & \wedge c = \psi(B \dots B) \\ & b = \psi(z_0 v_{n-1} \dots v_0)) \end{aligned}$$

- $\sigma(d,a)$: d codes a single symbol in $\psi^{-1}(a)$

$$\sigma(d,a) :\equiv \exists e \exists f (a = [e,d,f] \wedge d < \overline{p} \wedge \sim d = 0)$$

•

$$a = \psi(B \dots B) \quad :\equiv \forall d (\sigma(d,a) \rightarrow d = \hat{B})$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

1. $k_0 = B \dots Bz_0vB \dots B$
2. $k_i \vdash k_{i+1}$ for $i < t$
3. $|k_i| = |k_j|$ for all i, j
4. in no k_i if a $z \in Z$ first or last element
5. k_t is endconfiguraion

(1): start configuration of computation

•

$$\begin{aligned} & \exists k \, \exists m \exists a \, \exists b \exists c \\ (w &= [\hat{\$}, k, \hat{\$}, m] \\ &\wedge k = [a, b, c] \\ &\wedge a = \psi(B \dots B) \\ &\wedge c = \psi(B \dots B) \\ &b = \psi(z_0 v_{n-1} \dots v_0)) \end{aligned}$$

- $\sigma(d,a)$: d codes a single symbol in $\psi^{-1}(a)$

$$\sigma(d,a) :\equiv \exists e \, \exists f \, (a = [e,d,f] \, \wedge d < \overline{p} \wedge \sim d = 0)$$

•

$$a = \psi(B \dots B) \quad :\equiv \forall d \, (\sigma(d,a) \rightarrow d = \hat{B})$$

- Horner scheme for $\psi(z_0v)$

$$\begin{aligned} b = \psi(z_0v) &:\equiv \exists y_0 \dots \exists y_s \\ (y_s &= \widehat{z_0} \wedge \\ y_{s-1} &= \overline{p} \cdot y_s + \widehat{v_{s-1}} \wedge \\ &\dots \\ y_0 &= \overline{p} \cdot y_1 + \widehat{v_0} \wedge \\ b &= y_0) \end{aligned}$$

Consider 1-tape TM

$$M_u = (Z, A, \delta, z_0, E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

$$1. \quad k_0 = B \dots B z_0 v B \dots B$$

$$2. \quad k_i \vdash k_{i+1} \text{ for } i < t$$

$$3. \quad |k_i| = |k_j| \text{ for all } i, j$$

$$4. \quad \text{in no } k_i \text{ if a } z \in Z \text{ first or last element}$$

$$5. \quad k_t \text{ is endconfiguraion}$$

(2) \wedge (3): successor configuration

- in (codes of) neighboring configurations x and y we have ' $x \vdash y$ '

$$\begin{aligned} \forall r \forall x \forall y \forall z \quad (& (w = [r, \hat{\$}, x, \hat{\$}y, \hat{\$}, z] \wedge 'x \text{ codes no } \$' \wedge 'y \text{ codes no } \$') \\ & \rightarrow 'x \vdash y') \end{aligned}$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots Bz_0vB \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(2) \wedge (3): successor configuration

- in (codes of) neighboring configurations x and y we have $'x \vdash y'$
$$\forall r \forall x \forall y \forall z ((w = [r, \hat{\$}, x, \hat{\$}y, \hat{\$}, z] \wedge 'x \text{ codes no } \$' \wedge 'y \text{ codes no } \$') \rightarrow 'x \vdash y')$$

• $'x \text{ codes no } \$' :\equiv \quad \forall d (\sigma(d,x) \rightarrow \sim d = \hat{\$})$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

$$1. \quad k_0 = B \dots Bz_0vB \dots B$$

$$2. \quad k_i \vdash k_{i+1} \text{ for } i < t$$

$$\underline{3. \quad |k_i| = |k_j| \text{ for all } i,j}$$

$$4. \quad \text{in no } k_i \text{ if a } z \in Z \text{ first or last element}$$

$$5. \quad k_t \text{ is endconfiguraion}$$

(2) \wedge (3): successor configuration

- in (codes of) neighboring configurations x and y we have $'x \vdash y'$

$$\begin{aligned} \forall r \forall x \, \forall y \, \forall z \, ((w &= [r,\hat{\$},x,\hat{\$}y,\hat{\$},z] \wedge 'x \text{ codes no } \$' \wedge 'y \text{ codes no } \$') \\ &\rightarrow 'x \vdash y') \end{aligned}$$

-

$$'x \text{ codes no } \$' \, :\equiv \, \forall d \, (\sigma(d,x) \rightarrow \sim d = \hat{\$})$$

- next configuration

$$\begin{aligned} 'x \vdash y' \, &:\equiv \, \exists u \, \exists v \\ &(\bigvee_{z,q \in Z, \, a,c \in A, \, \delta(z,a)=(q,c,N)} x = [u,\hat{z},\hat{a},v] \wedge y = [u,\hat{q},\hat{c},v] \\ &\vee \bigvee_{z,q \in Z, \, a,c \in A, \, \delta(z,a)=(q,c,R)} x = [u,\hat{z},\hat{a},v] \wedge y = [u,\hat{c},\hat{q},v] \\ &\vee \bigvee_{z,q \in Z, \, a,b,c \in A, \, \delta(z,a)=(q,c,L)} x = [u,\hat{b},\hat{z},\hat{a},v] \wedge y = [u,\hat{q},\hat{b},\hat{c},v]) \end{aligned}$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots Bz_0vB \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(4): no state at border

$$\begin{aligned} & \forall x \forall y \forall v \\ ((w &= [x, \hat{\$}, y, \hat{\$}, v] \wedge 'y \text{ codes no } \$' \\ & \wedge \forall d (\sigma(d,y) \wedge (\bigvee_{z \in Z} d = \hat{z}))) \\ \rightarrow & \exists e \exists f (y = [e,d,f] \wedge \sim e = 0 \wedge \sim f = 0)) \end{aligned}$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- 1. $k_0 = B \dots Bz_0vB \dots B$
- 2. $k_i \vdash k_{i+1}$ for $i < t$
- 3. $|k_i| = |k_j|$ for all i, j
- 4. in no k_i if a $z \in Z$ first or last element
- 5. k_t is endconfiguraion

(4): no state at border

$$\begin{aligned} & \forall x \forall y \forall v \\ ((w &= [x, \hat{\$}, y, \hat{\$}, v] \wedge 'y \text{ codes no } \$' \\ & \wedge \forall d (\sigma(d,y) \wedge (\bigvee_{z \in Z} d = \hat{z}))) \\ \rightarrow & \exists e \exists f (y = [e, d, f] \wedge \sim e = 0 \wedge \sim f = 0)) \end{aligned}$$

(5) end state is reached

$$\exists e \exists f \exists t w = [e, t, f] \wedge (\bigvee_{z \in E} t = \hat{z})$$

Consider 1-tape TM

$$M_u = (Z,A,\delta,z_0,E)$$

and $v \in \mathbb{B}^*$. M_u started with v halts iff

$$\exists w. w = \$k_0\$ \dots \$k_t\$ \quad , \quad w \in (A \cup Z \cup \{\$\})^+$$

with

- $k_0 = B \dots Bz_0vB \dots B$
- $k_i \vdash k_{i+1}$ for $i < t$
- $|k_i| = |k_j|$ for all i, j
- in no k_i if a $z \in Z$ first or last element
- k_t is endconfiguraion

(4): no state at border

$$\begin{aligned} & \forall x \, \forall y \, \forall v \\ ((w &= [x, \hat{\$}, y, \hat{\$}, v] \wedge 'y \text{ codes no } \$' \\ & \wedge \forall d \, (\sigma(d, y) \wedge (\bigvee_{z \in Z} d = \hat{z}))) \\ \rightarrow & \exists e \, \exists f \, (y = [e, d, f] \wedge \sim e = 0 \wedge \sim f = 0)) \end{aligned}$$

(5) end state is reached

$$\exists e \, \exists f \, \exists t \, w = [e, t, f] \wedge (\bigvee_{z \in E} t = \hat{z})$$

reduction:

$$H(u\#v) :\equiv \exists w \, (1) \wedge (2) \wedge (3) \wedge (4) \wedge (5)$$

3 Towards the second incompleteness theorem

3.1 Preliminaries

For proof systems S denote by $\text{consis}(S)$ the statement, that S is consistent.

Lemma 7. *Consistency can be formulated in Z_E*

$$\text{consis}(S) \in L_E$$

3 Towards the second incompleteness theorem

3.1 Preliminaries

For proof systems S denote by $\text{consis}(S)$ the statement, that S is consistent.

Lemma 7. *Consistency can be formulated in Z_E*

$$\text{consis}(S) \in L_E$$

Proof. Machine M_m started with empty tape enumerates all proofs of S and records all proven statements p . If it finds proofs of p and $\sim p$ for some p it halts. Then

$$\text{consis}(S) :\equiv H(m\#)$$

□

3 Towards the second incompleteness theorem

3.1 Preliminaries

For proof systems S denote by $\text{consis}(S)$ the statement, that S is consistent.

Lemma 7. *Consistency can be formulated in Z_E*

$$\text{consis}(S) \in L_E$$

Proof. Machine M_m started with empty tape enumerates all proofs of S and records all proven statements p . If it finds proofs of p and $\sim p$ for some p it halts. Then

$$\text{consis}(S) :\equiv H(m\#)$$

□

Lemma 8. *If machine M_u started with v halts, then one can prove this in Z_e .*

$$H(u\#v) \text{ true} \Rightarrow S \vdash H(u\#v)$$

Proof. One proves this by writing down a sequence of configurations and checking that it is the computation on M_u started with v . This argument can be formalized in Z_E . □

3 Towards the second incompleteness theorem

3.1 Preliminaries

For proof systems S denote by $\text{consis}(S)$ the statement, that S is consistent.

Lemma 7. *Consistency can be formulated in Z_E*

$$\text{consis}(S) \in L_E$$

Proof. Machine M_m started with empty tape enumerates all proofs of S and records all proven statements p . If it finds proofs of p and $\sim p$ for some p it halts. Then

$$\text{consis}(S) :\equiv H(m\#)$$

□

Lemma 8. *If machine M_u started with v halts, then one can prove this in Z_e .*

$$H(u\#v) \text{ true} \Rightarrow S \vdash H(u\#v)$$

Proof. One proves this by writing down a sequence of configurations and checking that it is the computation on M_u started with v . This argument can be formalized in Z_E . □

Lemma 9. *If p is provable in Z_E , then one can prove, that it is provable*

$$S \vdash p \Rightarrow (S \vdash (S \vdash p))$$

Proof. One establishes the existence of a proof by writing down the proof and then checking (with a syntax check), that the proof rules are obeyed. This argument can be formalized in Z_e □

3.2 An explicit statement, which is independent of Z_E

def: independence Let $S = (\Sigma, L, A, R)$ be a proof system. A statement $v \in L$ is called independent of S if it can be neither proved nor disproved in S .

$$\sim S \vdash v \wedge \sim S \vdash \sim v$$

S is incomplete iff independent p exists. From now on assume that S extends Z_E , i.e. it is at least as powerful as Z_E .

$$Z_E \subseteq S$$

3.2 An explicit statement, which is independent of Z_E

def: independence Let $S = (\Sigma, L, A, R)$ be a proof system. A statement $v \in L$ is called independent of S if it can be neither proved nor disproved in S .

$$\sim S \vdash v \wedge \sim S \vdash \sim v$$

S is incomplete iff independent p exists. From now on assume that S extends Z_E , i.e. it is at least as powerful as Z_E .

$$Z_E \subseteq S$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

Q started with its own program q halts
as soon as it finds a proof
that it does not halt

3.2 An explicit statement, which is independent of Z_E

def: independence Let $S = (\Sigma, L, A, R)$ be a proof system. A statement $v \in L$ is called independent of S if it can be neither proved nor disproved in S .

$$\sim S \vdash v \wedge \sim S \vdash \sim v$$

S is incomplete iff independent p exists. From now on assume that S extends Z_E , i.e. it is at least as powerful as Z_E .

$$Z_E \subseteq S$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

Q started with its own program q halts
as soon as it finds a proof
that it does not halt

- assume $S \vdash H(q\#q)$ then

$$\Rightarrow \sim S \vdash \sim H(q\#q) \quad (S \text{ correct})$$

$$\Rightarrow \sim H(q\#q) \quad (\text{construction of } Q)$$

$$\Rightarrow S \text{ not correct}$$

3.2 An explicit statement, which is independent of Z_E

def: independence Let $S = (\Sigma, L, A, R)$ be a proof system. A statement $v \in L$ is called independent of S if it can be neither proved nor disproved in S .

$$\sim S \vdash v \wedge \sim S \vdash \sim v$$

S is incomplete iff independent p exists. From now on assume that S extends Z_E , i.e. it is at least as powerful as Z_E .

$$Z_E \subseteq S$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

• assume $S \vdash \sim H(q\#q)$ then

$\Rightarrow Q$ finds proof

$\Rightarrow H(q\#q)$ (construction of Q)

$\Rightarrow S$ not correct

Q started with its own program q halts
as soon as it finds a proof
that it does not halt

• assume $S \vdash H(q\#q)$ then

$\Rightarrow \sim S \vdash \sim H(q\#q)$ (S correct)

$\Rightarrow \sim H(q\#q)$ (construction of Q)

$\Rightarrow S$ not correct

3.2 An explicit statement, which is independent of Z_E

def: independence Let $S = (\Sigma, L, A, R)$ be a proof system. A statement $v \in L$ is called independent of S if it can be neither proved nor disproved in S .

$$\sim S \vdash v \wedge \sim S \vdash \sim v$$

S is incomplete iff independent p exists. From now on assume that S extends Z_E , i.e. it is at least as powerful as Z_E .

$$Z_E \subseteq S$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

- assume $S \vdash H(q\#q)$ then

$$\begin{aligned} \Rightarrow & \sim S \vdash \sim H(q\#q) \quad (S \text{ correct}) \\ \Rightarrow & \sim H(q\#q) \quad (\text{construction of } Q) \\ \Rightarrow & S \text{ not correct} \end{aligned}$$

- assume $S \vdash \sim H(q\#q)$ then

$$\begin{aligned} \Rightarrow & Q \text{ finds proof} \\ \Rightarrow & H(q\#q) \quad (\text{construction of } Q) \\ \Rightarrow & S \text{ not correct} \end{aligned}$$

Lemma 11. *Statement $H(q\#q)$ is false*

- assume M_q started with q halts,
- lemma 8: $H(q\#q)$ is provable
- impossible by lemma 10

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 12. *If S is consistent, then statement $\sim H(q\#q)$ is not provable in S .*

$$\text{consis}(S) \Rightarrow \sim S \vdash \sim H(q\#q)$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 12. *If S is consistent, then statement $\sim H(q\#q)$ is not provable in S .*

$$\text{consis}(S) \Rightarrow \sim S \vdash \sim H(q\#q)$$

asssume $S \vdash \sim H(q\#q)$. Then

$\Rightarrow Q$ finds proof and halts

$\Rightarrow H(q\#q)$

$\Rightarrow S \vdash H(q\#q)$ (lemma 8)

$\Rightarrow \sim \text{consis}(S)$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 12. *If S is consistent, then statement $\sim H(q\#q)$ is not provable in S .*

$$\text{consis}(S) \Rightarrow \sim S \vdash \sim H(q\#q)$$

asssume $S \vdash \sim H(q\#q)$. Then

$\Rightarrow Q$ finds proof and halts

$\Rightarrow H(q\#q)$

$\Rightarrow S \vdash H(q\#q)$ (lemma 8)

$\Rightarrow \sim \text{consis}(S)$

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

Proof. The proof of lemma 12 can be formalized in Z_E . □

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

3.4 The second incompleteness theorem

Lemma 14. *If $Z_E \subseteq S$ and S is consistent, then one cannot prove the consistency of S in S*

$$\text{consis}(S) \rightarrow \sim S \vdash \text{consis}(S)$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

3.4 The second incompleteness theorem

Lemma 14. *If $Z_E \subseteq S$ and S is consistent, then one cannot prove the consistency of S in S*

$$\text{consis}(S) \rightarrow \sim S \vdash \text{consis}(S)$$

- assume

$$S \vdash \underbrace{\text{consis}(S)}_A$$

- lemma 13:

$$S \vdash (\text{consis}(S) \rightarrow \underbrace{\sim S \vdash \sim H(q\#q)}_B)$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

3.4 The second incompleteness theorem

Lemma 14. *If $Z_E \subseteq S$ and S is consistent, then one cannot prove the consistency of S in S*

$$\text{consis}(S) \rightarrow \sim S \vdash \text{consis}(S)$$

- assume

$$S \vdash \underbrace{\text{consis}(S)}_A$$

- lemma 13:

$$S \vdash (\text{consis}(S) \rightarrow \underbrace{\sim S \vdash \sim H(q\#q)}_B)$$

- modus ponens

$$\frac{A, A \rightarrow B}{B}$$

-

$$S \vdash \underbrace{(\sim S \vdash \sim H(q\#q))}_B$$

- construction of Q :

$$B \rightarrow \underbrace{\sim H(q\#q)}_C$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

3.4 The second incompleteness theorem

Lemma 14. *If $Z_E \subseteq S$ and S is consistent, then one cannot prove the consistency of S in S*

$$\text{consis}(S) \rightarrow \sim S \vdash \text{consis}(S)$$

- assume

$$S \vdash \underbrace{\text{consis}(S)}_A$$

- lemma 13:

$$S \vdash (\text{consis}(S) \rightarrow \underbrace{\sim S \vdash \sim H(q\#q)}_B)$$

- modus ponens

$$\frac{A, A \rightarrow B}{B}$$

-

$$S \vdash \underbrace{(\sim S \vdash \sim H(q\#q))}_B$$

- construction of Q :

$$B \rightarrow \underbrace{\sim H(q\#q)}_C$$

- this argument can be formalized in Z_E

$$S \vdash B \rightarrow C$$

a statement: Consider Turing machine $Q = M_q$. Started with w it enumerates all provable statements of S , i.e. the set

$$P = \{v : S \vdash v\}$$

It halts as soon as it finds a proof of $\sim H(w\#w)$.

Lemma 10. *If S is correct, then statement $H(q\#q)$ is independent of S .*

3.3 weakening correctness to consistency

Lemma 13. *Lemma 12 is provable in S*

$$S \vdash (\text{consis}(S) \rightarrow \sim S \vdash \sim H(q\#q))$$

3.4 The second incompleteness theorem

Lemma 14. *If $Z_E \subseteq S$ and S is consistent, then one cannot prove the consistency of S in S*

$$\text{consis}(S) \rightarrow \sim S \vdash \text{consis}(S)$$

- assume

$$S \vdash \underbrace{\text{consis}(S)}_A$$

- lemma 13:

$$S \vdash (\text{consis}(S) \rightarrow \underbrace{\sim S \vdash \sim H(q\#q)}_B)$$

- modus ponens

$$\frac{A, A \rightarrow B}{B}$$

-

$$S \vdash \underbrace{(\sim S \vdash \sim H(q\#q))}_B$$

- construction of Q :

$$B \rightarrow \underbrace{\sim H(q\#q)}_C$$

- this argument can be formalized in Z_E

$$S \vdash B \rightarrow C$$

- modus ponens

$$\frac{B, B \rightarrow C}{C}$$

-

$$S \vdash \sim H(q\#q)$$

- lemma 9 (provability is provable)

$$S \vdash \underbrace{(S \vdash \sim H(q\#q))}_{\sim B}$$

S not consistent

warning

before you try to find such proofs yourself
for extended periods of time:

read vita of Cantor and Gödel in Wikipedia

there might be a mental health hazard

4 A non standard natural number

theories (as considered here) have

- a universe U . Here natural numbers.
- functions f with arguments and values in U , Here $+$, \times
- theory is *countable* if the set of axioms and proof rules is countable

4 A non standard natural number

theories (as considered here) have

- a universe U . Here natural numbers.
- functions f with arguments and values in U , Here $+$, \times
- theory is *countable* if the set of axioms and proof rules is countable

model of a theory : instantiation of universe U and functions f such that the axioms hold. For Z_E

- decimal numbers
- binary numbers
- unary numbers: finite sequences of vertical lines |
- all *isomorphic* elements of U in different models are just renamings of each other
- standard natural numbers

4 A non standard natural number

theories (as considered here) have

- a universe U . Here natural numbers.
- functions f with arguments and values in U , Here $+$, \times
- theory is *countable* if the set of axioms and proof rules is countable

model of a theory : instantiation of universe U and functions f such that the axioms hold. For Z_E

- decimal numbers
- binary numbers
- unary numbers: finite sequences of vertical lines |
- all *isomorphic* elements of U in different models are just renamings of each other
- standard natural numbers

do non standard models of Z_E exist?

4 A non standard natural number

theories (as considered here) have

- a universe U . Here natural numbers.
- functions f with arguments and values in U , Here $+$, \times
- theory is *countable* if the set of axioms and proof rules is countable

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

model of a theory : instantiation of universe U and functions f such that the axioms hold. For Z_E

- decimal numbers
- binary numbers
- unary numbers: finite sequences of vertical lines |
- all *isomorphic* elements of U in different models are just renamings of each other
- standard natural numbers

do non standard models of Z_E exist?

4 A non standard natural number

theories (as considered here) have

- a universe U . Here natural numbers.
- functions f with arguments and values in U , Here $+$, \times
- theory is *countable* if the set of axioms and proof rules in countable

model of a theory : instantiation of universe U and functions f such that the axioms hold. For Z_E

- decimal numbers
- binary numbers
- unary numbers: finite sequences of vertical lines |
- all *isomorphic* elements of U in different models are just renamings of each other
- standard natural numbers

do non standard models of Z_E exist?

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

- define α as the number of steps after which Q started with q halts.
- does not exist, because $H(q\#q)$ is false
- the statement that it exists is consistent.
- adding this statement to axioms of Z_E gives a consistent theory.
- has model by model existence theorem.
- it contains this 'step number'

5 A countable model of set theory and the real numbers

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

5 A countable model of set theory and the real numbers

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

- established proof systems for set theory (including the reals) exist
- e.g. ZF: Zermelo-Fraenkel set theory
- countable!
- so there exist countable models

5 A countable model of set theory and the real numbers

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

- established proof systems for set theory (including the reals) exist
- e.g. ZF: Zermelo-Fraenkel set theory
- countable!
- so there exist countable models

how can this be?

5 A countable model of set theory and the real numbers

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

- established proof systems for set theory (including the reals) exist
- e.g. ZF: Zermelo-Fraenkel set theory
- countable!
- so there exist countable models

how can this be?

- in a language L you can only define countably many entities
- include in model only the ones you can define/talk about

5 A countable model of set theory and the real numbers

model existence theorem of logic:

Lemma 15. *Every consistent countable theory has a countable model.*

- established proof systems for set theory (including the reals) exist
- e.g. ZF: Zermelo-Fraenkel set theory
- countable!
- so there exist countable models

how can this be?

- in a language L you can only define countably many entities
- include in model only the ones you can define/talk about

- question: do things you cannot talk about, really exist?