

The Recursive Functions

early 20th century

mankind trying to grasp what is computability

1 What functions are computable in the intuitive sense?

attempts at definitions

- primitive recursive functions, didn't work; extend to μ -recursive functions
- Turing machines (*extremely* simple) you have not seen them yet
- MIPS with infinite registers and memory (from I2EA)
- C with infinite data types and memory (from I2EA)
- more

1 What functions are computable in the intuitive sense?

attempts at definitions

- primitive recursive functions, didn't work; extend to μ -recursive functions
- Turing machines (*extremely* simple) you have not seen them yet
- MIPS with infinite registers and memory (from I2EA)
- C with infinite data types and memory (from I2EA)
- more

all were shown equivalent

- the interesting case shown by Church: Turing machines are simulated by μ -recursive functions
- *Church's thesis*: the above definition(s) capture the intuitive concept of computability
- all sufficiently powerful programming languages simulate each other
- if you know one (sufficiently powerful) programming language: you know them all. Just find the constructs you are used to in the new language.

or implement them

2 prelude: countability

def: countable set A set A is countable if A is finite or if there is a bijection

$$b : \mathbb{N}_0 \rightarrow A$$

Note that the infinite sequence

$$(b(n)) = (b(0), b(1), b(2), \dots)$$

enumerates A .

example:

$$A = \{n \in \mathbb{N}_0 : n \text{ even}\} \quad , \quad b(n) = 2n$$

2 prelude: countability

def: countable set A set A is countable if A is finite or if there is a bijection

$$b : \mathbb{N}_0 \rightarrow A$$

Note that the infinite sequence

$$(b(n)) = (b(0), b(1), b(2), \dots)$$

enumerates A .

example:

$$A = \{n \in \mathbb{N}_0 : n \text{ even}\} \quad , \quad b(n) = 2n$$

Lemma 1. *If A is finite, then A^* is countable*

enumerate

- A^0, A^1, A^2, \dots
- each A^n in lexicographic order

$$\mathbb{B}^* = (\{\epsilon, 0, 1, 00, 01, 10, 11, 000, \dots\})$$

2 prelude: countability

def: countable set A set A is countable if A is finite or if there is a bijection

$$b : \mathbb{N}_0 \rightarrow A$$

Note that the infinite sequence

$$(b(n)) = (b(0), b(1), b(2), \dots)$$

enumerates A .

example:

$$A = \{n \in \mathbb{N}_0 : n \text{ even}\} \quad , \quad b(n) = 2n$$

Lemma 1. *If A is finite, then A^* is countable*

enumerate

- A^0, A^1, A^2, \dots
- each A^n in lexicographic order

$$\mathbb{B}^* = (\{\epsilon, 0, 1, 00, 01, 10, 11, 000, \dots\})$$

Lemma 2. $\mathbb{N}_0 \times \mathbb{N}_0$ is countable

enumerate the sets

- $S_i = \{(a, b) : a + b = i\}$ for $i = 0, 1, 2, \dots$
- each S_i in lexicographic order

$$\mathbb{N}_0 \times \mathbb{N}_0 = \{(0, 0), (0, 1), (1, 0), (0, 2), (1, 1), (2, 0), \dots\}$$

2 prelude: countability

def: countable set A set A is countable if A is finite or if there is a bijection

$$b : \mathbb{N}_0 \rightarrow A$$

Note that the infinite sequence

$$(b(n)) = (b(0), b(1), b(2), \dots)$$

enumerates A .

example:

$$A = \{n \in \mathbb{N}_0 : n \text{ even}\} \quad , \quad b(n) = 2n$$

Lemma 1. *If A is finite, then A^* is countable*

enumerate

- A^0, A^1, A^2, \dots
- each A^n in lexicographic order

$$\mathbb{B}^* = (\{\epsilon, 0, 1, 00, 01, 10, 11, 000, \dots\})$$

Lemma 2. $\mathbb{N}_0 \times \mathbb{N}_0$ is countable

enumerate the sets

- $S_i = \{(a, b) : a + b = i\}$ for $i = 0, 1, 2, \dots$
- each S_i in lexicographic order

$$\mathbb{N}_0 \times \mathbb{N}_0 = \{(0, 0), (0, 1), (1, 0), (0, 2), (1, 1), (2, 0), \dots\}$$

notation: for pairs $p = (x, y)$ we denote - as for vectors - its components as

$$p_1 = x, p_2 = y \quad \text{maybe helpful for exercises}$$

Lemma 3. *If A and B are countable, then $A \times B$ is countable*

Proof. exercise □

Lemma 4. *For all k holds: \mathbb{N}^k is countable.*

proof: exercise

3 a first glance (in this lecture) at diagonalisation

Lemma 5. *Let*

$$F = \{f \mid f: \mathbb{N}_0 \rightarrow \mathbb{B}\}$$

then F is not countable.

3 a first glance (in this lecture) at diagonalisation

Lemma 5. *Let*

$$F = \{f \mid f: \mathbb{N}_0 \rightarrow \mathbb{B}\}$$

then F is not countable.

- assume F can be enumerated as

$$F = \{f_0, f_1, f_2, \dots\}$$

- consider the infinite matrix of table 1

$f_0(0)$	$f_0(1)$	$f_0(2)$	\dots	$f_0(x)$	\dots
$f_1(0)$	$f_1(1)$	$f_1(2)$	\dots	$f_1(x)$	\dots
$f_2(0)$	$f_2(1)$	$f_2(2)$	\dots	$f_2(x)$	\dots
			\dots		
$f_x(0)$	$f_x(1)$	$f_x(2)$	\dots	$f_x(x)$	\dots
			\dots		

Table 1: row x of this matrix is the function table of function f_x . Function g is defined such that it differs from f_x at argument x , i.e. on the diagonal of the matrix.

3 a first glance (in this lecture) at diagonalisation

Lemma 5. *Let*

$$F = \{f \mid f : \mathbb{N}_0 \rightarrow \mathbb{B}\}$$

then F is not countable.

- assume F can be enumerated as

$$F = \{f_0, f_1, f_2, \dots\}$$

- consider the infinite matrix of table 1

$f_0(0)$	$f_0(1)$	$f_0(2)$	\dots	$f_0(x)$	\dots
$f_1(0)$	$f_1(1)$	$f_1(2)$	\dots	$f_1(x)$	\dots
$f_2(0)$	$f_2(1)$	$f_2(2)$	\dots	$f_2(x)$	\dots
			\dots		
$f_x(0)$	$f_x(1)$	$f_x(2)$	\dots	$f_x(x)$	\dots
			\dots		

Table 1: row x of this matrix is the function table of function f_x . Function g is defined such that it differs from f_x at argument x , i.e. on the diagonal of the matrix.

- define $g : \mathbb{N}_0 \rightarrow \mathbb{B}$ such that it differs from f_x at argument x

$$\forall x. \quad g(x) = f_x(x) \oplus 1$$

- then $g \notin \{f_0, f_1, f_2, \dots\}$: otherwise $g = f_x$ for some x and we get contradiction

$$f_x(x) = g(x) = f_x(x) \oplus 1$$

4 definition of primitive recursive functions

you have seen it
where?

Inductive definition of a set PR of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

4 definition of primitive recursive functions

you have seen it
where?

Inductive definition of a set PR of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

base cases:

1. constant functions $c_s^r \in PR$ where

$$c_s^r : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$c_s^r(x) = s, s \in \mathbb{N}_0$$

2. projections $p_i^r \in PR$ where

$$p_i^r(x) : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$p_i^r(x) = x_i$$

3. successor function $S \in PR$

$$s : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

$$S(x) = x + 1$$

where have you seen it first?

4 definition of primitive recursive functions

you have seen it
where?

Inductive definition of a set PR of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

base cases:

1. constant functions $c_s^r \in PR$ where

$$c_s^r : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$c_s^r(x) = s, s \in \mathbb{N}_0$$

2. projections $p_i^r \in PR$ where

$$p_i^r(x) : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$p_i^r(x) = x_i$$

3. successor function $S \in PR$

$$s : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

$$S(x) = x + 1$$

where have you seen it first?

induction steps of definition:

4. function composition. If the following function are all in PR

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in PR$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in PR

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in PR$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$\begin{aligned} f(0, x) &= g(x) \\ f(n+1, x) &= h(n, f(n, x), x) \end{aligned}$$

6. these are all

excluding everything else

1. constant functions $c_s^r \in PR$ where

$$c_s^r : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$c_s^r(x) = s, s \in \mathbb{N}_0$$

2. projections $p_i^r \in PR$ where

$$p_i^r(x) : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$p_i^r(x) = x_i$$

3. successor function $S \in PR$

$$s : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

4. function composition. If the following function are all in PR

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in PR$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in PR

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in PR$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$f(0, x) = g(x)$$

$$f(n+1, x) = h(n, f(n, x), x)$$

examples you know

as KIU CS students :)

- addition

$$f(0, x) = x = p_1^1(x)$$

$$f(n+1, x) = S(f(n, x))$$

- multiplication

$$g(0, x) = 0 = c_0^1(x)$$

$$g(n+1, x) = f(x, g(n, x))$$

- exponentiation

$$h(0, x) = 1$$

$$h(n+1, x) = g(n, h(n, x))$$

recursion theory was for you
like mother's milk

1. constant functions $c_s^r \in PR$ where

$$c_s^r : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$c_s^r(x) = s, s \in \mathbb{N}_0$$

2. projections $p_i^r \in PR$ where

$$p_i^r(x) : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

$$p_i^r(x) = x_i$$

3. successor function $S \in PR$

$$s : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

4. function composition. If the following function are all in PR

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in PR$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in PR

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in PR$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$f(0, x) = g(x)$$

$$f(n+1, x) = h(n, f(n, x), x)$$

examples you know

- addition

$$f(0, x) = x = p_1^1(x)$$

$$f(n+1, x) = S(f(n, x))$$

- multiplication

$$g(0, x) = 0 = c_0^1(x)$$

$$g(n+1, x) = f(x, g(n, x))$$

- exponentiation

$$h(0, x) = 1$$

$$h(n+1, x) = g(n, h(n, x))$$

explanation for math students, why...

- in computer architecture you should be able to explain/prove that your adders work
- so why does $1 + 1 = 10$ make sense?
- decimal counter part: $9 + 1 = 10$ **theorem!**
- using
 - Z = number of fingers/toes
 - $9+1 = Z$ **definition**
 - **$10 = 1 \cdot Z^1 + 0 \cdot Z^0$**
- thus need to define exponentiation without decimal numbers

5 the pr functions cannot be all computable functions

a modern view

example for defining +

$$\begin{aligned}f_1(x) &= p_1^1(x) \\f_2(x) &= S(x) \\f_3(0,x) &= f_1(x) \\f_3(n+1,y) &= f_2(f_3(n,x))\end{aligned}$$

5 the pr functions cannot be all computable functions

a modern view

example for defining +

$$\begin{aligned}f_1(x) &= p_1^1(x) \\f_2(x) &= S(x) \\f_3(0,x) &= f_1(x) \\f_3(n+1,y) &= f_2(f_3(n,x))\end{aligned}$$

def: derivation A *derivation* of a pr function f is a finite sequence of definitions of functions

$$(f_1, f_2, \dots, f_s)$$

such that for each i

- either f_i is pr by rules 1 to 3
- or f_i is defined by rule 4 or 5 using only previously listed functions f_j with $j < i$.
- $f = f_s$ is the last function defined in the sequence

5 the pr functions cannot be all computable functions

a modern view

example for defining +

$$\begin{aligned}f_1(x) &= p_1^1(x) \\f_2(x) &= S(x) \\f_3(0,x) &= f_1(x) \\f_3(n+1,y) &= f_2(f_3(n,x))\end{aligned}$$

def: derivation A *derivation* of a pr function f is a finite sequence of definitions of functions

$$(f_1, f_2, \dots, f_s)$$

such that for each i

- either f_i is pr by rules 1 to 3
- or f_i is defined by rule 4 or 5 using only previously listed functions f_j with $j < i$.
- $f = f_s$ is the last function defined in the sequence

observe:

- derivations d are formed by symbols of a finite alphabet A , thus $d \in A^*$
- one can write a parser P , which decides if an input d is a derivation of a function $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$
- enumerate all elements $w \in A^*$ and test each w by the parser P .
- for all x define d_x as the derivation of the x 'th string which passes the test and define

$$f_x : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

as the function defined by derivation d_x .

- this enumerates the pr functions $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$

$$\{f_0, f_1, \dots\} = \{f \mid f : \mathbb{N}_0 \rightarrow \mathbb{N}_0, f \text{ is pr}\}$$

5 the pr functions cannot be all computable functions

a modern view

example for defining +

$$\begin{aligned}f_1(x) &= p_1^1(x) \\f_2(x) &= S(x) \\f_3(0,x) &= f_1(x) \\f_3(n+1,y) &= f_2(f_3(n,x))\end{aligned}$$

def: derivation A *derivation* of a pr function f is a finite sequence of definitions of functions

$$(f_1, f_2, \dots, f_s)$$

such that for each i

- either f_i is pr by rules 1 to 3
- or f_i is defined by rule 4 or 5 using only previously listed functions f_j with $j < i$.
- $f = f_s$ is the last function defined in the sequence

observe:

- derivations d are formed by symbols of a finite alphabet A , thus $d \in A^*$
- one can write a parser P , which decides if an input d is a derivation of a function $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$
- enumerate all elements $w \in A^*$ and test each w by the parser P .
- for all x define d_x as the derivation of the x 'th string which passes the test and define

$$f_x : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

as the function defined by derivation d_x .

- this enumerates the pr functions $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$

$$\{f_0, f_1, \dots\} = \{f \mid f : \mathbb{N}_0 \rightarrow \mathbb{N}_0, f \text{ is pr}\}$$

it's obviously over:

5 the pr functions cannot be all computable functions

observe:

- derivations d are formed by symbols of a finite alphabet A , thus $d \in A^*$
- one can write a parser P , which decides if an input d is a derivation of a function $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$
- enumerate all elements $w \in A^*$ and test each w by the parser P .
- for all x define d_x as the derivation of the x 'th string which passes the test and define

$$f_x : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

as the function defined by derivation d_x .

- this enumerates the pr functions $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$

$$\{f_0, f_1, \dots\} = \{f \mid f : \mathbb{N}_0 \rightarrow \mathbb{N}_0, f \text{ is pr}\}$$

it's obviously over:

- diagonalisation: the function

$$g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

defined by

$$\forall x. g(x) = f_x(x) + 1$$

is not pr. **diagonalisation as before**

- we can write an interpreter I which given a derivation d_x and an input x evaluates $f_x(x)$
- now compute $g(x)$ as i) enumerate words $w \in A^*$ and test each w by parser P until d_x is found. ii) using the interpreter compute $f_x(x)$ iii) then add 1

5 the pr functions cannot be all computable functions

observe:

- derivations d are formed by symbols of a finite alphabet A , thus $d \in A^*$
- one can write a parser P , which decides if an input d is a derivation of a function $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$
- enumerate all elements $w \in A^*$ and test each w by the parser P .
- for all x define d_x as the derivation of the x 'th string which passes the test and define

$$f_x : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

as the function defined by derivation d_x .

- this enumerates the pr functions $f : \mathbb{N}_0 \rightarrow \mathbb{N}_0$

$$\{f_0, f_1, \dots\} = \{f \mid f : \mathbb{N}_0 \rightarrow \mathbb{N}_0, f \text{ is pr}\}$$

it's obviously over:

- diagonalisation: the function

$$g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

defined by

$$\forall x. g(x) = f_x(x) + 1$$

is not pr. **diagonalisation as before**

- we can write an interpreter I which given a derivation d_x and an input x evaluates $f_x(x)$
- now compute $g(x)$ as i) enumerate words $w \in A^*$ and test each w by parser P until d_x is found. ii) using the interpreter compute $f_x(x)$ iii) then add 1

so it's computable

Lemma 6. *There exists a total computable function, which is not primitive recursive*

6 more pr functions

- modified predecessor

$$p(x) = \begin{cases} x-1 & x > 0 \\ 0 & x = 0 \end{cases}$$

$$\begin{aligned} u(0) &= 0 \\ u(x+1) &= x \end{aligned}$$

6 more pr functions

- modified predecessor

$$p(x) = \begin{cases} x-1 & x > 0 \\ 0 & x = 0 \end{cases}$$

$$\begin{aligned} u(0) &= 0 \\ u(x+1) &= x \end{aligned}$$

- modified difference

$$x \dot{-} y = \begin{cases} x-y & x \geq y \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} x \dot{-} 0 &= x \\ x \dot{-} (y+1) &= u(x \dot{-} y) \end{aligned}$$

- sign

$$sg(x) = \begin{cases} 1 & x > 0 \\ 0 & x = 0 \end{cases}$$

$$\begin{aligned} sg(0) &= 0 \\ sg(x+1) &= 1 \end{aligned}$$

alternatively

$$\begin{aligned} \overline{sg} &= 1 \dot{-} x \\ sg(x) &= \overline{sg}(\overline{sg}(x)) \end{aligned}$$

7 pr predicates

def: predicate on N_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

7 pr predicates

def: predicate on N_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

def: characteristic function of predicate P :

$$C_P : \mathbb{N}_0^r \rightarrow \mathbb{B}$$

$$C_P(x) = \begin{cases} 1 & P(x) = true \\ 0 & P(x) = false \end{cases}$$

def: pr predicates Predicate P is pr iff C_P is pr.

7 pr predicates

def: predicate on N_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

def: characteristic function of predicate P :

$$C_P : \mathbb{N}_0^r \rightarrow \mathbb{B}$$

$$C_P(x) = \begin{cases} 1 & P(x) = true \\ 0 & P(x) = false \end{cases}$$

def: pr predicates Predicate P is pr iff C_P is pr.

examples

- $x < y$

$$C_{x < y} = sg(x - y)$$

7 pr predicates

def: predicate on \mathbb{N}_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

def: characteristic function of predicate P :

$$C_P : \mathbb{N}_0^r \rightarrow \mathbb{B}$$

$$C_P(x) = \begin{cases} 1 & P(x) = true \\ 0 & P(x) = false \end{cases}$$

def: pr predicates Predicate P is pr iff C_P is pr.

examples

- $x < y$

$$C_{x < y} = sg(x - y)$$

- if P is pr predicate on \mathbb{N}_0^r and

$$g_1, \dots, g_r : \mathbb{N}^m \rightarrow \mathbb{N}_0$$

are pr, then

$$Q(x) = P(g_1(x), \dots, g_r(x))$$

is pr predicate on \mathbb{N}_0^m

- $2x < y$

def: predicate on \mathbb{N}_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

def: characteristic function of predicate P :

$$C_P : \mathbb{N}_0^r \rightarrow \mathbb{B}$$

$$C_P(x) = \begin{cases} 1 & P(x) = true \\ 0 & P(x) = false \end{cases}$$

def: pr predicates Predicate P is pr iff C_P is pr.

examples

- $x < y$

$$C_{x < y} = sg(x - y)$$

- if P is pr predicate on \mathbb{N}_0^r and

$$g_1, \dots, g_r : \mathbb{N}^m \rightarrow \mathbb{N}_0$$

are pr, then

$$Q(x) = P(g_1(x), \dots, g_r(x))$$

is pr predicate on \mathbb{N}_0^m

- $2x < y$

Lemma 7. If P and Q are pr predicates, then so are

$$P \wedge Q, P \vee Q, \sim P$$

def: predicate on \mathbb{N}_0^r for the time being just a function

$$P : \mathbb{N}_0^r \rightarrow \{true, false\}$$

examples: with $r = 2$

$$x \leq y, x|y, 2x < y$$

def: characteristic function of predicate P :

$$C_P : \mathbb{N}_0^r \rightarrow \mathbb{B}$$

$$C_P(x) = \begin{cases} 1 & P(x) = true \\ 0 & P(x) = false \end{cases}$$

def: pr predicates Predicate P is pr iff C_P is pr.

examples

- $x < y$

$$C_{x < y} = sg(x \dot{-} y)$$

- if P is pr predicate on \mathbb{N}_0^r and

$$g_1, \dots, g_r : \mathbb{N}^m \rightarrow \mathbb{N}_0$$

are pr, then

$$Q(x) = P(g_1(x), \dots, g_r(x))$$

is pr predicate on \mathbb{N}_0^m

- $2x < y$

Lemma 7. If P and Q are pr predicates, then so are

$$P \wedge Q, P \vee Q, \sim P$$

$$\begin{aligned} C_{P \wedge Q}(x) &= C_P(x) \cdot C_Q(x) \\ C_{P \vee Q}(x) &= sg(C_P(x) + C_Q(x)) \\ C_{\sim P}(x) &= \overline{sg}(P(x)) \end{aligned}$$

- $x = y$

$$x = y \leftrightarrow \sim ((x < y) \vee (y < x))$$

- $x \leq y$

$$x \leq y \leftrightarrow x(x < y) \vee (x = y)$$

definition by case split

Lemma 8. *If*

$$P_1, \dots, P_k$$

are pr predicates on \mathbb{N}_0^r and

$$f_1, \dots, f_r : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

are pr functions and for each $x \in \mathbb{N}_0^r$ there is at most one i such that $P_i(x)$ is true. Then

$$g(x) = \begin{cases} f_1(x) & P_1(x) \\ \dots & \\ f_k(x) & P_k(x) \\ 0 & \text{otherwise} \end{cases}$$

is pr.

$$g(x) = C_{P_1}(x) \cdot f_1(x) + \dots + C_{P_k}(x) \cdot f_k(x)$$

definition by case split

Lemma 8. *If*

$$P_1, \dots, P_k$$

are pr predicates on \mathbb{N}_0^r and

$$f_1, \dots, f_k : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

are pr functions and for each $x \in \mathbb{N}_0^r$ there is at most one i such that $P_i(x)$ is true. Then

$$g(x) = \begin{cases} f_1(x) & P_1(x) \\ \dots & \\ f_k(x) & P_k(x) \\ 0 & \text{otherwise} \end{cases}$$

is pr.

$$g(x) = C_{P_1}(x) \cdot f_1(x) + \dots + C_{P_k}(x) \cdot f_k(x)$$

finite functions

Lemma 9. *Let $f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$. If $f(x) \neq 0$ for finitely many x , then f is pr.*

8 bounded μ -operator

def: bounded μ -operator For

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

we define

$$\mu_b f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

by

$$\mu_b f(n, x) = \begin{cases} \min\{m : f(m, x) = 0, m \leq n\} & \text{if it exists} \\ 0 & \text{otherwise} \end{cases}$$

$\mu_b f(n, x)$ returns smallest solution of equation $f(m, x) = 0$ in the bounded range $[0 : n]$ if it exists...

8 bounded μ -operator

def: bounded μ -operator For

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

we define

$$\mu_b f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

by

$$\mu_b f(n, x) = \begin{cases} \min\{m : f(m, x) = 0, m \leq n\} & \text{if it exists} \\ 0 & \text{otherwise} \end{cases}$$

$\mu_b f(n, x)$ returns smallest solution of equation $f(m, x) = 0$ in the bounded range $[0 : n]$ if it exists...

Lemma 10. *If f is pr, then $\mu_b f$ is pr*

$$\begin{aligned} \mu_b f(n, 0) &= 0 \\ \mu_b f(n+1, x) &= \begin{cases} \mu_b f(n, x) & \mu_b f(n, x) \neq 0 \\ n+1 & f(n+1, x) = 0 \wedge \mu_b f(n, x) = 0 \wedge f(0, x) \neq 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

8 bounded μ -operator

def: bounded μ -operator For

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

we define

$$\mu_b f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

by

$$\mu_b f(n, x) = \begin{cases} \min\{m : f(m, x) = 0, m \leq n\} & \text{if it exists} \\ 0 & \text{otherwise} \end{cases}$$

$\mu_b f(n, x)$ returns smallest solution of equation $f(m, x) = 0$ in the bounded range $[0 : n]$ if it exists...

examples:

- $\lfloor x/y \rfloor$

$$\lfloor x/y \rfloor = \min\{m : m \leq x \wedge (m+1)y > x\}$$

- divides: $y|x$

$$rem(x, y) = x - y \cdot \lfloor x/y \rfloor$$

$$C|(x, y) = \overline{sg}(rem(x, y))$$

Lemma 10. *If f is pr, then $\mu_b f$ is pr*

$$\mu_b f(n, 0) = 0$$

$$\mu_b f(n+1, x) = \begin{cases} \mu_b f(n, x) & \mu_b f(n, x) \neq 0 \\ n+1 & f(n+1, x) = 0 \wedge \mu_b f(n, x) = 0 \wedge f(0, x) \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

Lemma 11. *For all $k \leq 2$ and $i \in [1 : k]$ there are pr functions*

$$b^{(k)} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \quad \text{bijective}$$

$$b_i^{(k)} : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

such that for all $x \in \mathbb{N}_0^k$ and all i

$$b_i^{(k)}(b^{(k)}(x_1, \dots, x_k)) = x_i$$

Lemma 11. For all $k \leq 2$ and $i \in [1 : k]$ there are pr functions

$$b^{(k)} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \quad \text{bijective}$$

$$b_i^{(k)} : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

such that for all $x \in \mathbb{N}_0^k$ and all i

$$b_i^{(k)}(b^{(k)}(x_1, \dots, x_k)) = x_i$$

- $k = 2$: enumerate as in figure 1




Figure 1: map $(0,0) \rightarrow 0$, $(1,0) \rightarrow 1$, $(0,1) \rightarrow 2$ etc

- diagonals $1, 2, \dots$ each with positions $0, 1, \dots$

primitive recursive bijections

Lemma 11. For all $k \leq 2$ and $i \in [1 : k]$ there are pr functions

$$b^{(k)} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \quad \text{bijective}$$

$$b_i^{(k)} : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

such that for all $x \in \mathbb{N}_0^k$ and all i

$$b_i^{(k)}(b^{(k)}(x_1, \dots, x_k)) = x_i$$

- $k = 2$: enumerate as in figure 1




Figure 1: map $(0,0) \rightarrow 0$, $(1,0) \rightarrow 1$, $(0,1) \rightarrow 2$ etc

- diagonals $1, 2, \dots$ each with positions $0, 1, \dots$

- then

(n, m) stands on diagonal $n + m + 1$ at position m

-

$$\begin{aligned} b^{(2)}(n, m) &= \sum_{i=1}^{n+m} i + m \\ &= \frac{(n+m)(n+m+1)}{2} + m \quad \text{this is pr} \end{aligned}$$

primitive recursive bijections

Lemma 11. For all $k \leq 2$ and $i \in [1 : k]$ there are pr functions

$$b^{(k)} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \quad \text{bijective}$$

$$b_i^{(k)} : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

such that for all $x \in \mathbb{N}_0^k$ and all i

$$b_i^{(k)}(b^{(k)}(x_1, \dots, x_k)) = x_i$$

- $k = 2$: enumerate as in figure 1




Figure 1: map $(0,0) \rightarrow 0$, $(1,0) \rightarrow 1$, $(0,1) \rightarrow 2$ etc

- diagonals $1, 2, \dots$ each with positions $0, 1, \dots$

- then

(n, m) stands on diagonal $n + m + 1$ at position m

-

$$\begin{aligned} b^{(2)}(n, m) &= \sum_{i=1}^{n+m} i + m \\ &= \frac{(n+m)(n+m+1)}{2} + m \quad \text{this is pr} \end{aligned}$$

- inverse mappings: given

$$b^{(2)}(n, m) = \sum_{i=1}^s i + m = x \quad \text{solve } s = ?$$

$$\begin{aligned} s(x) &= \max\{j : \frac{j(j+1)}{2} \leq x, j \leq x\} \\ &= \min\{j : \frac{j(j+1)}{2} > x, j \leq x\} \quad \text{this is pr} \end{aligned}$$

$$b_2^{(2)}(x) = x - \frac{s(x) \cdot (s(x) + 1)}{2} \quad (= m)$$

$$b_1^{(2)}(x) = s(x) - b_2^{(2)}(x) \quad (= n)$$

Lemma 11. *For all $k \leq 2$ and $i \in [1 : k]$ there are pr functions*

$$b^{(k)} : \mathbb{N}_0^k \rightarrow \mathbb{N}_0 \quad \text{bijective}$$

$$b_i^{(k)} : \mathbb{N}_0 \rightarrow \mathbb{N}_0$$

such that for all $x \in \mathbb{N}_0^k$ and all i

$$b_i^{(k)}(b^{(k)}(x_1, \dots, x_k)) = x_i$$

- $k > 2$: the fun part:

$$b^{(k)}(x_1, \dots, x_k) = b^{(2)}(x_1, b^{(k-1)}(x_2, \dots, x_k))$$

$$b_1^{(k)}(x) = b_1^{(2)}(x)$$

$$b_i^{(k)}(x) = b_{i-1}^{(k-1)}(b_2^{(2)}(x)) \quad \text{for } i > 1$$

9 μ -operator and μ -recursive functions

def: (unbounded) μ -operator For

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

we define

$$\mu f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

by

$$\mu f(n, x) = \begin{cases} \min\{m : f(m, x) = 0\} & \text{if it exists} \\ \Omega & \text{(undefined) otherwise} \end{cases}$$

$\mu f(n, x)$ returns smallest solution of equation $f(m, x) = 0$ if it exists. Could be implemented as unbounded while loop:

```
m=0;  
while f(m, x) != 0 {m = m+1}; return m
```

9 μ -operator and μ -recursive functions

def: (unbounded) μ -operator For

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

we define

$$\mu f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

by

$$\mu f(n, x) = \begin{cases} \min\{m : f(m, x) = 0\} & \text{if it exists} \\ \Omega & \text{(undefined) otherwise} \end{cases}$$

$\mu f(n, x)$ returns smallest solution of equation $f(m, x) = 0$ if it exists. Could be implemented as unbounded while loop:

```
m=0;
while f(m, x) != 0 {m = m+1}; return m
```

Inductive definition of a set R of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

1. constant functions $c_s^r \in R$

2. projections $p_i^r \in R$

3. successor function $S \in R$

4. substitution. If the following function are all in R

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in R$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in R

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in R$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$f(0, x) = g(x)$$

$$f(n+1, x) = h(n, f(n, x), x)$$

6. if $f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$ is in R , then μf is in R

7. these are all

9 μ -operator and μ -recursive functions

Inductive definition of a set R of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

1. constant functions $c_s^r \in R$

2. projections $p_i^r \in R$

3. successor function $S \in R$

4. substitution. If the following function are all in R

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in R$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in R

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in R$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$f(0, x) = g(x)$$

$$f(n+1, x) = h(n, f(n, x), x)$$

6. if $f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$ is in R , then μf is in R

7. these are all

what can be computed with functions in R ?

- Answer: everything that can be computed at all
- this is known as *Church's thesis*.
- We cannot prove it (based on what definition or axiom could we do that?)
- we *can* supply evidence, and we *will* do it. Of course here

9 μ -operator and μ -recursive functions

Inductive definition of a set R of computable functions:

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N}_0$$

1. constant functions $c_s^r \in R$

2. projections $p_i^r \in R$

3. successor function $S \in R$

4. substitution. If the following function are all in R

$$f : \mathbb{N}_0^r \rightarrow \mathbb{N} \text{ and } g_1, \dots, g_r : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

then also $h \in R$ where

$$h : \mathbb{N}_0^m \rightarrow \mathbb{N}_0$$

$$h(x) = f(g_1(x), \dots, g_r(x))$$

5. primitive recursion. If the following functions are in R

$$g : \mathbb{N}_0^r \rightarrow \mathbb{N}_0, h : \mathbb{N}_0^{r+2} \rightarrow \mathbb{N}_0$$

then also $f \in R$ where

$$f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$$

$$f(0, x) = g(x)$$

$$f(n+1, x) = h(n, f(n, x), x)$$

6. if $f : \mathbb{N}_0^{r+1} \rightarrow \mathbb{N}_0$ is in R , then μf is in R

7. these are all

what can be computed with functions in R ?

- Answer: everything that can be computed at all
- this is known as *Church's thesis*.
- We cannot prove it (based on what definition or axiom could we do that?)
- we *can* supply evidence, and we *will* do it. Of course here

hope in favor of Church's thesis proof that pr functions do not compute all computable functions relied on fact, that all pr functions are total. That proof collapses for μ -recursive functions. Exercise: in which line?