
ExPetDB (training + internal test) vs ArcPL opx (external held-out) data distribution
(a) ExPetDB opx_liq n=600
(b) ExPetDB opx_only n=1035
(c) ArcPL opx_liq held-out n=197
(d) ArcPL opx_only held-out n=197
count
P (kbar)
T (°C)
30 kbar
15 kbar
5 kbar
ExPetDB (train+test, gray underlay)
n total = 600
n train = 426
n test = 174
n citations = 93
T (1-99%) = 850-1660 °C
P (1-99%) = 0.0-40.0 kbar
regime counts:
shallow_crustal 137
deep_crustal_MASH 245
lithospheric_mantle 178
deeper_mantle 40
n total = 1035
n train = 845
n test = 190
n citations = 123
T (1-99%) = 850-1657 °C
P (1-99%) = 0.0-60.0 kbar
regime counts:
shallow_crustal 245
deep_crustal_MASH 383
lithospheric_mantle 259
deeper_mantle 148
n total = 197
partition = external holdout
(no train/test split)
n citations = 19
T (1-99%) = 825-1212 °C
P (1-99%) = 1.0-14.6 kbar
regime counts:
shallow_crustal 80
deep_crustal_MASH 116
lithospheric_mantle 1
deeper_mantle 0
Regime colors; train/test markers for ExPetDB panels (a, b); gray ExPetDB underlay under ArcPL points in panels (c, d)
shallow_crustal
deep_crustal_MASH
lithospheric_mantle
deeper_mantle
test (dark edge)
train (faded)
ExPetDB underlay (panels c, d)