

Reminder about homology: $\text{Top} \xrightarrow[\text{sing. complex}]{S} (\text{simplicial sets}) \xrightarrow[\text{linearization}]{C(-;A)} (\text{chain complex}) \xrightarrow[\text{n-th homology group}]{H_n} (\text{abelian groups})$

- For a space X , the singular complex $S(X)$ is the simplicial set with $S(X)_n = \text{maps}^{\text{dis}}(\Delta^n, X)$

$$\Delta^n = \text{topological } n\text{-simplex} = \{ (x_0, \dots, x_n) \in \mathbb{R}^{n+1} : x_i \geq 0, x_0 + \dots + x_n = 1 \}$$

- For a simplicial set Y and an abelian group, the linearization is the chain complex $C(Y; A)$ with

$$C_n(Y; A) = A[Y_n] \quad A\text{-linearization of } Y_n \quad (C_n(Y; A) = 0 \text{ for } n < 0)$$

$$d_n = \sum_{i=0, \dots, n} (-1)^i \cdot d_i^* : A[Y_n] \rightarrow A[Y_{n-1}]$$

- For a chain complex C and $n \in \mathbb{Z}$, the n -th homology group $H_n(C)$ is
$$\frac{\text{Ker}(d_n : C_n \rightarrow C_{n-1})}{\text{Im}(d_{n+1} : C_{n+1} \rightarrow C_n)}$$

Variation: Cohomology

Def: A cocchain complex C consists of abelian groups C^n for $n \in \mathbb{Z}$ and homomorphisms $d^n : C^n \rightarrow C^{n+1}$ such that $d^{n+1} \circ d^n = 0 : C^n \rightarrow C^{n+2}$. A morphism $f : C \rightarrow D$ of cocchain complexes

(cocchain map) consists of homomorphisms $f^n : C^n \rightarrow D^n$ such that $d_D^n \circ f^n = f^{n+1} \circ d_C^n$

The n -th cohomology group of a cocchain complex C is

$$H^n C = \frac{\text{Ker}(d^n : C^n \rightarrow C^{n+1})}{\text{Im}(d^{n-1} : C^{n-1} \rightarrow C^n)}$$

$$\begin{array}{ccc} C^n & \xrightarrow{f^n} & D^n \\ d^n \downarrow & & \downarrow d^{n+1} \\ C^{n+1} & \xrightarrow{f^{n+1}} & D^{n+1} \end{array}$$

A cocchain homotopy between two morphisms $f, g : C \rightarrow D$ of cocchain complexes consists of homomorphisms

$$s^n : C^n \rightarrow D^{n-1} \text{ for all } n \in \mathbb{Z} \text{ such that}$$

$$d^{n+1} \circ s^n + s^{n+1} \circ d^n = f^n - g^n \text{ for all } n \in \mathbb{Z}.$$

The main tools and properties carry over from chain complexes to cocchain complexes, with essentially the same proofs, such as:

- a morphism $f : C \rightarrow D$ of cocchain complexes induces a homomorphism $H^n f : H^n C \rightarrow H^n D$ of cohomology groups by $(H^n f)[x] = [f^n(x)]$ $x \in \text{Ker}(d^n : C^n \rightarrow C^{n+1})$
- cocchain homotopic morphisms $f, g : C \rightarrow D$ between cocchain complexes induce the same map in cohomology, i.e. $H^n f = H^n g$.
- every short exact sequence of cocchain complexes $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ given rise to a long exact sequence of cohomology groups:

$$\dots \rightarrow H^n A \xrightarrow{H^n f} H^n B \xrightarrow{H^n g} H^n C \xrightarrow{\partial} H^{n+1}(A) \rightarrow \dots$$

where the connecting homomorphism ∂ is defined as follows:

given $x \in C^n$ with $d^n(x) = 0$, choose $\tilde{x} \in B^n$ such that $g^n(\tilde{x}) = x$,

then $g^{n+1}(d_B^n(\tilde{x})) = d_C^n(g^n(\tilde{x})) = d_C^n(x) = 0$, so there is a

unique $y \in A^{n+1}$ such that $f^{n+1}(y) = d_B^{n+1}(\tilde{x})$. Set

$$\partial[x] = [y] \in H^{n+1}(A).$$

An isomorphism of categories:

We define a functor $D : (\text{chain complex}, \text{chain maps}) \xrightarrow{\cong} (\text{cocchain complex and cocchain maps})$

On objects: $(DC)^n = C_{-n}$, $d_{DC}^n = d_{C_{-n}} : C_{-n} \rightarrow C_{-n-1} = (DC)^{n+1}$

On morphisms: let $f : C \rightarrow C'$ be a chain map. Then

$$Df : DC \rightarrow DC' \text{ is the cocchain map with } (Df)^n = f_{-n} : C_{-n} \rightarrow C'_{-n} = (DC')^n$$

The process is reversible on the nose (and not just up to natural isomorphism), so

we have defined an isomorphism of categories.

The process is reversible on the nose (and not just up to natural isomorphism), so we have defined an isomorphism of categories.

$$(\mathcal{D}C)^n$$

Moreover:

$$H^n(\mathcal{D}C) = H_{-n}(C)$$

$f, g: C \rightarrow C'$ are chain homotopic $\Leftrightarrow Df, Dg: \mathcal{D}C \rightarrow \mathcal{D}C'$ are cochain homotopic

Construction: Let C be a chain complex and A an abelian group. We define a cochain complex $\text{Hom}(C, A)$ by

$$\text{Hom}(C, A)^n = \text{Hom}(C_n, A) \quad \text{with differential}$$

$$d^n: \text{Hom}(C, A)^n \rightarrow \text{Hom}(C, A)^{n+1}$$

$$\text{Then: } d^{n+1} \circ d^n = \text{Hom}(d_{n+2}, A) \circ \text{Hom}(d_{n+1}, A)$$

$$\text{Hom}(d_{n+2}, A): \text{Hom}(C_n, A) \rightarrow \text{Hom}(C_{n+2}, A)$$

$$= \text{Hom}(d_{n+2} \circ d_{n+1}, A) = \text{Hom}(0, A) = 0.$$

This construction becomes a contravariant functor from chain complexes to cochain complexes

$$\text{Hom}(-, A): (\text{chain complexes})^{\text{op}} \rightarrow (\text{cochain complexes})$$

On chain morphisms $f: C \rightarrow C'$, $\text{Hom}(f, A): \text{Hom}(C', A) \rightarrow \text{Hom}(C, A)$ is given by

$$\text{Hom}(f, A)^n = \text{Hom}(f_n, A): \text{Hom}(C'_n, A) \rightarrow \text{Hom}(C_n, A)$$

Lemma: Let $f, g: C \rightarrow C'$ be chain homotopic chain maps. Then the cochain maps $\text{Hom}(f, A), \text{Hom}(g, A): \text{Hom}(C', A) \rightarrow \text{Hom}(C, A)$ are cochain homotopic.

Proof: Suppose that $s = \{s_n: C_n \rightarrow C'_{n+2}\}_{n \in \mathbb{Z}}$ is chain homotopy, then

$$\{\text{Hom}(s_n, A): \text{Hom}(C'_{n+2}, A) \rightarrow \text{Hom}(C_n, A)\}_{n \in \mathbb{Z}} \text{ is a cochain homotopy between } \text{Hom}(f, A) \text{ and } \text{Hom}(g, A).$$

The singular cohomology of spaces and simplicial sets

Def: Let Y be a simplicial set and A an abelian group. The cohomology of Y with coefficients in A is

$$H^n(Y, A) = H^n(\text{Hom}(C(Y, \mathbb{Z}), A))$$

If $Y' \subseteq Y$ is a simplicial subset, the relative cohomology of the pair (Y, Y') is

$$H^n(Y, Y'; A) = H^n\left(\text{Hom}\left(\frac{C(Y, \mathbb{Z})}{C(Y', \mathbb{Z})}, A\right)\right).$$

If X is a space, the cohomology with coefficients in A is

$$H^n(X, A) = H^n(S(X), A) = H^n(\text{Hom}(C(S(X), \mathbb{Z}), A)).$$

If X' is a subspace of X , the relative cohomology $H^n(X, X'; A)$ is the relative cohomology of the pair $(S(X), S(X'))$.

The definition can be made more concrete / verifiable, as follows:

Construction: Let Y be a simplicial set, A an abelian group. We define a cochain complex $C^*(Y, A)$ by

$$C^n(Y, A) = \text{map}(Y_n, A) = \text{abelian group of maps } f: Y_n \rightarrow A \text{ under pointwise addition.}$$

The differential is defined by:

$$d^n(f)(y) = \sum_{i=0, \dots, n+1} (-1)^i \cdot f(d_i^*(y)) \quad \text{for } y \in Y_{n+1}$$

Omitted: verification that

$$d^{n+1}(d^n(f)) = 0.$$

If Y' is a simplicial subset of Y , we define $C^*(Y, Y'; A)$ as follows.

$$C^n(Y, Y'; A) = \{f: Y_n \rightarrow A : f(Y'_n) = 0\}$$

Omitted: this defines a sub-cochain complex of $C^*(Y, A)$.

Lemma: Let Y be a simplicial set and A an abelian group. Then there is an isomorphism of cochain complexes

$$H_0(C(Y, \mathbb{Z}), A) \cong C^*(Y, A), \text{ hence an isomorphism of cohomology groups}$$

$H^n(Y, A) \cong H^n(C^*(Y, A))$. These isomorphisms are natural for morphisms of simplicial sets. For a simplicial subset Y' of Y there is an isomorphism of cochain complexes

$$\text{Hom} \left(\frac{C(Y; \mathbb{Z})}{C(Y'; \mathbb{Z})}, A \right) \cong C^*(Y, Y'; A), \text{ natural for morphisms of pairs of simplicial sets.}$$


Proof: In the absolute case: define a specific isomorphism $\varphi: \text{Hom}(C(Y; \mathbb{Z}), A) \rightarrow C^*(Y, A)$ in dimension $n \geq 0$

$$\varphi^n: \text{Hom}(C_n(Y; \mathbb{Z}), A) = \text{Hom}(\mathbb{Z}[Y_n], A) \xrightarrow{\cong} \text{map}(Y_n, A)$$

evaluation at the generators of the free abelian group $\mathbb{Z}[Y_n]$, is an isomorphism of n -cops.

Omitted: the φ^n 's form a morphism of cochain complexes. So $\varphi = \{\varphi^n\}$ is an isomorphism of cochain complexes. \square

Schematically:



the key properties of singular homology all have analogs for cohomology:

Homotopy invariance: Let $f, g: X \rightarrow Y$ be homotopic continuous maps. Then for all $n \geq 0$ and all abelian groups A ,

$$H^n(f; A) = H^n(g; A): H^n(Y, A) \rightarrow H^n(X, A).$$

Proof: Since f and g are homotopic, $C(f, \mathbb{Z}), C(g, \mathbb{Z}): C(X, \mathbb{Z}) \rightarrow C(Y, \mathbb{Z})$ are chain homotopic.

By an earlier lemma, $\text{Hom}(C(f, \mathbb{Z}), A)$ and $\text{Hom}(C(g, \mathbb{Z}), A)$ are cochain homotopic, so they induce the same map on cohomology groups. \square

Long exact sequence: Let Y' be a simplicial subset of a simplicial set Y . Then we have a short exact sequence of cochain complexes

$$0 \rightarrow C^*(Y, Y'; A) \rightarrow C^*(Y, A) \xrightarrow{\text{inclusion}} C^*(Y', A) \rightarrow 0$$

\Rightarrow long exact sequence of cohomology groups

$$\dots \rightarrow H^n(Y, Y'; A) \rightarrow H^n(Y, A) \rightarrow H^n(Y', A) \xrightarrow{\partial} H^{n+1}(Y, Y'; A) \rightarrow \dots$$

For a subspace X' of a space X , we can apply this to the pair $(S(X), S(X'))$ to get a long exact sequence of singular cohomology groups.

Excision: Let (X, Y, U) be an excisive triple of spaces, i.e. $U \subseteq Y \subseteq X$ and $\bar{U} \subseteq \bar{Y}$.

In the proof of excision for homology we showed that the inclusions induce a quasi-isomorphism of chain complexes

$$i: \frac{C(S(X \setminus U); \mathbb{Z})}{C(S(Y \setminus U); \mathbb{Z})} \longrightarrow \frac{C(S(X), \mathbb{Z})}{C(S(Y), \mathbb{Z})}, \text{ i.e. it induces an}$$

isomorphism of all homology groups.

Prop: Let $f: C \rightarrow D$ be a quasi-isomorphism of chain complexes of free abelian groups. Then f is a chain homotopy equivalence.

Proof: Deferred to a separate video.

Since i is a chain homotopy equivalence, $\text{Hom}(i, A): \text{Hom}\left(\frac{C(S(X), \mathbb{Z})}{C(S(Y), \mathbb{Z})}, A\right) \rightarrow \text{Hom}\left(\frac{C(S(X \setminus U), \mathbb{Z})}{C(S(Y \setminus U), \mathbb{Z})}, A\right)$ is a cochain homotopy equivalence.

So $\text{Hom}(i, A)$ induces isomorphisms of cohomology groups

$$H^n(X, Y; A) = H^n\left(\text{Hom}\left(\frac{C(S(X), \mathbb{Z})}{C(S(Y), \mathbb{Z})}, A\right)\right) \xrightarrow{\cong} H^n\left(\text{Hom}\left(\frac{C(S(X \setminus U), \mathbb{Z})}{C(S(Y \setminus U), \mathbb{Z})}, A\right)\right) \xrightarrow{\vee} H^n(X \setminus U, Y \setminus U; A)$$

$$\begin{array}{ccc} C(S^1, \mathbb{Z}) & \xrightarrow{\cong} & C(S(X \cup U), \mathbb{Z}) \\ & \searrow \text{inclusion} & \downarrow \\ & & H^n(X \cup U, \mathbb{Z}) \end{array}$$

Given the same fundamental formal properties, the basic calculations for singular homology can be repeated in much the same way for singular cohomology:

$$H^n(S^n; A) \cong \begin{cases} A & \text{for } n=0, n \\ 0 & \text{otherwise} \end{cases}$$

$$H^n(D^n, S^{n-1}; A) \cong \begin{cases} A & \text{for } n=n \\ 0 & \text{otherwise} \end{cases}$$

Reminder: Let X be an absolute CW-complex with skeleton $\{X_n\}_{n \geq 0}$. The cellular chain complex is given by

$$C_n^{\text{cell}}(X; \mathbb{Z}) = H_n(X_n, X_{n-1}; \mathbb{Z}) \quad , \text{ with cellular differential defined as the composite}$$

$$C_n^{\text{cell}}(X; \mathbb{Z}) = H_n(X_n, X_{n-1}; \mathbb{Z}) \xrightarrow{\partial} H_{n-1}(X_{n-1}; \mathbb{Z}) \longrightarrow H_{n-1}(X_{n-1}, X_{n-2}; \mathbb{Z}) = C_{n-1}^{\text{cell}}(X; \mathbb{Z})$$

We define the cellular cochain complex of X with coefficients in A as

$$C_{\text{cell}}^*(X; A) = \text{Hom}(C_*^{\text{cell}}(X; \mathbb{Z}), A)$$

Thm: There is an isomorphism $H^n(C_{\text{cell}}^*(X; A)) \cong H^n(X; A)$ that is moreover natural for cellular maps in X .

Proof: Copy for the proof for homology. \square

Example: Let X be a CW-complex with no cells in any odd dimension.

Then $C_*^{\text{cell}}(X; \mathbb{Z})$ has trivial differentials. So $\text{Hom}(C_*^{\text{cell}}(X; \mathbb{Z}), A)$ has trivial differentials and so

$$\begin{aligned} H^n(X; A) &\cong H^n(C_{\text{cell}}^*(X; A)) = \text{Hom}(C_n^{\text{cell}}(X; \mathbb{Z}), A) \stackrel{\parallel}{=} C_{\text{cell}}^{*n}(X; A) \\ &\cong \text{Hom}(\mathbb{Z}[J_n], A) \cong \text{map}(J_n, A) \\ &\quad \uparrow \\ &\quad \text{set of } n\text{-cells} \end{aligned}$$

Example \mathbb{CP}^∞ has a CW-structure with exactly 1 cell in every even dimension and no cells in odd dimensions.

$$H^n(\mathbb{CP}^\infty; A) \cong \begin{cases} A & \text{for } n \geq 0 \text{ even} \\ 0 & \text{for } n \text{ odd} \end{cases}$$