

IT575 Computational Shape Modeling

Module - 3: Discrete Geometry

Aditya Tatu

Sampling a Surface

- Let M be a surface. A finite subset $P \in M$ is called a *sample* set of M , and the set P is often referred to as a *point cloud*.




Figure : (left) Original Surface, (right) Point cloud representation.

Quality of Sampling

- Measure of uniformity: A sample-set P is said to be an *r-covering* of the surface M if

$$\bigcup_{p \in P} B_r(p) = M \Leftrightarrow \forall x \in M, d_M(x, P) \leq r.$$

- ▶ Issue: If P is an *r-covering*, is it a minimal set?
- A sample set P is said to be *r-separated* if
 $d_M(p, q) \geq r, \forall p, q \in P, \text{ with } p \neq q.$

Quality of Sampling

- Measure of uniformity: A sample-set P is said to be an *r-covering* of the surface M if

$$\cup_{p \in P} B_r(p) = M \Leftrightarrow \forall x \in M, d_M(x, P) \leq r.$$

- ▶ Issue: If P is an *r-covering*, is it a minimal set?
- A sample set P is said to be *r-separated* if
 $d_M(p, q) \geq r, \forall p, q \in P, \text{ with } p \neq q.$

Quality of Sampling

- Measure of uniformity: A sample-set P is said to be an *r-covering* of the surface M if

$$\cup_{p \in P} B_r(p) = M \Leftrightarrow \forall x \in M, d_M(x, P) \leq r.$$

- ▶ Issue: If P is an *r-covering*, is it a minimal set?
- A sample set P is said to be *r-separated* if $d_M(p, q) \geq r$, $\forall p, q \in P$, with $p \neq q$.

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____-covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____ -covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____-covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a _____-separated and _____-covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____ -covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____ -covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____ -covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Sampling strategy: Farthest Point Sampling

- ▶ Let $p_1 = x_1 \in M$, and $P = \{p_1\}$.
- ▶ How should the second point be selected?
- ▶ Let $p_2 = \arg \max_{x \in M} d_M(x, P)$, and $P = \{p_1, p_2\}$.
- ▶ P is an _____-separated set and an _____ -covering set.
- ▶ Similarly, $p_3 = \arg \max_{x \in M} d_M(x, P)$, and $P = P \cup \{p_3\}$.
- ▶ Let $P = \{p_1, \dots, p_n\}$ be the sample set. It is a
_____ -separated and _____ -covering sample set.
- Issues with Farthest point sampling?

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

Voronoi Tessellation

- Let P be a sample-set of the surface M . Every sample point represents a small nbhd of M .
- For $p_i \in P$, let $V_i(P) = \{x \in M \mid d_M(x, p_i) < d_M(x, p_j), j \neq i, p_j \in P\}$.
- $V_i(P)$ is called the *Voronoi region* of p_i with respect to the sample-set P , and the collection of Voronoi regions of M is called the *Voronoi decomposition* of M generated by P .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j)$, x is said to be on the *Voronoi edge* V_{ij} of the Voronoi regions V_i, V_j .
- If for $x \in M$, $d_M(x, p_i) = d_M(x, p_j) = d_M(x, p_k)$, x is said to be a *Voronoi vertex* V_{ijk} of the Voronoi regions V_i, V_j, V_k .
- Demo: <http://alexbeutel.com/webgl/voronoi.html>.

- Let \bar{V}_i denote the closure of V_i . Note $M = \cup_i \bar{V}_i$, and $V_i \cap V_j = \emptyset$.
- The decomposition of M into $\{V_i \mid p_i \in P\}$ such that $M = \cup_i V_i$, and $V_i \cap V_j = \emptyset$ is called a *tessellation or cell complex* of M .
- If the tiles V_i are Voronoi regions, we call it a *Voronoi tessellation*.




Figure : Voronoi cells. Image from: *Bronstein, Bronstein, Kimmel, Numerical Geometry of Nonrigid shapes, Springer 2008*, henceforth - BBK08

- Let \bar{V}_i denote the closure of V_i . Note $M = \cup_i \bar{V}_i$, and $V_i \cap V_j = \emptyset$.
 - The decomposition of M into $\{V_i \mid p_i \in P\}$ such that $M = \cup_i V_i$, and $V_i \cap V_j = \emptyset$ is called a *tessellation or cell complex* of M .
- If the tiles V_i are Voronoi regions, we call it a *Voronoi tessellation*.




Figure : Voronoi cells. Image from: *Bronstein, Bronstein, Kimmel, Numerical Geometry of Nonrigid shapes, Springer 2008*, henceforth - BBK08

- Let \bar{V}_i denote the closure of V_i . Note $M = \cup_i \bar{V}_i$, and $V_i \cap V_j = \emptyset$.
- The decomposition of M into $\{V_i \mid p_i \in P\}$ such that $M = \cup_i V_i$, and $V_i \cap V_j = \emptyset$ is called a *tessellation or cell complex* of M .
- If the tiles V_i are Voronoi regions, we call it a *Voronoi tessellation*.




Figure : Voronoi cells. Image from: *Bronstein, Bronstein, Kimmel, Numerical Geometry of Nonrigid shapes, Springer 2008*, henceforth - BBK08

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.
- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

- Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.
- If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.
- Thus, $p_i = \frac{\int_{V_i} x dx}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.

- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

► Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.

► If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.

► Thus, $p_i = \frac{\int_{V_i} x da}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.
- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

- Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.
- If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.
- Thus, $p_i = \frac{\int_{V_i} x dx}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.
- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

- Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.
- If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.
- Thus, $p_i = \frac{\int_{V_i} x dx}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.

- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

► Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.

► If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.

► Thus, $p_i = \frac{\int_{V_i} x da}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.

- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

► Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.

► If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.

► Thus, $p_i = \frac{\int_{V_i} x da}{\int_{V_i} da}$ is the *centroid* of the Voronoi cell V_i .

Sampling and Voronoi tessellation

- Let P be a sample set of a surface M .
- Let $s : M \rightarrow P$ be defined by

$$s(x) = p_i \text{ where } p_i \text{ is such that } x \in V_i(M).$$

- By representing M by P , $\forall x \in M$, error incorporated is $e(x) = d_M(x, s(x) = p_i)$.
- Total error:

$$e(P) = \int_M d_M^2(x, s(x)) = \sum_{i=1}^N \int_{V_i(P)} d_M^2(x, p_i) da$$

- Thus, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} d_M^2(x, x^*) da$.
- If $M \subset \mathbb{R}^2$, $p_i = \arg \min_{x^* \in V_i} \int_{V_i} \|x - x^*\|^2 dx$.
- Thus, $p_i = \frac{\int_{V_i} x dx}{\int_{V_i} dx}$ is the *centroid* of the Voronoi cell V_i .

Lloyd-Max algorithm

- Given a surface M , and an initial sample-set P .
- Construct voronoi tessellation of M generated by P .
- Update: $P = \text{centroids of voronoi cells}$. Repeat till convergence.

Lloyd-Max algorithm

- Given a surface M , and an initial sample-set P .
- Construct voronoi tessellation of M generated by P .
- Update: $P = \text{centroids of voronoi cells}$. Repeat till convergence.

Lloyd-Max algorithm

- Given a surface M , and an initial sample-set P .
- Construct voronoi tessellation of M generated by P .
- Update: $P = \text{centroids of voronoi cells}$. Repeat till convergence.




Figure : (left) Inadequate (right) Adequate sampling density. The voronoi regions can be deformed to unit disks in the example on the right. Image from BBK08.

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
- ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood* of p_k .
- ▶ Thus P with the above edges gives an *undirected binary graph*.
- ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
- ▶ The edges give a one-dimensional approximation of the surface M .

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
 - ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood* of p_k .
 - ▶ Thus P with the above edges gives an *undirected binary graph*.
 - ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
 - ▶ The edges give a one-dimensional approximation of the surface M .

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
- ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood of p_k* .
- ▶ Thus P with the above edges gives an *undirected binary graph*.
- ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
- ▶ The edges give a one-dimensional approximation of the surface M .

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
- ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood* of p_k .
- ▶ Thus P with the above edges gives an *undirected binary graph*.
- ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
- ▶ The edges give a one-dimensional approximation of the surface M .

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
- ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood* of p_k .
- ▶ Thus P with the above edges gives an *undirected binary graph*.
- ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
- ▶ The edges give a one-dimensional approximation of the surface M .

Edge structure

- The sample-set P is a zero-dimensional approximation of the surface M .
- Connect two distinct sample points p_i and p_j if $d_M(p_i, p_j) < r$.
- ▶ The set $N(p_k) = \{p_q \in P \mid d_M(p_k, p_q) < r\}$ is the *Neighborhood* of p_k .
- ▶ Thus P with the above edges gives an *undirected binary graph*.
- ▶ Approximate Edge lengths $l(p_i, p_j) \simeq \|p_i - p_j\|_2$.
- ▶ The edges give a one-dimensional approximation of the surface M .

Two dimensional approximation

- Let p_i, p_j and p_k be three sample-sets whose Voronoi cells are adjacent to each other and thus yield a vertex V_{ijk} .
- Connect p_i with p_j by a curve of shortest length: Γ_{ij} . Similarly curves Γ_{jk} and Γ_{ki} connect points p_j with p_k and p_k with p_i , resp., forming a *geodesic triangle*.
- Degenerate cases: (a) Three *collinear* points, (b) Four *cocircular* points.
- Decomposition of M into these geodesic triangles is called *Delaunay tessellation*.

Two dimensional approximation

- Let p_i, p_j and p_k be three sample-sets whose Voronoi cells are adjacent to each other and thus yield a vertex V_{ijk} .
- Connect p_i with p_j by a curve of shortest length: Γ_{ij} . Similarly curves Γ_{jk} and Γ_{ki} connect points p_j with p_k and p_k with p_i , resp., forming a *geodesic triangle*.
- Degenerate cases: (a) Three *collinear* points, (b) Four *cocircular* points.
- Decomposition of M into these geodesic triangles is called *Delaunay tessellation*.

Two dimensional approximation

- Let p_i, p_j and p_k be three sample-sets whose Voronoi cells are adjacent to each other and thus yield a vertex V_{ijk} .
- Connect p_i with p_j by a curve of shortest length: Γ_{ij} . Similarly curves Γ_{jk} and Γ_{ki} connect points p_j with p_k and p_k with p_i , resp., forming a *geodesic triangle*.
- Degenerate cases: (a) Three *collinear* points, (b) Four *cocircular* points.
- Decomposition of M into these geodesic triangles is called *Delaunay tessellation*.

Two dimensional approximation

- Let p_i, p_j and p_k be three sample-sets whose Voronoi cells are adjacent to each other and thus yield a vertex V_{ijk} .
- Connect p_i with p_j by a curve of shortest length: Γ_{ij} . Similarly curves Γ_{jk} and Γ_{ki} connect points p_j with p_k and p_k with p_i , resp., forming a *geodesic triangle*.
- Degenerate cases: (a) Three *collinear* points, (b) Four *cocircular* points.
- Decomposition of M into these geodesic triangles is called *Delaunay tessellation*.

Triangular meshes

- We can approximate the geodesic triangles by Euclidean/planar triangles, giving us a triangular mesh:




Figure : Triangular mesh surface approximation.

- Triangular meshes are by far the most preferred discretization.
 - ▶ Approximate surface locally via planar polygons - we prefer triangles.
 - ▶ Optimized software and hardware available for processing triangles.

- Triangular meshes are by far the most preferred discretization.
- ▶ Approximate surface locally via planar polygons - we prefer triangles.
- ▶ Optimized software and hardware available for processing triangles.

- Triangular meshes are by far the most preferred discretization.
- ▶ Approximate surface locally via planar polygons - we prefer triangles.
- ▶ Optimized software and hardware available for processing triangles.

Datastructure for Mesh

- Data: Geometry & Topology.

- ▶ Geometry: For a mesh with n vertices, the geometry consists of an *ordered* $n \times 3$ array of coordinates of the n vertices.
- ▶ Topology: Connectivity information stored as an $m \times 3$ integer array, where m is the number of triangles in the mesh, and each row consisting of indices i, j, k such that the vertices v_i, v_j, v_k form a triangle.
- ▶ A triangle is often referred to as a *face*. The surface represented by the triangular mesh is:

$$M' = \bigcup_{i=1}^m \text{conv}(x_{t_i}, y_{t_i}, z_{t_i})$$

- ▶ Additional information may be stored, for example list of edges as a $k \times 2$ array consisting of the index pairs of vertices forming an edge.

Datastructure for Mesh

- Data: Geometry & Topology.
- ▶ Geometry: For a mesh with n vertices, the geometry consists of an *ordered* $n \times 3$ array of coordinates of the n vertices.
- ▶ Topology: Connectivity information stored as an $m \times 3$ integer array, where m is the number of triangles in the mesh, and each row consisting of indices i, j, k such that the vertices v_i, v_j, v_k form a triangle.
- ▶ A triangle is often referred to as a *face*. The surface represented by the triangular mesh is:

$$M' = \bigcup_{i=1}^m \text{conv}(x_{t_i}, y_{t_i}, z_{t_i})$$

- ▶ Additional information may be stored, for example list of edges as a $k \times 2$ array consisting of the index pairs of vertices forming an edge.

Datastructure for Mesh

- Data: Geometry & Topology.
- ▶ Geometry: For a mesh with n vertices, the geometry consists of an *ordered* $n \times 3$ array of coordinates of the n vertices.
- ▶ Topology: Connectivity information stored as an $m \times 3$ integer array, where m is the number of triangles in the mesh, and each row consisting of indices i, j, k such that the vertices v_i, v_j, v_k form a triangle.
- ▶ A triangle is often referred to as a *face*. The surface represented by the triangular mesh is:

$$M' = \bigcup_{i=1}^m \text{conv}(x_{t_i}, y_{t_i}, z_{t_i})$$

- ▶ Additional information may be stored, for example list of edges as a $k \times 2$ array consisting of the index pairs of vertices forming an edge.

Datastructure for Mesh

- Data: Geometry & Topology.
- ▶ Geometry: For a mesh with n vertices, the geometry consists of an *ordered* $n \times 3$ array of coordinates of the n vertices.
- ▶ Topology: Connectivity information stored as an $m \times 3$ integer array, where m is the number of triangles in the mesh, and each row consisting of indices i, j, k such that the vertices v_i, v_j, v_k form a triangle.
- ▶ A triangle is often referred to as a *face*. The surface represented by the triangular mesh is:

$$M' = \cup_{i=1}^m \text{conv}(x_{t_i}, y_{t_i}, z_{t_i})$$

- ▶ Additional information may be stored, for example list of edges as a $k \times 2$ array consisting of the index pairs of vertices forming an edge.

Datastructure for Mesh

- Data: Geometry & Topology.
- ▶ Geometry: For a mesh with n vertices, the geometry consists of an *ordered* $n \times 3$ array of coordinates of the n vertices.
- ▶ Topology: Connectivity information stored as an $m \times 3$ integer array, where m is the number of triangles in the mesh, and each row consisting of indices i, j, k such that the vertices v_i, v_j, v_k form a triangle.
- ▶ A triangle is often referred to as a *face*. The surface represented by the triangular mesh is:

$$M' = \bigcup_{i=1}^m \text{conv}(x_{t_i}, y_{t_i}, z_{t_i})$$

- ▶ Additional information may be stored, for example list of edges as a $k \times 2$ array consisting of the index pairs of vertices forming an edge.

Degenerate cases

- Non-manifold meshes:




Figure : Examples of non-manifold meshes. Image from *Polygonal Mesh Processing*, Botsch et al., A.K.Peters, 2010, henceforth BKPAL10

- ▶ Assumptions: No non-manifold vertices or edges.

Degenerate cases

- Non-manifold meshes:




Figure : Examples of non-manifold meshes. Image from *Polygonal Mesh Processing*, Botsch et al., A.K.Peters, 2010, henceforth BKPAL10

- ▶ Assumptions: No non-manifold vertices or edges.

Discrete Differential Geometry

Differential geometric properties

- A triangular mesh is a linear approximation of the original surface.
- Most of differential geometry depends on derivatives of different quantities.
- We would want to compute the values of the derivatives of given quantities/signal/function at the vertices.
- Given a manifold M , and a function $f : M \rightarrow \mathbb{R}$ on it, the directional derivative of f along $v \in T_p M$ at p is defined as $(D_v f)(p) = \lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(p)}{t}$, where $\gamma : (-\epsilon, \epsilon) \rightarrow M$ is a smooth curve on M such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v \in T_p M$.

Differential geometric properties

- A triangular mesh is a linear approximation of the original surface.
- Most of differential geometry depends on derivatives of different quantities.
- We would want to compute the values of the derivatives of given quantities/signal/function at the vertices.
- Given a manifold M , and a function $f : M \rightarrow \mathbb{R}$ on it, the directional derivative of f along $v \in T_p M$ at p is defined as $(D_v f)(p) = \lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(p)}{t}$, where $\gamma : (-\epsilon, \epsilon) \rightarrow M$ is a smooth curve on M such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v \in T_p M$.

Differential geometric properties

- A triangular mesh is a linear approximation of the original surface.
- Most of differential geometry depends on derivatives of different quantities.
- We would want to compute the values of the derivatives of given quantities/signal/function at the vertices.
- Given a manifold M , and a function $f : M \rightarrow \mathbb{R}$ on it, the directional derivative of f along $v \in T_p M$ at p is defined as $(D_v f)(p) = \lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(p)}{t}$, where $\gamma : (-\epsilon, \epsilon) \rightarrow M$ is a smooth curve on M such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v \in T_p M$.

Differential geometric properties

- A triangular mesh is a linear approximation of the original surface.
- Most of differential geometry depends on derivatives of different quantities.
- We would want to compute the values of the derivatives of given quantities/signal/function at the vertices.
- Given a manifold M , and a function $f : M \rightarrow \mathbb{R}$ on it, the directional derivative of f along $v \in T_p M$ at p is defined as $(D_v f)(p) = \lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(p)}{t}$, where $\gamma : (-\epsilon, \epsilon) \rightarrow M$ is a smooth curve on M such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v \in T_p M$.

Approximating derivatives

- Let us first consider sampling of a 1 D signal as shown in the figure below.




Figure : Sampling of a 1D signal.

- In the absence of the original signal f , we assume that the original signal lies in the span of the following *piece-wise linear* basis:




Figure : Piece-wise linear basis, note that $b_n(x) = 0, x \notin [x_{n-1}, x_{n+1}]$.

- Thus

$$\tilde{f}(x) = \dots + a_{n-1}b_{n-1}(x) + a_n b_n(x) + a_{n+1}b_{n+1}(x) + \dots$$

- Coefficients a_i 's can be found by requiring that $\tilde{f}(x = x_n) = f_n$, the given sample values.
- For this particular choice of basis it is clear that $a_n = f_n, \forall n$, which gives for any $x \in [x_n, x_{n+1}]$,

$$\tilde{f}(x) = \frac{x_{n+1} - x}{x_{n+1} - x_n} f_n + \frac{x - x_n}{x_{n+1} - x_n} f_{n+1}.$$

- In the absence of the original signal f , we assume that the original signal lies in the span of the following *piece-wise linear* basis:




Figure : Piece-wise linear basis, note that $b_n(x) = 0, x \notin [x_{n-1}, x_{n+1}]$.

- Thus
 $\tilde{f}(x) = \dots + a_{n-1}b_{n-1}(x) + a_n b_n(x) + a_{n+1}b_{n+1}(x) + \dots$
- Coefficients a_i 's can be found by requiring that
 $\tilde{f}(x = x_n) = f_n$, the given sample values.
- For this particular choice of basis it is clear that $a_n = f_n, \forall n$, which gives for any $x \in [x_n, x_{n+1}]$,

$$\tilde{f}(x) = \frac{x_{n+1} - x}{x_{n+1} - x_n} f_n + \frac{x - x_n}{x_{n+1} - x_n} f_{n+1}.$$

- In the absence of the original signal f , we assume that the original signal lies in the span of the following *piece-wise linear* basis:




Figure : Piece-wise linear basis, note that $b_n(x) = 0, x \notin [x_{n-1}, x_{n+1}]$.

- Thus
 $\tilde{f}(x) = \dots + a_{n-1}b_{n-1}(x) + a_n b_n(x) + a_{n+1}b_{n+1}(x) + \dots$
- Coefficients a_i 's can be found by requiring that
 $\tilde{f}(x = x_n) = f_n$, the given sample values.
- For this particular choice of basis it is clear that $a_n = f_n, \forall n$, which gives for any $x \in [x_n, x_{n+1}]$,

$$\tilde{f}(x) = \frac{x_{n+1} - x}{x_{n+1} - x_n} f_n + \frac{x - x_n}{x_{n+1} - x_n} f_{n+1}.$$

- In the absence of the original signal f , we assume that the original signal lies in the span of the following *piece-wise linear* basis:




Figure : Piece-wise linear basis, note that $b_n(x) = 0, x \notin [x_{n-1}, x_{n+1}]$.

- Thus
 $\tilde{f}(x) = \dots + a_{n-1}b_{n-1}(x) + a_n b_n(x) + a_{n+1}b_{n+1}(x) + \dots$
- Coefficients a_i 's can be found by requiring that
 $\tilde{f}(x = x_n) = f_n$, the given sample values.
- For this particular choice of basis it is clear that $a_n = f_n, \forall n$, which gives for any $x \in [x_n, x_{n+1}]$,

$$\tilde{f}(x) = \frac{x_{n+1} - x}{x_{n+1} - x_n} f_n + \frac{x - x_n}{x_{n+1} - x_n} f_{n+1}.$$

- As should be clear from the equation, \tilde{f} is simply a linear interpolation of the sample values.




Figure : Original signal f and interpolated signal \tilde{f} .

- This allows us to compute derivatives of \tilde{f} at any interior point $x \in (x_n, x_{n+1})$ as

$$\frac{d}{dx} \tilde{f}(x) = f_n \frac{d}{dx} b_n(x) + f_{n+1} \frac{d}{dx} b_{n+1}(x)$$

- As should be clear from the equation, \tilde{f} is simply a linear interpolation of the sample values.




Figure : Original signal f and interpolated signal \tilde{f} .

- This allows us to compute derivatives of \tilde{f} at any interior point $x \in (x_n, x_{n+1})$ as

$$\frac{d}{dx} \tilde{f}(x) = f_n \frac{d}{dx} b_n(x) + f_{n+1} \frac{d}{dx} b_{n+1}(x)$$




Figure : Piece-wise linear basis

- Obviously, for $x \in (x_n, x_{n+1})$,

$$\frac{d}{dx} b_n(x) = \frac{-1}{x_{n+1} - x_n} \text{ and } \frac{d}{dx} b_{n+1}(x) = \frac{1}{x_{n+1} - x_n}$$

- Thus for $x \in (x_n, x_{n+1})$,

$$\frac{d}{dx} \tilde{f}(x) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$$




Figure : Piece-wise linear basis

- Obviously, for $x \in (x_n, x_{n+1})$,

$$\frac{d}{dx} b_n(x) = \frac{-1}{x_{n+1} - x_n} \text{ and } \frac{d}{dx} b_{n+1}(x) = \frac{1}{x_{n+1} - x_n}$$

- Thus for $x \in (x_n, x_{n+1})$,

$$\frac{d}{dx} \tilde{f}(x) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$$

- At $x = x_n$?
 - Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
$$D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$$
, where $\omega \in [0, 1]$.
 - ▶ In other words, compute an *average* of the two.

- At $x = x_n$?
- Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
$$D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$$
, where $\omega \in [0, 1]$.
 - ▶ In other words, compute an *average* of the two.

- At $x = x_n$?
- Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
$$D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$$
, where $\omega \in [0, 1]$.
- ▶ In other words, compute an *average* of the two.

- At $x = x_n$?
- Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
 $D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$, where $\omega \in [0, 1]$.
 - ▶ In other words, compute an average of the two.

- At $x = x_n$?
- Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
$$D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$$
, where $\omega \in [0, 1]$.
- ▶ In other words, compute an average of the two.

- At $x = x_n$?
- Several choices:
 - ▶ Forward Difference: $D_+ \tilde{f}(x_n) = \frac{f_{n+1} - f_n}{x_{n+1} - x_n}$.
 - ▶ Backward Difference: $D_- \tilde{f}(x_n) = \frac{f_n - f_{n-1}}{x_n - x_{n-1}}$.
 - ▶ Any convex combination:
$$D_\omega \tilde{f}(x_n) = \omega D_+ \tilde{f}(x_n) + (1 - \omega) D_- \tilde{f}(x_n)$$
, where $\omega \in [0, 1]$.
 - ▶ In other words, compute an *average* of the two.

Gradients of functions on a mesh

- Linear basis on a triangle of the mesh:




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- Any piece-wise linear function on the mesh with values at the vertices f_i can be written as

$$f(x) = f_i B_i(x) + f_j B_j(x) + f_k B_k(x)$$

- The gradient of f at any $x \in \text{Int}(\text{tri}(x_i, x_j, x_k))$ is then

$$\nabla f(x) = f_i \nabla B_i(x) + f_j \nabla B_j(x) + f_k \nabla B_k(x)$$

Gradients of functions on a mesh

- Linear basis on a triangle of the mesh:




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- Any piece-wise linear function on the mesh with values at the vertices f_i can be written as

$$f(x) = f_i B_i(x) + f_j B_j(x) + f_k B_k(x)$$

- The gradient of f at any $x \in \text{Int}(\text{tri}(x_i, x_j, x_k))$ is then

$$\nabla f(x) = f_i \nabla B_i(x) + f_j \nabla B_j(x) + f_k \nabla B_k(x)$$

Gradients of functions on a mesh

- Linear basis on a triangle of the mesh:




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- Any piece-wise linear function on the mesh with values at the vertices f_i can be written as

$$f(x) = f_i B_i(x) + f_j B_j(x) + f_k B_k(x)$$

- The gradient of f at any $x \in \text{Int}(\text{tri}(x_i, x_j, x_k))$ is then

$$\nabla f(x) = f_i \nabla B_i(x) + f_j \nabla B_j(x) + f_k \nabla B_k(x)$$




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- What is $\nabla B_i(x)$?

► $\nabla B_i(x) = \frac{(x_k - x_j)^\perp}{2A_T}$.

- Note that $B_i(x) + B_j(x) + B_k(x) = 1$ gives
 $\nabla B_i(x) = -(\nabla B_j(x) + \nabla B_k(x))$.

► Putting it together, we get

$$\nabla f(x) = (f_j - f_i)\nabla B_j(x) + (f_k - f_i)\nabla B_k(x)$$

- Note that $\nabla B_i(x) = \frac{N \times (x_k - x_j)}{2A_T}$, which gives

$$\nabla f(x) = f_i \frac{N \times (x_k - x_j)}{2A_T} + f_j \frac{N \times (x_i - x_k)}{2A_T} + f_k \frac{N \times (x_j - x_i)}{2A_T}$$




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- What is $\nabla B_i(x)$?
- ▶ $\nabla B_i(x) = \frac{(x_k - x_j)^\perp}{2A_T}$.
- Note that $B_i(x) + B_j(x) + B_k(x) = 1$ gives
 $\nabla B_i(x) = -(\nabla B_j(x) + \nabla B_k(x))$.
- ▶ Putting it together, we get

$$\nabla f(x) = (f_j - f_i)\nabla B_j(x) + (f_k - f_i)\nabla B_k(x)$$

- Note that $\nabla B_i(x) = \frac{N \times (x_k - x_j)}{2A_T}$, which gives

$$\nabla f(x) = f_i \frac{N \times (x_k - x_j)}{2A_T} + f_j \frac{N \times (x_i - x_k)}{2A_T} + f_k \frac{N \times (x_j - x_i)}{2A_T}$$




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- What is $\nabla B_i(x)$?
- ▶ $\nabla B_i(x) = \frac{(x_k - x_j)^\perp}{2A_T}$.
- Note that $B_i(x) + B_j(x) + B_k(x) = 1$ gives
 $\nabla B_i(x) = -(\nabla B_j(x) + \nabla B_k(x))$.
- ▶ Putting it together, we get

$$\nabla f(x) = (f_j - f_i)\nabla B_j(x) + (f_k - f_i)\nabla B_k(x)$$

- Note that $\nabla B_i(x) = \frac{N \times (x_k - x_j)}{2A_T}$, which gives

$$\nabla f(x) = f_i \frac{N \times (x_k - x_j)}{2A_T} + f_j \frac{N \times (x_i - x_k)}{2A_T} + f_k \frac{N \times (x_j - x_i)}{2A_T}$$




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- What is $\nabla B_i(x)$?
- ▶ $\nabla B_i(x) = \frac{(x_k - x_j)^\perp}{2A_T}$.
- Note that $B_i(x) + B_j(x) + B_k(x) = 1$ gives $\nabla B_i(x) = -(\nabla B_j(x) + \nabla B_k(x))$.
- ▶ Putting it together, we get

$$\nabla f(x) = (f_j - f_i)\nabla B_j(x) + (f_k - f_i)\nabla B_k(x)$$

- Note that $\nabla B_i(x) = \frac{N \times (x_k - x_j)}{2A_T}$, which gives

$$\nabla f(x) = f_i \frac{N \times (x_k - x_j)}{2A_T} + f_j \frac{N \times (x_i - x_k)}{2A_T} + f_k \frac{N \times (x_j - x_i)}{2A_T}$$




Figure : Three piece-wise linear basis per triangle. Image from BKPAL10.

- What is $\nabla B_i(x)$?
- ▶ $\nabla B_i(x) = \frac{(x_k - x_j)^\perp}{2A_T}$.
- Note that $B_i(x) + B_j(x) + B_k(x) = 1$ gives
 $\nabla B_i(x) = -(\nabla B_j(x) + \nabla B_k(x))$.
- ▶ Putting it together, we get

$$\nabla f(x) = (f_j - f_i)\nabla B_j(x) + (f_k - f_i)\nabla B_k(x)$$

- Note that $\nabla B_i(x) = \frac{N \times (x_k - x_j)}{2A_T}$, which gives

$$\nabla f(x) = f_i \frac{N \times (x_k - x_j)}{2A_T} + f_j \frac{N \times (x_i - x_k)}{2A_T} + f_k \frac{N \times (x_j - x_i)}{2A_T}$$

Surface Normal

- Surface normal for a triangle T with vertices (x_i, x_j, x_k) , given in AC order seen from outside of the surface:


$$N(T) = \frac{(x_j - x_i) \times (x_k - x_i)}{\|(x_j - x_i) \times (x_k - x_i)\|}.$$

- On a vertex v : Weighted average of unit normals of all incident triangles.

$$N(v) = \frac{\sum_{T \in N_1(v)} a_T N(T)}{\|\sum_{T \in N_1(v)} a_T N(T)\|},$$

where a_T 's are non-negative real scalars.

- *One-Ring*: Set of triangles incident to a given vertex v is called the one-ring of v and is denoted by $N_1(v)$.




Surface Normal

- Surface normal for a triangle T with vertices (x_i, x_j, x_k) , given in AC order seen from outside of the surface:
$$N(T) = \frac{(x_j - x_i) \times (x_k - x_i)}{\|(x_j - x_i) \times (x_k - x_i)\|}.$$
- On a vertex v : Weighted average of unit normals of all incident triangles.

$$N(v) = \frac{\sum_{T \in N_1(v)} a_T N(T)}{\|\sum_{T \in N_1(v)} a_T N(T)\|},$$

where a_T 's are non-negative real scalars.

- *One-Ring*: Set of triangles incident to a given vertex v is called the one-ring of v and is denoted by $N_1(v)$.




Surface Normal

- Surface normal for a triangle T with vertices (x_i, x_j, x_k) , given in AC order seen from outside of the surface:
$$N(T) = \frac{(x_j - x_i) \times (x_k - x_i)}{\|(x_j - x_i) \times (x_k - x_i)\|}.$$
- On a vertex v : Weighted average of unit normals of all incident triangles.

$$N(v) = \frac{\sum_{T \in N_1(v)} a_T N(T)}{\|\sum_{T \in N_1(v)} a_T N(T)\|},$$

where a_T 's are non-negative real scalars.

- *One-Ring*: Set of triangles incident to a given vertex v is called the one-ring of v and is denoted by $N_1(v)$.



Laplacian




Figure : Original signal f and interpolated signal \tilde{f} .

- For now let's assume $x_n = \Delta n$, where Δ is a uniform sampling interval.
- $\frac{d^2\tilde{f}}{dx^2}(x) = 0, \forall x \in (x_n, x_{n+1}).$
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{\frac{d\tilde{f}}{dx}\left(x_{n+\frac{1}{2}}\right) - \frac{d\tilde{f}}{dx}\left(x_{n-\frac{1}{2}}\right)}{\Delta} = \frac{\frac{f_{n+1}-f_n}{\Delta} - \frac{f_n-f_{n-1}}{\Delta}}{\Delta}.$$
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1} + f_{n-1} - 2f_n}{\Delta^2}.$$

Laplacian




Figure : Original signal f and interpolated signal \tilde{f} .

- For now let's assume $x_n = \Delta n$, where Δ is a uniform sampling interval.

- $\frac{d^2\tilde{f}}{dx^2}(x) = 0, \forall x \in (x_n, x_{n+1}).$

- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{\frac{d\tilde{f}}{dx}\left(x_{n+\frac{1}{2}}\right) - \frac{d\tilde{f}}{dx}\left(x_{n-\frac{1}{2}}\right)}{\Delta} = \frac{\frac{f_{n+1}-f_n}{\Delta}}{\Delta} - \frac{\frac{f_n-f_{n-1}}{\Delta}}{\Delta}.$$

- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1} + f_{n-1} - 2f_n}{\Delta^2}.$$

Laplacian




Figure : Original signal f and interpolated signal \tilde{f} .

- For now let's assume $x_n = \Delta n$, where Δ is a uniform sampling interval.
- $\frac{d^2\tilde{f}}{dx^2}(x) = 0, \forall x \in (x_n, x_{n+1}).$
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{\frac{d\tilde{f}}{dx}\left(x_{n+\frac{1}{2}}\right) - \frac{d\tilde{f}}{dx}\left(x_{n-\frac{1}{2}}\right)}{\Delta} = \frac{\frac{f_{n+1}-f_n}{\Delta} - \frac{f_n-f_{n-1}}{\Delta}}{\Delta}.$$
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1} + f_{n-1} - 2f_n}{\Delta^2}.$$

Laplacian




Figure : Original signal f and interpolated signal \tilde{f} .

- For now let's assume $x_n = \Delta n$, where Δ is a uniform sampling interval.
- $\frac{d^2\tilde{f}}{dx^2}(x) = 0, \forall x \in (x_n, x_{n+1})$.
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{\frac{d\tilde{f}}{dx}\left(x_{n+\frac{1}{2}}\right) - \frac{d\tilde{f}}{dx}\left(x_{n-\frac{1}{2}}\right)}{\Delta} = \frac{\frac{f_{n+1}-f_n}{\Delta} - \frac{f_n-f_{n-1}}{\Delta}}{\Delta}.$$
- $$\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1} + f_{n-1} - 2f_n}{\Delta^2}.$$

- $\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1}-f_n}{\Delta^2} + \frac{f_{n-1}-f_n}{\Delta^2}$




Figure : Signal domain interpreted as a graph.

- Denoting the Laplacian operator by L , we have

$$(L\tilde{f})_n = \sum_{j \in N(n)} w_{jn} (f_j - f_n), \quad w_{jn} = \frac{1}{\Delta^2}.$$

- Define $w_{jn} = \frac{1}{\Delta^2}$ if $j \in N(n)$, else $w_{jn} = 0$. We get

$$(L\tilde{f})_n = \sum_j w_{jn} (f_j - f_n).$$

- $\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1}-f_n}{\Delta^2} + \frac{f_{n-1}-f_n}{\Delta^2}$




Figure : Signal domain interpreted as a graph.

- Denoting the Laplacian operator by L , we have

$$(L\tilde{f})_n = \sum_{j \in N(n)} w_{jn} (f_j - f_n), \quad w_{jn} = \frac{1}{\Delta^2}.$$

► Define $w_{jn} = \frac{1}{\Delta^2}$ if $j \in N(n)$, else $w_{jn} = 0$. We get

$$(L\tilde{f})_n = \sum_j w_{jn} (f_j - f_n).$$

- $\frac{d^2\tilde{f}}{dx^2}(x_n) = \frac{f_{n+1}-f_n}{\Delta^2} + \frac{f_{n-1}-f_n}{\Delta^2}$




Figure : Signal domain interpreted as a graph.

- Denoting the Laplacian operator by L , we have

$$(L\tilde{f})_n = \sum_{j \in N(n)} w_{jn}(f_j - f_n), \quad w_{jn} = \frac{1}{\Delta^2}.$$

- Define $w_{jn} = \frac{1}{\Delta^2}$ if $j \in N(n)$, else $w_{jn} = 0$. We get

$$(L\tilde{f})_n = \sum_j w_{jn}(f_j - f_n).$$

- As a matrix, we get

$$\begin{bmatrix} | \\ (L\tilde{f})_n \\ | \end{bmatrix} = \begin{bmatrix} & & & | & & \\ - & w_{nn-1} & w_{nn} & w_{nn+1} & - & \\ & & | & & & \end{bmatrix} \begin{bmatrix} | \\ f_{n-1} \\ f_n \\ f_{n+1} \\ | \end{bmatrix},$$

where $w_{n,n} = -\sum_j w_{jn}$.

- The matrix A such that $A(i,j) = w_{ij}$ with $A(i,i) = 0$ is called the *Adjacency matrix* of the graph, while the diagonal matrix D with $D(i,i) = \sum_j A(i,j)$ is called the *Degree matrix* of the graph.
- Thus $L = A - D$.
- In graph theory, $\tilde{L} = D - A$ is called the *Graph Laplacian*.

- As a matrix, we get

$$\begin{bmatrix} | \\ (L\tilde{f})_n \\ | \end{bmatrix} = \begin{bmatrix} & & & | & & \\ - & w_{nn-1} & w_{nn} & w_{nn+1} & - & \\ & & | & & & \end{bmatrix} \begin{bmatrix} | \\ f_{n-1} \\ f_n \\ f_{n+1} \\ | \end{bmatrix},$$

where $w_{n,n} = -\sum_j w_{jn}$.

- The matrix A such that $A(i,j) = w_{ij}$ with $A(i,i) = 0$ is called the *Adjacency matrix* of the graph, while the diagonal matrix D with $D(i,i) = \sum_j A(i,j)$ is called the *Degree matrix* of the graph.
- Thus $L = A - D$.
- In graph theory, $\tilde{L} = D - A$ is called the *Graph Laplacian*.

- As a matrix, we get

$$\begin{bmatrix} | \\ (L\tilde{f})_n \\ | \end{bmatrix} = \begin{bmatrix} & & & | & & \\ - & w_{nn-1} & w_{nn} & w_{nn+1} & - & \\ & & | & & & \end{bmatrix} \begin{bmatrix} | \\ f_{n-1} \\ f_n \\ f_{n+1} \\ | \end{bmatrix},$$

where $w_{n,n} = -\sum_j w_{jn}$.

- The matrix A such that $A(i,j) = w_{ij}$ with $A(i,i) = 0$ is called the *Adjacency matrix* of the graph, while the diagonal matrix D with $D(i,i) = \sum_j A(i,j)$ is called the *Degree matrix* of the graph.
- Thus $L = A - D$.
- In graph theory, $\tilde{L} = D - A$ is called the *Graph Laplacian*.

- As a matrix, we get

$$\begin{bmatrix} | \\ (L\tilde{f})_n \\ | \end{bmatrix} = \begin{bmatrix} & & & | & & \\ - & w_{nn-1} & w_{nn} & w_{nn+1} & - & \\ & & | & & & \end{bmatrix} \begin{bmatrix} | \\ f_{n-1} \\ f_n \\ f_{n+1} \\ | \end{bmatrix},$$

where $w_{n,n} = -\sum_j w_{jn}$.

- The matrix A such that $A(i,j) = w_{ij}$ with $A(i,i) = 0$ is called the *Adjacency matrix* of the graph, while the diagonal matrix D with $D(i,i) = \sum_j A(i,j)$ is called the *Degree matrix* of the graph.
- Thus $L = A - D$.
- In graph theory, $\tilde{L} = D - A$ is called the *Graph Laplacian*.

Laplacian for meshes

- A mesh is a graph; note $\Delta = d(x_n, x_{n+1})$.




Figure : Mesh as a graph, $d_{jn} = \|x_j - x_n\|^2$

- If vertex i and j are edge-connected $A(i,j) = \frac{1}{d_{i,j}}$, else $A(i,j) = 0$.
- ▶ Construct the degree matrix D with $D(i,i) = \sum_j A(i,j)$, and the Laplacian $L = A - D$.

Laplacian for meshes

- A mesh is a graph; note $\Delta = d(x_n, x_{n+1})$.




Figure : Mesh as a graph, $d_{jn} = \|x_j - x_n\|^2$

- If vertex i and j are edge-connected $A(i,j) = \frac{1}{d_{i,j}}$, else $A(i,j) = 0$.
- ▶ Construct the degree matrix D with $D(i,i) = \sum_j A(i,j)$, and the Laplacian $L = A - D$.

Laplacian for meshes

- A mesh is a graph; note $\Delta = d(x_n, x_{n+1})$.




Figure : Mesh as a graph, $d_{jn} = \|x_j - x_n\|^2$

- If vertex i and j are edge-connected $A(i,j) = \frac{1}{d_{i,j}}$, else $A(i,j) = 0$.
- ▶ Construct the degree matrix D with $D(i,i) = \sum_j A(i,j)$, and the Laplacian $L = A - D$.

Thumb rules for discrete estimation

- Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- Order of error: Finer sampling \Rightarrow Error should go down.

For example, if the function is smooth enough, then the error of the estimate is proportional to the square of the width of the bins.

Thumb rules for discrete estimation

- ▶ Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- ▶ Order of error: Finer sampling \Rightarrow Error should go down.
- ▶ Estimate should be robust.

Thumb rules for discrete estimation

- ▶ Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- ▶ Order of error: Finer sampling \Rightarrow Error should go down.
- ▶ Estimate should be robust.

Thumb rules for discrete estimation

- ▶ Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- ▶ Order of error: Finer sampling \Rightarrow Error should go down.
- ▶ Estimate should be robust. Robustness \rightarrow average (a.k.a low pass filter).

Thumb rules for discrete estimation

- ▶ Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- ▶ Order of error: Finer sampling \Rightarrow Error should go down.
- ▶ Estimate should be robust. Robustness \rightarrow average (a.k.a low pass filter).

Thumb rules for discrete estimation

- ▶ Properties of the measure in the continuous setting should be satisfied by the discrete estimate.
- ▶ Order of error: Finer sampling \Rightarrow Error should go down.
- ▶ Estimate should be robust. Robustness \rightarrow average (a.k.a low pass filter).

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \text{div} \circ \text{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:
 - ▶ We know that $A(\sigma) = \iint \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
 - ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
 - ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \operatorname{div} \circ \operatorname{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:
 - ▶ We know that $A(\sigma) = \iint \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
 - ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
 - ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \operatorname{div} \circ \operatorname{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:

- ▶ We know that $A(\sigma) = \iint \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
- ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
- ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \operatorname{div} \circ \operatorname{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:
 - ▶ We know that $A(\sigma) = \int \int \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
 - ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
 - ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \operatorname{div} \circ \operatorname{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:
 - ▶ We know that $A(\sigma) = \int \int \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
 - ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
 - ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

Laplace-Beltrami Operator (LBO)

- **Definition 1:** $\Delta := \operatorname{div} \circ \operatorname{grad}$.
- In \mathbb{R}^n , $\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$, LBO = Laplacian.
- Properties/Characteristics of LBO:
 - ▶ We know that $A(\sigma) = \int \int \sqrt{EG - F^2} da$, for $\sigma : \Omega \subset \mathbb{R}^2 \rightarrow S \subset \mathbb{R}^3$.
 - ▶ For $\sigma(u, v) = (u, v, f(u, v))$, $EG - F^2 = 1 + f_u^2 + f_v^2$.
 - ▶ Characterize an σ^* such that $A(\sigma^*)$ is minimum, given a fixed boundary $\sigma|_{\partial\Omega} = \Gamma$.

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = -\frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
 - Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
 - As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
 - It can be derived that

$$\nabla A(f) = -\frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$

- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes

$$\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v)), \text{ with } g|_{\partial\Omega} = 0.$$

- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
- Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
- As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
- It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$


- Let $g : \Omega \rightarrow \mathbb{R}$ be a perturbation added to f by a scalar amount s , so that the surface σ becomes $\tilde{\sigma}_s = (u, v, f(u, v) + sg(u, v))$, with $g|_{\partial\Omega} = 0$.
- One can then define the Directional derivative of A at f in the direction g as $D_g A(f) := \frac{d}{ds} A(f + sg)|_{s=0}$.
 - Another notation for DD: $\frac{\partial A}{\partial g}(f)$.
 - As in finite dimensional cases, $\frac{\partial A}{\partial g}(f) = \langle \nabla A(f), g \rangle$.
 - It can be derived that

$$\nabla A(f) = - \frac{(1 + f_v^2)f_{uu} + (1 + f_u^2)f_{vv} - 2f_u f_v f_{uv}}{(1 + f_u^2 + f_v^2)^{\frac{3}{2}}}.$$

- Thus f^* represents a minimal surface if $\nabla A(f^*) = 0$.
- Note $E = 1 + f_u^2$, $F = f_u f_v$, $G = 1 + f_v^2$.
- Also

$$L = \langle \sigma_{uu}, n \rangle = \frac{f_{uu}}{\sqrt{1+f_u^2+f_v^2}}, M = \frac{f_{uv}}{\sqrt{1+f_u^2+f_v^2}}, N = \frac{f_{vv}}{\sqrt{1+f_u^2+f_v^2}}.$$


- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.



Minimal surfaces are surfaces that minimize their area given boundary conditions.


Minimal surfaces are surfaces that minimize their area given boundary conditions.

- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.




Minimal surfaces are surfaces that locally minimize their area.

- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.



- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.



- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.




Figure : Area minimization surface evolution

The famous Mean Curvature Flow $\frac{d}{dt}f = 2Hn$

- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
- Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- ▶ Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- ▶ Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.




Figure : Area minimization surface evolution

The famous Mean Curvature Flow $\frac{d}{dt}f = 2Hn$

- Thus $\nabla A(f) = -\frac{GL+EN-2FM}{EG-F^2}$.
 - Note $H = \text{Tr}([S]) = \text{Tr}(\mathcal{F}_1^{-1}\mathcal{F}_2) = \frac{GL+EN-2FM}{2(EG-F^2)}$.
- Thus, $\nabla A(f) = -2H$.
- Given a surface σ (or f), how can one turn it into a minimal surface?
- Gradient descent: $\frac{\partial f}{\partial t} = -\nabla A(f)$, or $\frac{\partial \sigma}{\partial t} = -\nabla A(f)\hat{z}$, where $\hat{z} = (0, 0, 1)$.




Figure : Area minimization surface evolution

- The famous **Mean Curvature Flow**: $\frac{\partial \sigma}{\partial t} = 2Hn$.

What has this got to do with LBO?

- *Isothermal or Conformal parameterization:* If $F = 0$ and $E = G$ at every point of the surface, we say that the parameters are *isothermal* or *conformal*.
 - ▶ Assuming such parameters¹ u, v exist and $E = G = 1$, $H = L + N$.
 - ▶ In this case, $\langle \sigma_u, \sigma_u \rangle = 1$, thus $\sigma_{uu} + \sigma_{vv} \parallel n$.
 - ▶ Conformal parameterization $\Rightarrow \Delta \sigma = Hn$.
- **Definition 2²:** LBO is the operator that assigns to each point $p \in M$ the *mean curvature vector* $H(p)n(p)$.

¹Gauss(1822) proved existence of local conformal parameterization for any 2D surface, while global existence was shown by Lichtenstein (1916).

²Meyer et al., Discrete Differential-Geometry Operators for Triangulated 2-Manifolds, Intl. w'shop on Vis. & Math., 2002

What has this got to do with LBO?

- *Isothermal or Conformal parameterization:* If $F = 0$ and $E = G$ at every point of the surface, we say that the parameters are *isothermal* or *conformal*.
- ▶ Assuming such parameters¹ u, v exist and $E = G = 1$, $H = L + N$.
- ▶ In this case, $\langle \sigma_u, \sigma_u \rangle = 1$, thus $\sigma_{uu} + \sigma_{vv} \parallel n$.
- ▶ Conformal parameterization $\Rightarrow \Delta \sigma = Hn$.
- **Definition 2²:** LBO is the operator that assigns to each point $p \in M$ the *mean curvature vector* $H(p)n(p)$.

¹Gauss(1822) proved existence of local conformal parameterization for any 2D surface, while global existence was shown by Lichtenstein (1916).

²Meyer et al., Discrete Differential-Geometry Operators for Triangulated 2-Manifolds, Intl. w'shop on Vis. & Math., 2002

What has this got to do with LBO?

- *Isothermal or Conformal parameterization*: If $F = 0$ and $E = G$ at every point of the surface, we say that the parameters are *isothermal* or *conformal*.
- ▶ Assuming such parameters¹ u, v exist and $E = G = 1$, $H = L + N$.
- ▶ In this case, $\langle \sigma_u, \sigma_u \rangle = 1$, thus $\sigma_{uu} + \sigma_{vv} \parallel n$.
- ▶ Conformal parameterization $\Rightarrow \Delta \sigma = Hn$.
- **Definition 2²**: LBO is the operator that assigns to each point $p \in M$ the *mean curvature vector* $H(p)n(p)$.

¹Gauss(1822) proved existence of local conformal parameterization for any 2D surface, while global existence was shown by Lichtenstein (1916).

²Meyer et al., Discrete Differential-Geometry Operators for Triangulated 2-Manifolds, Intl. w'shop on Vis. & Math., 2002

What has this got to do with LBO?

- *Isothermal or Conformal parameterization:* If $F = 0$ and $E = G$ at every point of the surface, we say that the parameters are *isothermal* or *conformal*.
- ▶ Assuming such parameters¹ u, v exist and $E = G = 1$, $H = L + N$.
- ▶ In this case, $\langle \sigma_u, \sigma_u \rangle = 1$, thus $\sigma_{uu} + \sigma_{vv} \parallel n$.
- ▶ Conformal parameterization $\Rightarrow \Delta\sigma = Hn$.
- **Definition 2²:** LBO is the operator that assigns to each point $p \in M$ the *mean curvature vector* $H(p)n(p)$.

¹Gauss(1822) proved existence of local conformal parameterization for any 2D surface, while global existence was shown by Lichtenstein (1916).

²Meyer et al., Discrete Differential-Geometry Operators for Triangulated 2-Manifolds, Intl. w'shop on Vis. & Math., 2002

What has this got to do with LBO?

- *Isothermal or Conformal parameterization*: If $F = 0$ and $E = G$ at every point of the surface, we say that the parameters are *isothermal* or *conformal*.
- ▶ Assuming such parameters¹ u, v exist and $E = G = 1$, $H = L + N$.
- ▶ In this case, $\langle \sigma_u, \sigma_u \rangle = 1$, thus $\sigma_{uu} + \sigma_{vv} \parallel n$.
- ▶ Conformal parameterization $\Rightarrow \Delta\sigma = Hn$.
- **Definition 2²**: LBO is the operator that assigns to each point $p \in M$ the *mean curvature vector* $H(p)n(p)$.

¹Gauss(1822) proved existence of local conformal parameterization for any 2D surface, while global existence was shown by Lichtenstein (1916).

²Meyer et al., Discrete Differential-Geometry Operators for Triangulated 2-Manifolds, Intl. w'shop on Vis. & Math., 2002

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \iint \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \iint \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \iint \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta\sigma = 0$.
- ▶ Functions that satisfy³ $\Delta\sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \int \int \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \int \int \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \int \int \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta \sigma = 0$.
- ▶ Functions that satisfy³ $\Delta \sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \int \int \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \int \int \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \int \int \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta \sigma = 0$.
- ▶ Functions that satisfy³ $\Delta \sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \int \int \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \int \int \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \int \int \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta \sigma = 0$.
- ▶ Functions that satisfy³ $\Delta \sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \int \int \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \int \int \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \int \int \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta\sigma = 0$.
- ▶ Functions that satisfy³ $\Delta\sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Dirichlet Energy

- A parameterization σ of a surface maps points from \mathbb{R}^2 to points in \mathbb{R}^3 .
- ▶ A measure of how rapidly σ varies:

$$E(\sigma) = \frac{1}{2} \int \int \left\| \frac{\partial \sigma}{\partial u} \right\|^2 + \left\| \frac{\partial \sigma}{\partial v} \right\|^2 du dv$$

- ▶ Typically written as $E(\sigma) = \frac{1}{2} \int \int \|\nabla \sigma\|^2 du dv$ or $E(\sigma) = \frac{1}{2} \int \int \text{Tr}(J\sigma^t J\sigma) du dv$, and is called the *Dirichlet Energy*.
- ▶ With conformal parameterization $\sqrt{EG - F^2} = E$, thus $A(\sigma) = E(\sigma)$.
- ▶ The minimizer of $E(\sigma)$ also satisfies $\nabla A(\sigma) = Hn = \Delta\sigma = 0$.
- ▶ Functions that satisfy³ $\Delta\sigma = 0$ are called *Harmonic functions*.

³This is the famous Laplace equation.

Self-Adjoint operator

- Let $V := \{f : [0, 2\pi) \rightarrow \mathbb{R} \mid f \text{ is smooth and periodic}\}$, equipped with the standard L_2 inner product.
- Note the following:

$$\int_0^{2\pi} \Delta f(x) g(x) dx = [\nabla f(x)g(x)]_0^{2\pi} - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx$$

Since $f, g \in V, \nabla f(0) = \nabla f(2\pi), g(0) = g(2\pi)$

$$\begin{aligned} \int_0^{2\pi} \Delta f(x) g(x) dx &= - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx \\ &= - [f(x)g(x)]_0^{2\pi} + \int_0^{2\pi} f(x)\Delta g(x) dx \\ &= \int_0^{2\pi} f(x)\Delta g(x) dx \end{aligned}$$

- In terms of inner product, we get

$$\langle \Delta f, g \rangle_{L_2} = \langle f, \Delta g \rangle_{L_2} \Rightarrow \Delta^* = \Delta.$$

Self-Adjoint operator

- Let $V := \{f : [0, 2\pi) \rightarrow \mathbb{R} \mid f \text{ is smooth and periodic}\}$, equipped with the standard L_2 inner product.
- Note the following:

$$\int_0^{2\pi} \Delta f(x) g(x) dx = [\nabla f(x)g(x)]_0^{2\pi} - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx$$

Since $f, g \in V, \nabla f(0) = \nabla f(2\pi), g(0) = g(2\pi)$

$$\begin{aligned} \int_0^{2\pi} \Delta f(x) g(x) dx &= - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx \\ &= - [f(x)g(x)]_0^{2\pi} + \int_0^{2\pi} f(x)\Delta g(x) dx \\ &= \int_0^{2\pi} f(x)\Delta g(x) dx \end{aligned}$$

- In terms of inner product, we get

$$\langle \Delta f, g \rangle_{L_2} = \langle f, \Delta g \rangle_{L_2} \Rightarrow \Delta^* = \Delta.$$

Self-Adjoint operator

- Let $V := \{f : [0, 2\pi) \rightarrow \mathbb{R} \mid f \text{ is smooth and periodic}\}$, equipped with the standard L_2 inner product.
- Note the following:

$$\int_0^{2\pi} \Delta f(x) g(x) dx = [\nabla f(x)g(x)]_0^{2\pi} - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx$$

Since $f, g \in V, \nabla f(0) = \nabla f(2\pi), g(0) = g(2\pi)$

$$\begin{aligned} \int_0^{2\pi} \Delta f(x) g(x) dx &= - \int_0^{2\pi} \nabla f(x)\nabla g(x) dx \\ &= - [f(x)g(x)]_0^{2\pi} + \int_0^{2\pi} f(x)\Delta g(x) dx \\ &= \int_0^{2\pi} f(x)\Delta g(x) dx \end{aligned}$$

- In terms of inner product, we get

$$\langle \Delta f, g \rangle_{L_2} = \langle f, \Delta g \rangle_{L_2} \Rightarrow \Delta^* = \Delta.$$

- Thus Δ can be diagonalized, and for the vector space V in the previous slide, the eigenfunctions of Δ are:
 $e_n(x) = \sin(nx)$ or $e_n(x) = \cos(nx), \forall n \in \mathbb{Z}$.

If complex valued functions are allowed, we have the familiar exponential Fourier series basis:

$$e_n(x) = \exp(jnx), \forall n \in \mathbb{Z}$$

- Thus Δ can be diagonalized, and for the vector space V in the previous slide, the eigenfunctions of Δ are:
 $e_n(x) = \sin(nx)$ or $e_n(x) = \cos(nx), \forall n \in \mathbb{Z}$.

If complex valued functions are allowed, we have the familiar exponential Fourier series basis:

$$e_n(x) = \exp(jnx), \forall n \in \mathbb{Z}$$

- Thus Δ can be diagonalized, and for the vector space V in the previous slide, the eigenfunctions of Δ are:
 $e_n(x) = \sin(nx)$ or $e_n(x) = \cos(nx), \forall n \in \mathbb{Z}$.
- ▶ If complex valued functions are allowed, we have the familiar exponential Fourier series basis:

$$e_n(x) = \exp(jnx), \forall n \in \mathbb{Z}.$$

Eigenfunctions of discrete LBO




Figure : Can you interpret these eigenfunctions?

Deriving discrete LBO

- **Derivation 1.:** Minimizing Dirichlet energy⁴.

- Let a surface be parameterized by the function f ; the corresponding Dirichlet energy is

$$E_D(f) = \frac{1}{2} \int_{\Omega} \|\nabla f\|^2 \ da$$

⁴Pinkall & Polthier, Computing Discrete Minimal Surfaces & their conjugates, Experimental Mathematics, vol. 2, no. 1, 1993

Deriving discrete LBO

- **Derivation 1.:** Minimizing Dirichlet energy⁴.
- Let a surface be parameterized by the function f ; the corresponding Dirichlet energy is

$$E_D(f) = \frac{1}{2} \int_{\Omega} \|\nabla f\|^2 \ da$$

⁴Pinkall & Polthier, Computing Discrete Minimal Surfaces & their conjugates, Experimental Mathematics, vol. 2, no. 1, 1993

- Let us look at E_D triangle-wise.




Figure : Triangles in local coordinates mapping to Triangles on the surface.

- For a discrete triangulated surface, we will assume that f is defined at vertices, while at all other points we will assume f to be linear.
- Moreover, $E_D(f) = \sum_{i=1}^m E_D(f_i)$, where m is the total number of triangles, and f_i is a mapping between corresponding triangles i .

- Let us look at E_D triangle-wise.




Figure : Triangles in local coordinates mapping to Triangles on the surface.

- For a discrete triangulated surface, we will assume that f is defined at vertices, while at all other points we will assume f to be linear.
- Moreover, $E_D(f) = \sum_{i=1}^m E_D(f_i)$, where m is the total number of triangles, and f_i is a mapping between corresponding triangles i .

- Let us look at E_D triangle-wise.




Figure : Triangles in local coordinates mapping to Triangles on the surface.

- For a discrete triangulated surface, we will assume that f is defined at vertices, while at all other points we will assume f to be linear.
- Moreover, $E_D(f) = \sum_{i=1}^m E_D(f_i)$, where m is the total number of triangles, and f_i is a mapping between corresponding triangles i .

- The linear map $f_i : \Delta_1 \rightarrow \Delta_2$ is defined by $f(v) = a, f(w) = b$.




Figure : f_i between two corresponding triangles.

- The map f can also be written as $f = \psi \circ \phi^{-1}$, as shown in the following figure.




Figure : The map f via the standard triangle.

- Using Chain rule & Inverse Function theorem,

$$Jf = J\psi \circ J(\phi^{-1}) = J\psi (J\phi)^{-1}.$$

- Then,

$$\begin{aligned} Tr(Jf^t Jf) &= Tr((J\psi (J\phi)^{-1})^t J\psi (J\phi)^{-1}) \\ &= Tr(((J\phi)^{-1})^t J\psi^t J\psi (J\phi)^{-1}) \\ &= Tr(J\psi^t J\psi (J\phi)^{-1} ((J\phi)^{-1})^t) \end{aligned}$$

- Note that $J\psi = [a \ b]$, thus $J\psi^t J\psi = \begin{bmatrix} \langle a, a \rangle & \langle a, b \rangle \\ \langle b, a \rangle & \langle b, b \rangle \end{bmatrix}$.
- Similarly, $J\phi = [v \ w]$, thus

$$(J\phi)^{-1}((J\phi)^{-1})^t = ((J\phi)^t J\phi)^{-1} = \begin{bmatrix} \langle v, v \rangle & \langle v, w \rangle \\ \langle w, v \rangle & \langle w, w \rangle \end{bmatrix}^{-1}.$$

- Let $d = \det(J\phi)$ and $c = b - a$.

$$\begin{aligned} Tr(Jf^t Jf) &= \frac{1}{d} \left(\frac{\langle -w, v - w \rangle}{d} \|a\|^2 + \frac{\langle w - v, -v \rangle}{d} \|b\|^2 \right. \\ &\quad \left. + \frac{\langle v, w \rangle}{d} \|c\|^2 \right) \end{aligned}$$

- Using Chain rule & Inverse Function theorem,

$$Jf = J\psi \circ J(\phi^{-1}) = J\psi (J\phi)^{-1}.$$

- Then,

$$\begin{aligned} Tr(Jf^t Jf) &= Tr((J\psi (J\phi)^{-1})^t J\psi (J\phi)^{-1}) \\ &= Tr(((J\phi)^{-1})^t J\psi^t J\psi (J\phi)^{-1}) \\ &= Tr(J\psi^t J\psi (J\phi)^{-1} ((J\phi)^{-1})^t) \end{aligned}$$

- Note that $J\psi = [a \ b]$, thus $J\psi^t J\psi = \begin{bmatrix} \langle a, a \rangle & \langle a, b \rangle \\ \langle b, a \rangle & \langle b, b \rangle \end{bmatrix}$.

- Similarly, $J\phi = [v \ w]$, thus

$$(J\phi)^{-1}((J\phi)^{-1})^t = ((J\phi)^t J\phi)^{-1} = \begin{bmatrix} \langle v, v \rangle & \langle v, w \rangle \\ \langle w, v \rangle & \langle w, w \rangle \end{bmatrix}^{-1}.$$

- Let $d = \det(J\phi)$ and $c = b - a$.

$$\begin{aligned} Tr(Jf^t Jf) &= \frac{1}{d} \left(\frac{\langle -w, v - w \rangle}{d} \|a\|^2 + \frac{\langle w - v, -v \rangle}{d} \|b\|^2 \right. \\ &\quad \left. + \frac{\langle v, w \rangle}{d} \|c\|^2 \right) \end{aligned}$$

- Using Chain rule & Inverse Function theorem,

$$Jf = J\psi \circ J(\phi^{-1}) = J\psi (J\phi)^{-1}.$$

- Then,

$$\begin{aligned} Tr(Jf^t Jf) &= Tr((J\psi (J\phi)^{-1})^t J\psi (J\phi)^{-1}) \\ &= Tr(((J\phi)^{-1})^t J\psi^t J\psi (J\phi)^{-1}) \\ &= Tr(J\psi^t J\psi (J\phi)^{-1} ((J\phi)^{-1})^t) \end{aligned}$$

- Note that $J\psi = [a \ b]$, thus $J\psi^t J\psi = \begin{bmatrix} \langle a, a \rangle & \langle a, b \rangle \\ \langle b, a \rangle & \langle b, b \rangle \end{bmatrix}$.
- Similarly, $J\phi = [v \ w]$, thus

$$(J\phi)^{-1}((J\phi)^{-1})^t = ((J\phi)^t J\phi)^{-1} = \begin{bmatrix} \langle v, v \rangle & \langle v, w \rangle \\ \langle w, v \rangle & \langle w, w \rangle \end{bmatrix}^{-1}.$$

- Let $d = \det(J\phi)$ and $c = b - a$.

$$\begin{aligned} Tr(Jf^t Jf) &= \frac{1}{d} \left(\frac{\langle -w, v - w \rangle}{d} \|a\|^2 + \frac{\langle w - v, -v \rangle}{d} \|b\|^2 \right. \\ &\quad \left. + \frac{\langle v, w \rangle}{d} \|c\|^2 \right) \end{aligned}$$

- Using Chain rule & Inverse Function theorem,

$$Jf = J\psi \circ J(\phi^{-1}) = J\psi (J\phi)^{-1}.$$

- Then,

$$\begin{aligned} Tr(Jf^t Jf) &= Tr((J\psi (J\phi)^{-1})^t J\psi (J\phi)^{-1}) \\ &= Tr(((J\phi)^{-1})^t J\psi^t J\psi (J\phi)^{-1}) \\ &= Tr(J\psi^t J\psi (J\phi)^{-1} ((J\phi)^{-1})^t) \end{aligned}$$

- Note that $J\psi = [a \ b]$, thus $J\psi^t J\psi = \begin{bmatrix} \langle a, a \rangle & \langle a, b \rangle \\ \langle b, a \rangle & \langle b, b \rangle \end{bmatrix}$.
- Similarly, $J\phi = [v \ w]$, thus

$$(J\phi)^{-1}((J\phi)^{-1})^t = ((J\phi)^t J\phi)^{-1} = \begin{bmatrix} \langle v, v \rangle & \langle v, w \rangle \\ \langle w, v \rangle & \langle w, w \rangle \end{bmatrix}^{-1}.$$

- Let $d = \det(J\phi)$ and $c = b - a$.

$$\begin{aligned} Tr(Jf^t Jf) &= \frac{1}{d} \left(\frac{\langle -w, v - w \rangle}{d} \|a\|^2 + \frac{\langle w - v, -v \rangle}{d} \|b\|^2 \right. \\ &\quad \left. + \frac{\langle v, w \rangle}{d} \|c\|^2 \right) \end{aligned}$$

- Using Chain rule & Inverse Function theorem,

$$Jf = J\psi \circ J(\phi^{-1}) = J\psi (J\phi)^{-1}.$$

- Then,

$$\begin{aligned} Tr(Jf^t Jf) &= Tr((J\psi (J\phi)^{-1})^t J\psi (J\phi)^{-1}) \\ &= Tr(((J\phi)^{-1})^t J\psi^t J\psi (J\phi)^{-1}) \\ &= Tr(J\psi^t J\psi (J\phi)^{-1} ((J\phi)^{-1})^t) \end{aligned}$$

- Note that $J\psi = [a \ b]$, thus $J\psi^t J\psi = \begin{bmatrix} \langle a, a \rangle & \langle a, b \rangle \\ \langle b, a \rangle & \langle b, b \rangle \end{bmatrix}$.

- Similarly, $J\phi = [v \ w]$, thus

$$(J\phi)^{-1}((J\phi)^{-1})^t = ((J\phi)^t J\phi)^{-1} = \begin{bmatrix} \langle v, v \rangle & \langle v, w \rangle \\ \langle w, v \rangle & \langle w, w \rangle \end{bmatrix}^{-1}.$$

- Let $d = \det(J\phi)$ and $c = b - a$.

$$\begin{aligned} Tr(Jf^t Jf) &= \frac{1}{d} \left(\frac{\langle -w, v - w \rangle}{d} \|a\|^2 + \frac{\langle w - v, -v \rangle}{d} \|b\|^2 \right. \\ &\quad \left. + \frac{\langle v, w \rangle}{d} \|c\|^2 \right) \end{aligned}$$

- With $d = 2A$, and $d = ||v|| ||w|| \sin \theta_1 = ||w|| ||v - w|| \sin \theta_2 = ||w - v|| ||v|| \sin \theta_3$, we get

$$Tr(Jf^t Jf) = \frac{1}{2A} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2$$

- The Dirichlet energy is given by

$$\begin{aligned} E_D(f_i) &= \frac{1}{2} \int_{\Delta_1} Tr(Jf_i^t Jf_i) da \\ &= \frac{1}{2} Tr(Jf_i^t Jf_i) \int_{\Delta_1} da \\ &= \frac{1}{4} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2 \end{aligned}$$

- The total Dirichlet energy is then,

$$E_D(f) = \sum_{i=1}^m E_D(f_i)$$

- With $d = 2A$, and $d = ||v|| ||w|| \sin \theta_1 = ||w|| ||v - w|| \sin \theta_2 = ||w - v|| ||v|| \sin \theta_3$, we get

$$Tr(Jf^t Jf) = \frac{1}{2A} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2$$

- The Dirichlet energy is given by

$$\begin{aligned} E_D(f_i) &= \frac{1}{2} \int_{\Delta_1} Tr(Jf_i^t Jf_i) da \\ &= \frac{1}{2} Tr(Jf_i^t Jf_i) \int_{\Delta_1} da \\ &= \frac{1}{4} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2 \end{aligned}$$

- The total Dirichlet energy is then,

$$E_D(f) = \sum_{i=1}^m E_D(f_i)$$

- With $d = 2A$, and $d = ||v|| ||w|| \sin \theta_1 = ||w|| ||v - w|| \sin \theta_2 = ||w - v|| ||v|| \sin \theta_3$, we get

$$Tr(Jf^t Jf) = \frac{1}{2A} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2$$

- The Dirichlet energy is given by

$$\begin{aligned} E_D(f_i) &= \frac{1}{2} \int_{\Delta_1} Tr(Jf_i^t Jf_i) da \\ &= \frac{1}{2} Tr(Jf_i^t Jf_i) \int_{\Delta_1} da \\ &= \frac{1}{4} \sum_{i=1}^3 \cot \alpha_i ||a_i||^2 \end{aligned}$$

- The total Dirichlet energy is then,

$$E_D(f) = \sum_{i=1}^m E_D(f_i)$$

- To minimize Dirichlet energy, we differentiate w.r.t a vertex position and set it to zero:

$$\frac{\partial}{\partial p} E_D(f) = \frac{1}{2} \sum_{q \in N(p)} (\cot \alpha + \cot \beta)(p - q) = 0$$

- The minimizer can be interpreted as satisfying the (discrete) Laplace's equation: $Lf = (D - A)f = 0$, thus giving us the adjacency relation:

$$A_{ij} = \frac{1}{2}(\cot \alpha_{ij} + \cot \beta_{ij})$$




Figure : Weights for the Cotan Laplacian

- To minimize Dirichlet energy, we differentiate w.r.t a vertex position and set it to zero:

$$\frac{\partial}{\partial p} E_D(f) = \frac{1}{2} \sum_{q \in N(p)} (\cot \alpha + \cot \beta)(p - q) = 0$$

- The minimizer can be interpreted as satisfying the (discrete) Laplace's equation: $Lf = (D - A)f = 0$, thus giving us the adjacency relation:

$$A_{ij} = \frac{1}{2}(\cot \alpha_{ij} + \cot \beta_{ij})$$




Figure : Weights for the Cotan Laplacian

Derivation 2: Weak formulation of the Poisson equation

- Compute solution to the Poisson's equation: $\Delta u = f$, i.e., given f , compute u .
- ▶ The domain is a mesh, and the space of functions we work with is $V := \{u = \sum_j u_j B_j\}$, where B_j 's are the linear functions with value 1 at vertex j , else 0, as shown below.




Figure : Piece-wise linear

- ▶ Given f , can we compute $\hat{u} \in V$ as the best approximation to the actual solution u ?
- ▶ Idea 1: Given f , can we compute $\hat{u} \in V$ such that the laplacian of \hat{u} best approximates the function f ?
- ▶ Let error $e = \Delta \hat{u} - f$. We want $e \in V^\perp$.
 $\Rightarrow \langle \Delta \hat{u} - f, B_i \rangle = 0, \forall i$.

Derivation 2: Weak formulation of the Poisson equation

- Compute solution to the Poisson's equation: $\Delta u = f$, i.e., given f , compute u .
- ▶ The domain is a mesh, and the space of functions we work with is $V := \{u = \sum_j u_j B_j\}$, where B_j 's are the linear functions with value 1 at vertex j , else 0, as shown below.




Figure : Piece-wise linear

- ▶ Given f , can we compute $\hat{u} \in V$ as the best approximation to the actual solution u ?
- ▶ Idea 1: Given f , can we compute $\hat{u} \in V$ such that the laplacian of \hat{u} best approximates the function f ?
- ▶ Let error $e = \Delta \hat{u} - f$. We want $e \in V^\perp$.
 $\Rightarrow \langle \Delta \hat{u} - f, B_i \rangle = 0, \forall i$.

Derivation 2: Weak formulation of the Poisson equation

- Compute solution to the Poisson's equation: $\Delta u = f$, i.e., given f , compute u .
- ▶ The domain is a mesh, and the space of functions we work with is $V := \{u = \sum_j u_j B_j\}$, where B_j 's are the linear functions with value 1 at vertex j , else 0, as shown below.




Figure : Piece-wise linear

- ▶ Given f , can we compute $\hat{u} \in V$ as the best approximation to the actual solution u ?
- ▶ Idea 1: Given f , can we compute $\hat{u} \in V$ such that the laplacian of \hat{u} best approximates the function f ?
- ▶ Let error $e = \Delta \hat{u} - f$. We want $e \in V^\perp$.
 $\Rightarrow \langle \Delta \hat{u} - f, B_i \rangle = 0, \forall i$.

Derivation 2: Weak formulation of the Poisson equation

- Compute solution to the Poisson's equation: $\Delta u = f$, i.e., given f , compute u .
- ▶ The domain is a mesh, and the space of functions we work with is $V := \{u = \sum_j u_j B_j\}$, where B_j 's are the linear functions with value 1 at vertex j , else 0, as shown below.




Figure : Piece-wise linear

- ▶ Given f , can we compute $\hat{u} \in V$ as the best approximation to the actual solution u ?
- ▶ Idea 1: Given f , can we compute $\hat{u} \in V$ such that the laplacian of \hat{u} best approximates the function f ?
- ▶ Let error $e = \Delta \hat{u} - f$. We want $e \in V^\perp$.
 $\Rightarrow \langle \Delta \hat{u} - f, B_i \rangle = 0, \forall i$.

Derivation 2: Weak formulation of the Poisson equation

- Compute solution to the Poisson's equation: $\Delta u = f$, i.e., given f , compute u .
- ▶ The domain is a mesh, and the space of functions we work with is $V := \{u = \sum_j u_j B_j\}$, where B_j 's are the linear functions with value 1 at vertex j , else 0, as shown below.




Figure : Piece-wise linear

- ▶ Given f , can we compute $\hat{u} \in V$ as the best approximation to the actual solution u ?
- ▶ Idea 1: Given f , can we compute $\hat{u} \in V$ such that the laplacian of \hat{u} best approximates the function f ?
- ▶ Let error $e = \Delta \hat{u} - f$. We want $e \in V^\perp$.
 $\Rightarrow \langle \Delta \hat{u} - f, B_i \rangle = 0, \forall i$.

- Inner product on the mesh Ω : $\langle f, g \rangle_{\Omega} = \int_{\Omega} f(x)g(x) da$.
- ▶ Solve: $\langle \Delta \hat{u}, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega} \Rightarrow \sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- Given that \hat{u} (or B_j) $\in V$, how to define $\Delta \hat{u}$ (or ΔB_j).

- Inner product on the mesh Ω : $\langle f, g \rangle_{\Omega} = \int_{\Omega} f(x)g(x) da$.
- ▶ Solve: $\langle \Delta \hat{u}, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega} \Rightarrow \sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- Given that \hat{u} (or B_j) $\in V$, how to define $\Delta \hat{u}$ (or ΔB_j).

- Inner product on the mesh Ω : $\langle f, g \rangle_{\Omega} = \int_{\Omega} f(x)g(x) da$.
- ▶ Solve: $\langle \Delta \hat{u}, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega} \Rightarrow \sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- Given that \hat{u} (or B_j) $\in V$, how to define $\Delta \hat{u}$ (or ΔB_j).

- Integration by parts:

$$\int_{\Omega} f \Delta g da = \int_{\partial\Omega} f \langle \nabla g, n \rangle dl - \int_{\Omega} \langle \nabla f, \nabla g \rangle da,$$

where n is the outward unit normal to Ω on the boundary $\partial\Omega$.

- ▶ Note that ∇f and ∇g are vector fields on the mesh Ω , and the second term on the right hand side can be written as $\langle \nabla f, \nabla g \rangle_{\Omega}$.
- ▶ Idea 2: If Ω is a closed mesh, $\partial\Omega = \emptyset$, giving us

$$\langle f, \Delta g \rangle_{\Omega} = -\langle \nabla f, \nabla g \rangle_{\Omega}.$$

- ▶ Thus rather than solving $\sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$, we solve $-\sum_j u_j \langle \nabla B_j, \nabla B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- ▶ Let $u = (u_1, \dots, u_n)$, $\tilde{f} = (\langle f, B_1 \rangle_{\Omega}, \dots, \langle f, B_n \rangle_{\Omega})$, and $L_{ij} = \langle \nabla B_i, \nabla B_j \rangle_{\Omega}$. Using these notations, solve $-Lu = \tilde{f}$.

- Integration by parts:

$$\int_{\Omega} f \Delta g da = \int_{\partial\Omega} f \langle \nabla g, n \rangle dl - \int_{\Omega} \langle \nabla f, \nabla g \rangle da,$$

where n is the outward unit normal to Ω on the boundary $\partial\Omega$.

- ▶ Note that ∇f and ∇g are vector fields on the mesh Ω , and the second term on the right hand side can be written as $\langle \nabla f, \nabla g \rangle_{\Omega}$.
- ▶ Idea 2: If Ω is a closed mesh, $\partial\Omega = \emptyset$, giving us

$$\langle f, \Delta g \rangle_{\Omega} = -\langle \nabla f, \nabla g \rangle_{\Omega}.$$

- ▶ Thus rather than solving $\sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$, we solve $-\sum_j u_j \langle \nabla B_j, \nabla B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- ▶ Let $u = (u_1, \dots, u_n)$, $\tilde{f} = (\langle f, B_1 \rangle_{\Omega}, \dots, \langle f, B_n \rangle_{\Omega})$, and $L_{ij} = \langle \nabla B_i, \nabla B_j \rangle_{\Omega}$. Using these notations, solve $-Lu = \tilde{f}$.

- Integration by parts:

$$\int_{\Omega} f \Delta g da = \int_{\partial\Omega} f \langle \nabla g, n \rangle dl - \int_{\Omega} \langle \nabla f, \nabla g \rangle da,$$

where n is the outward unit normal to Ω on the boundary $\partial\Omega$.

- ▶ Note that ∇f and ∇g are vector fields on the mesh Ω , and the second term on the right hand side can be written as $\langle \nabla f, \nabla g \rangle_{\Omega}$.
- ▶ Idea 2: If Ω is a closed mesh, $\partial\Omega = \emptyset$, giving us

$$\langle f, \Delta g \rangle_{\Omega} = -\langle \nabla f, \nabla g \rangle_{\Omega}.$$

- ▶ Thus rather than solving $\sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$, we solve $-\sum_j u_j \langle \nabla B_j, \nabla B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- ▶ Let $u = (u_1, \dots, u_n)$, $\tilde{f} = (\langle f, B_1 \rangle_{\Omega}, \dots, \langle f, B_n \rangle_{\Omega})$, and $L_{ij} = \langle \nabla B_i, \nabla B_j \rangle_{\Omega}$. Using these notations, solve $-Lu = \tilde{f}$.

- Integration by parts:

$$\int_{\Omega} f \Delta g da = \int_{\partial\Omega} f \langle \nabla g, n \rangle dl - \int_{\Omega} \langle \nabla f, \nabla g \rangle da,$$

where n is the outward unit normal to Ω on the boundary $\partial\Omega$.

- ▶ Note that ∇f and ∇g are vector fields on the mesh Ω , and the second term on the right hand side can be written as $\langle \nabla f, \nabla g \rangle_{\Omega}$.
- ▶ Idea 2: If Ω is a closed mesh, $\partial\Omega = \emptyset$, giving us

$$\langle f, \Delta g \rangle_{\Omega} = -\langle \nabla f, \nabla g \rangle_{\Omega}.$$

- ▶ Thus rather than solving $\sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$, we solve $-\sum_j u_j \langle \nabla B_j, \nabla B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- ▶ Let $u = (u_1, \dots, u_n)$, $\tilde{f} = (\langle f, B_1 \rangle_{\Omega}, \dots, \langle f, B_n \rangle_{\Omega})$, and $L_{ij} = \langle \nabla B_i, \nabla B_j \rangle_{\Omega}$. Using these notations, solve $-Lu = \tilde{f}$.

- Integration by parts:

$$\int_{\Omega} f \Delta g da = \int_{\partial\Omega} f \langle \nabla g, n \rangle dl - \int_{\Omega} \langle \nabla f, \nabla g \rangle da,$$

where n is the outward unit normal to Ω on the boundary $\partial\Omega$.

- ▶ Note that ∇f and ∇g are vector fields on the mesh Ω , and the second term on the right hand side can be written as $\langle \nabla f, \nabla g \rangle_{\Omega}$.
- ▶ Idea 2: If Ω is a closed mesh, $\partial\Omega = \emptyset$, giving us

$$\langle f, \Delta g \rangle_{\Omega} = -\langle \nabla f, \nabla g \rangle_{\Omega}.$$

- ▶ Thus rather than solving $\sum_j u_j \langle \Delta B_j, B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$, we solve $-\sum_j u_j \langle \nabla B_j, \nabla B_i \rangle_{\Omega} = \langle f, B_i \rangle_{\Omega}$.
- ▶ Let $u = (u_1, \dots, u_n)$, $\tilde{f} = (\langle f, B_1 \rangle_{\Omega}, \dots, \langle f, B_n \rangle_{\Omega})$, and $L_{ij} = \langle \nabla B_i, \nabla B_j \rangle_{\Omega}$. Using these notations, solve $-Lu = \tilde{f}$.

- The discrete Laplacian of the function u on the mesh at vertex i is $(Lu)_i$.

- $\langle \nabla B_i, \nabla B_i \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_i \rangle_{\Delta_k}$.
- $\langle \nabla B_i, \nabla B_i \rangle_{\Delta_k} = \frac{1}{2}(\cot \alpha + \cot \beta)$.
- $\langle \nabla B_i, \nabla B_j \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_j \rangle_{\Delta_k}$
- $\langle \nabla B_i, \nabla B_j \rangle_{\Delta_k} = -\frac{1}{2} \cot \theta$.




Figure : Laplacian at a point in the mesh

- The discrete Laplacian of the function u on the mesh at vertex i is $(Lu)_i$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_i \rangle_{\Delta_k}$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Delta_k} = \frac{1}{2}(\cot \alpha + \cot \beta)$.
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_j \rangle_{\Delta_k}$
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Delta_k} = -\frac{1}{2} \cot \theta$.




Figure : Laplacian at a point in the mesh

- The discrete Laplacian of the function u on the mesh at vertex i is $(Lu)_i$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_i \rangle_{\Delta_k}$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Delta_k} = \frac{1}{2}(\cot \alpha + \cot \beta)$.
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_j \rangle_{\Delta_k}$
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Delta_k} = -\frac{1}{2} \cot \theta$.




Figure : Laplacian at a point in the mesh

- The discrete Laplacian of the function u on the mesh at vertex i is $(Lu)_i$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_i \rangle_{\Delta_k}$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Delta_k} = \frac{1}{2}(\cot \alpha + \cot \beta)$.
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_j \rangle_{\Delta_k}$
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Delta_k} = -\frac{1}{2} \cot \theta$.




Figure : Laplacian at a point in the mesh

- The discrete Laplacian of the function u on the mesh at vertex i is $(Lu)_i$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_i \rangle_{\Delta_k}$.
- ▶ $\langle \nabla B_i, \nabla B_i \rangle_{\Delta_k} = \frac{1}{2}(\cot \alpha + \cot \beta)$.
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Omega} = \sum_k \langle \nabla B_i, \nabla B_j \rangle_{\Delta_k}$
- ▶ $\langle \nabla B_i, \nabla B_j \rangle_{\Delta_k} = -\frac{1}{2} \cot \theta$.




Figure : Laplacian at a point in the mesh

- Thus $\langle \nabla B_i, \nabla B_i \rangle_\Omega = \frac{1}{2} \sum_{\Delta_i} (\cot \alpha_i + \cot \beta_i)$ and $\langle \nabla B_i, \nabla B_j \rangle_\Omega = -\frac{1}{2}(\cot \theta_1 + \cot \theta_2)$.
- The Discrete LBO is

$$L_{ii} = \frac{1}{2} \sum_{i \sim k} (\cot \alpha_k + \cot \beta_k)$$

$$L_{ij} = -\frac{1}{2}(\cot \alpha_j + \cot \beta_j), \quad \text{if } i \sim j.$$

$$L_{ij} = 0, \quad \text{otherwise.}$$

- ▶ Note that in this case $L = D - A$, you may want to use $\hat{L} = -L$.

- Thus $\langle \nabla B_i, \nabla B_i \rangle_\Omega = \frac{1}{2} \sum_{\Delta_i} (\cot \alpha_i + \cot \beta_i)$ and $\langle \nabla B_i, \nabla B_j \rangle_\Omega = -\frac{1}{2}(\cot \theta_1 + \cot \theta_2)$.
- The Discrete LBO is

$$L_{ii} = \frac{1}{2} \sum_{i \sim k} (\cot \alpha_k + \cot \beta_k)$$

$$L_{ij} = -\frac{1}{2}(\cot \alpha_j + \cot \beta_j), \quad \text{if } i \sim j.$$

$$L_{ij} = 0, \quad \text{otherwise.}$$

- Note that in this case $L = D - A$, you may want to use $\hat{L} = -L$.

- Thus $\langle \nabla B_i, \nabla B_i \rangle_\Omega = \frac{1}{2} \sum_{\Delta_i} (\cot \alpha_i + \cot \beta_i)$ and $\langle \nabla B_i, \nabla B_j \rangle_\Omega = -\frac{1}{2}(\cot \theta_1 + \cot \theta_2)$.
- The Discrete LBO is

$$L_{ii} = \frac{1}{2} \sum_{i \sim k} (\cot \alpha_k + \cot \beta_k)$$

$$L_{ij} = -\frac{1}{2}(\cot \alpha_j + \cot \beta_j), \quad \text{if } i \sim j.$$

$$L_{ij} = 0, \quad \text{otherwise.}$$

- Note that in this case $L = D - A$, you may want to use $\hat{L} = -L$.

Cotan Laplacian⁵

- Using the definition of LBO: $\nabla = \text{div} \circ \text{grad}$:

$$\int_{A_i} \Delta f(x) da = \int_{A_i} \text{div} \nabla f(x) da = \int_{\partial A_i} \nabla f(x) \cdot n(x) dl,$$

where A_i refers to a local integration (averaging) area, typically the Voronoi cell of a vertex.

- This reduces to

$$\int_{A_i} \Delta f(x) da = \frac{1}{2} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

- Thus the LBO on function f is:

$$\Delta f(v_i) = \frac{1}{2A_i} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

⁵Meyer et.al., Discrete Differential Geometry Operators for Triangulated 2-Manifolds, Vis. & Math - 3, 2003

Cotan Laplacian⁵

- Using the definition of LBO: $\nabla = \text{div} \circ \text{grad}$:

$$\int_{A_i} \Delta f(x) da = \int_{A_i} \text{div} \nabla f(x) da = \int_{\partial A_i} \nabla f(x) \cdot n(x) dl,$$

where A_i refers to a local integration (averaging) area, typically the Voronoi cell of a vertex.

- This reduces to

$$\int_{A_i} \Delta f(x) da = \frac{1}{2} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

- Thus the LBO on function f is:

$$\Delta f(v_i) = \frac{1}{2A_i} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

⁵Meyer et.al., Discrete Differential Geometry Operators for Triangulated 2-Manifolds, Vis. & Math - 3, 2003

Cotan Laplacian⁵

- Using the definition of LBO: $\nabla = \text{div} \circ \text{grad}$:

$$\int_{A_i} \Delta f(x) da = \int_{A_i} \text{div} \nabla f(x) da = \int_{\partial A_i} \nabla f(x) \cdot n(x) dl,$$

where A_i refers to a local integration (averaging) area, typically the Voronoi cell of a vertex.

- This reduces to

$$\int_{A_i} \Delta f(x) da = \frac{1}{2} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

- Thus the LBO on function f is:

$$\Delta f(v_i) = \frac{1}{2A_i} \sum_{j \in N(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(f_j - f_i)$$

⁵Meyer et.al., Discrete Differential Geometry Operators for Triangulated 2-Manifolds, Vis. & Math - 3, 2003

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?

- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?
- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?
- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?
- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?
- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

- Solution to Poisson eqn: $\Delta u = f$. What if $f \in V$?
- ▶ $\langle f, B_i \rangle_{\Omega} = \langle \sum_j f_j B_j, B_i \rangle_{\Omega} = \sum_j f_j \langle B_j, B_i \rangle_{\Omega}$.
- ▶ Let $A_{ij} = \int_{\Omega} B_i(x) B_j(x) da$. Then $\tilde{f} = A(f_1, \dots, f_n)^t = A\bar{f}$.
- ▶ The matrix A is called the *Mass matrix*, while the discrete Poisson eqn becomes $Lu = A\bar{f}$.
- ▶ The matrix $A^{-1}L$ is called the *Area normalized Cotan Laplacian*.
- ▶ One can derive the following:

$$A_{ii} = \frac{\text{Area of one-ring of vertex } i}{6}$$

$$A_{ij} = \frac{\text{Area of common triangles}}{12}$$

Application 1: Heat Equation

- $\frac{\partial u}{\partial t} = \Delta u.$




Figure : Heat diffusion on a cat.

Application 2: Compression




Figure : Mesh recovered using (top-left) 1, (top-right) 10, (bottom-left) 50 and (bottom-right) 200 coefficients, out of a total 3400 coefficients.

