

Computational Science M.S. Thesis Proposal

Production of massless spin-0 particles in analog (3+1)-dimensional de Sitter universes from a ground state BEC using high-performance computing

Nathaniel W. Chapman

Central Washington University

May 8, 2024

The story so far - Analog Universes

- ① Big Bang and rapid expansion $\implies$ could observe particles
- ② Big Bang was 14 billion years ago $\implies$ we can't observe particles
- ③ Condensed matter dynamics match particle dynamics (see Methods)
- ④ Can make condensed matter in the lab $\implies$ can effectively observe Big Bang particles
- ⑤ Lab based condensed matter system = “analog universe”

The story so far - Previous Research

Previous research has been done investigating a variety of different systems

- ① “Hyper-phononic” quasiparticles and Lorentz Violation
- ② Analog massive particles
- ③ Analog particles with non-zero quantum spin
- ④ Analog universes with different expansion behavior
- ⑤ Analog universes with non-zero background velocity
- ⑥ Experimental realizations (some extremely relevant and recent)

The story so far - 2D & Low-Performance Computing

- 1 Analog $(2+1)$ -dimensional universe with massless, scalar particles in a de Sitter expansion has been investigated
- 2 Previous investigation provides theory and guide for conceptual analysis
- 3 3d analog universe needs high-performance computing because of increased size of wave vector domain (see Methods)
- 4 Without 3d, 2d error is unknown

My story

My focus is on

- ① $(3+1)$ D analog universe with high-performance computing
- ② linear dispersion & phononic quasiparticle
- ③ analog massless particles
- ④ analog spin-0 particles
- ⑤ analog de Sitter expansion
- ⑥ no background velocity

Goals

- ① Provide insight into the dynamics of particles created due to the expansion of the universe in the moments just after the Big Bang.
- ② Quantitatively describe the error in using a reduced-dimensional physical model as compared to a full-dimensional model.
- ③ Quantitatively describe the efficiency of using a full-dimensional model with high-performance computing.
- ④ Provide an open-access, “ready to use”, packaged, high-performance computational model.

Methods Overview

① Physical Theory

- ① Setting up the Analog
- ② Speed of Sound & Universe Expansion
- ③ The Field Equation in Context
- ④ Particle Production
- ⑤ Computational Parameters

② Computation

- ① Computational Algorithms
- ② Programming Language

Methods - Physical Theory - Setting up the analog

- 1 Tangible: Ground state Bose-Einstein condensate with no thermal or quantum fluctuations
- 2 Mathematical: Gross-Pitaevskii equation (GPE) under a Bogoliubov mean-field approximation (aka the *nonlinear Schrödinger equation*),

$$i\hbar\partial_t\psi(\mathbf{x}, t) = \left[-\frac{\hbar^2}{2m}\nabla^2 + V_{\text{ext}}(x) + U|\psi(\mathbf{x}, t)|^2 \right] \psi(\mathbf{x}, t). \quad (1)$$

Methods - Physical Theory - Setting up the analog

- 1 Substitute the linearized Madelung density-phase representation (below) into eq. (1)

$$\psi(\mathbf{x}, t) = \sqrt{n_0 + \hat{n}(\mathbf{x}, t)} e^{i(\theta_0 + \hat{\theta}(\mathbf{x}, t))} \quad (2)$$

- 2 Yields the phase field equation in d spatial dimensions ($g := \det g_{\mu\nu}$)

The Field Equation

$$\frac{1}{\sqrt{-g}} \partial_\mu \left[\sqrt{-g} g^{\mu\nu} \partial_\nu \hat{\theta} \right] = 0 \quad g_{\mu\nu} = \left(\frac{n_0}{c(t)} \right)^{2/(d-1)} \begin{bmatrix} -c(t)^2 & \vdots & 0 \\ \dots & & \dots \\ 0 & \vdots & \delta_{ij} \end{bmatrix} \quad (3)$$

Methods - Physical Theory - Setting up the analog

The Analog

The field equation describes the dynamics of

- the phase of collective oscillations in the BEC
- particles produced due to the expansion of the universe

The Field Equation

$$\frac{1}{\sqrt{-g}} \partial_\mu [\sqrt{-g} g^{\mu\nu} \partial_\nu \hat{\theta}] = 0 \quad g_{\mu\nu} = \left(\frac{n_0}{c(t)} \right)^{2/(d-1)} \begin{bmatrix} -c(t)^2 & \vdots & 0 \\ \dots & & \dots \\ 0 & \vdots & \delta_{ij} \end{bmatrix}$$

Methods - Physical Theory - Speed of Sound & Universe Expansion

Speed Analog

Speed of sound in BEC = $c(t)$ = Speed of light in analog universe

- 1 Define $c(t)$ in the context of BEC

$$c(t)^2 = \frac{U(t)n_0}{m} = \frac{4\pi\hbar^2}{m^2}n_0a(t) \quad (4)$$

- 2 Scaling function $b(t)$

$$\overbrace{U(t) = U_0 b(t)}^{\text{BEC Interaction Strength}} \implies \overbrace{c(t) = c_0 \sqrt{b(t)}}^{\text{Analog Universe Light Speed}} \quad (5)$$

Methods - Physical Theory - Speed of Sound & Universe Expansion

- A decreasing speed of light is indistinguishable from an expanding universe (i.e. $c'(t) \propto 1/\text{expansion}$)

Expansion Analog

A decreasing BEC interaction strength yields an expanding analog universe

- Can tune interaction strength in the lab to induce a specific expansion
- $b(t)$ defines the expansion

Methods - Physical Theory - The Field Equation in Context

- 1 Explicitly expansion dependent phase field equation in (2+1)D (left) and conformal time (right)

$$\partial_\eta^2 \hat{\theta} - \frac{1}{2} \frac{b'(\eta)}{b(\eta)} \partial_\eta \hat{\theta} - c_0^2 \nabla^2 \hat{\theta} = 0 \quad d\eta = \sqrt{b(t)} dt \quad (6)$$

- 2 Field Fourier Expansions in wave vector $\mathbf{k}$

$$\hat{\theta}(\mathbf{x}, t) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{x}} \chi_{\mathbf{k}}(t) \hat{b}_{\mathbf{k}}(t) + e^{-i\mathbf{k} \cdot \mathbf{x}} \chi_{\mathbf{k}}^*(t) \hat{b}_{\mathbf{k}}^\dagger(t) \quad (7a)$$

$$\hat{n}(\mathbf{x}, t) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{x}} n_{\mathbf{k}}(t) \hat{b}_{\mathbf{k}}(t) + e^{-i\mathbf{k} \cdot \mathbf{x}} n_{\mathbf{k}}^*(t) \hat{b}_{\mathbf{k}}^\dagger(t) \quad (7b)$$

Methods - Physical Theory - The Field Equation in Context

- ③ Choose de Sitter expansion for early universe (t_s is the “rate of expansion”)

$$b(t) = e^{-t/t_s}, \quad 0 \leq t \leq t_f \quad (8)$$

- ④ Phase Field equation with de Sitter expansion

The Fourier de Sitter Field Equation

$$\partial_\eta^2 \chi_{\mathbf{k}} - \frac{1}{\eta} \partial_\eta \chi_{\mathbf{k}} + c_0^2 k^2 \chi_{\mathbf{k}} = 0 \quad (9)$$

- ⑤ Initial conditions from the continuity of $\chi_{\mathbf{k}}$ and $n_{\mathbf{k}} = -\frac{\hbar}{U} \partial_t \chi_{\mathbf{k}}$ at $t = 0$.

Methods - Physical Theory - Particle Production

- 1 Can represent phase and density amplitudes in same basis

$$u_k(t) = \frac{1}{2\sqrt{n_0}} n_k(t) + i\sqrt{n_0} \chi_k(t) \quad (10a)$$

$$v_k(t) = \frac{1}{2\sqrt{n_0}} n_k(t) - i\sqrt{n_0} \chi_k(t), \quad (10b)$$

- 2 Find $u_k(t)$ and $v_k(t)$ out of and during the expansion

Particle Production

$$N_k(t) = \left| u_k^{\text{out}*}(t) v_k^{\text{exp}*}(t) - v_k^{\text{out}*}(t) u_k^{\text{exp}*}(t) \right|^2 \quad (11)$$

Methods - Physical Theory - Particle Production Algorithm

- 1 Solve the Fourier de Sitter field equation (9) for $\chi_{\mathbf{k}}(t)$ at a point in space $\mathbf{x}$
- 2 Differentiate $\chi_{\mathbf{k}}(t)$ to get $n_{\mathbf{k}}(t)$
- 3 Combine $\chi_{\mathbf{k}}(t)$ and $n_{\mathbf{k}}(t)$ to get the mixed Fourier amplitudes as in equations (10)
- 4 Calculate the number of particles at position $\mathbf{x}$ at time t with wave vector $\mathbf{k}$ as in equation (11)

Methods - Physical Theory - Computational Parameters

	Previous	New
Universe Size $\mathbf{x}$	128×128	$128 \times 128 \times 128$
Time Scale t_s	10^{-5} to 10^{-3}	10^{-5} to 10^{-3}
Wave Vector Domain $\mathbf{k}$	$\ \mathbf{k}\ \leq 32 \times 2\pi$	$\ \mathbf{k}\ < k_c$

Table: Previous and new computational domains

- This system only valid for $\|\mathbf{k}\| < k_c$ such that $\hbar^2 k_c^2 / 2m = Un_0$
- Time step such that the total particle number for the entire field changed by much less than one, or that the change in norm was small enough i.e. $\Delta t : \Delta \|\psi(\mathbf{x}, t)\| \leq 10^{-9}$

Methods - Computational Methods - Algorithms & Hardware

Physics	Method
Fourier Field Equation	4-th order Runge-Kutta
Density-Phase Transformation	4-th order explicit central FD
Particle Production	vectorized arithmetic

Table: Computational methods

Use the “Turing” supercomputer for its

- “tight”-cluster configuration
- dedicated head, storage, and several compute nodes
- NVIDIA GPUs allowing for CUDA massive parallel computing

Methods - Computational Methods - Programming Language

- Will choose between Mathematica or Julia programming languages considering the following:

	Experience	Open-source	Free	Scientific	Parallel	Cluster	GPU
Mathematica	X			X	X	X	X
Julia		X	X	X	X	?	X

Table: Language comparison

- Other high-performance languages like C++ and Java could also be considered if performance is needed
- Mathematica is not good for reproducibility

Timeline Overview

- ① Performance Scaling Phase - Summer '23
- ② Domain Scaling Phase - Fall '23
- ③ Writing Phase - Winter '24
- ④ Dissemination Phase - Spring '24

Timeline - Performance Scaling Phase - Summer '23

- 1 Convert hybrid analytic-numerical algorithms to fully numeric low-performance algorithms (LPNA) in final language of choice
- 2 Test the LPNAs in the 2D model and compare against previous results
- 3 Translating the LPNAs to high-performance fully-numerical algorithms (HPNA) that make use of high-performance hardware
- 4 Test the HPNAs in the 2D model and compare against previous results, measuring the efficiency when compared to the LPNA

Timeline - Domain Scaling Phase - Fall '23

- ① Convert 2D theory to 3D
- ② If not generalized, convert the 2D HPNAs to 3D
- ③ Test the 3D HPNAs on a small spatial domain and small momentum domain
- ④ Test the 3D HPNAs on a small spatial domain and large momentum domain
- ⑤ Do the “real” run of the 3D HPNAs on a large spatial domain and large momentum domain
- ⑥ Revise and repeat runs until satisfactory results are achieved.

Timeline - Writing & Dissemination Phases - Winter & Spring '24

① Writing Phase - Winter '24

- ① Write report
- ② Prepare for conferences

② Dissemination Phase - Spring '24

- ① Defend thesis
- ② Present at conferences
- ③ Submit final document to be considered for journal publication and graduation

Anticipated Results

Refer back to goals

- Insight into particle production during the moments just after the Big Bang
- Quantitatively describe the error in lower-dimensional models
- Quantitatively describe the worth of using full-dimensional model with high-performance computing

Anticipated Results - Particle Production Insight

- 2D and 3D field equations are qualitatively equal

$$\text{2D: } \partial_t^2 \theta - \frac{3}{2} \frac{b'(t)}{b(t)} \partial_t \theta - c_0^2 b(t) \nabla^2 \theta = 0 \quad (12a)$$

$$\text{3D: } \partial_t^2 \theta - \frac{b'(t)}{b(t)} \partial_t \theta - c_0^2 b(t) \nabla^2 \theta = 0 \quad (12b)$$

- I expect qualitatively similar results from previous research

Anticipated Results - Low-dimensional Model Error

- Qualitatively similar results should yield small error
- Will determine how small
- Small enough to not warrant future research using HPC based 3D model?
- Big enough that more accurate results are sometimes warranted?

Anticipated Results - Efficiency

- Recreated analytic 2D results




Figure: The momentum distribution when the universe stops expanding for multiple effective rates of expansion

Anticipated Results - Efficiency




Figure: Momentum distribution throughout the expansion

Anticipated Results - Efficiency

- Used analytic solution for particle production
- Calculations still took a few seconds
- I expect the worth to be dependant on the domains of the parameters e.g. large spatial grid over a long time needs HPC but otherwise modern technology will yield resonable calculation times

Dissemination

- Journal-ready thesis submitted to Physical Review D
- Present at SOURCE
- Present at APS Northwest
- Possibly present at APS April Meeting

Thank you & Code source

Thank you for your attention.

The code for this project is available at:

- GitHub (QR code): github.com/nondairyneutrino/thesis
- My website: nathanwchapman.com



Bibliography I

- Anderson, M. H., J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell (1995). "Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor". In: *Science* 269.5221, pp. 198–201. DOI: [10.1126/science.269.5221.198](https://www.science.org/doi/pdf/10.1126/science.269.5221.198). eprint: <https://www.science.org/doi/pdf/10.1126/science.269.5221.198>.
- Barceló, C., S. Liberati, and M. Visser (Nov. 2003). "Probing semiclassical analog gravity in Bose-Einstein condensates with widely tunable interactions". In: *Phys. Rev. A* 68 (5), p. 053613. DOI: [10.1103/PhysRevA.68.053613](https://doi.org/10.1103/PhysRevA.68.053613).
- Barceló, C., S. Liberati, and M. Visser (Dec. 2005). "Analogue Gravity". In: *Living Reviews in Relativity* 8.1. DOI: [10.12942/lrr-2005-12](https://doi.org/10.12942/lrr-2005-12).

Bibliography II

- Birrell, N. D. and P. C. W. Davies (1982). *Quantum Fields in Curved Space*. Cambridge Monographs on Mathematical Physics. Cambridge University Press. DOI: [10.1017/CB09780511622632](https://doi.org/10.1017/CB09780511622632).
- Cornish, S. L., N. R. Claussen, J. L. Roberts, E. A. Cornell, and C. E. Wieman (Aug. 2000). "Stable ^{85}Rb Bose-Einstein Condensates with Widely Tunable Interactions". In: *Phys. Rev. Lett.* 85 (9), pp. 1795–1798. DOI: [10.1103/PhysRevLett.85.1795](https://doi.org/10.1103/PhysRevLett.85.1795).
- Fedichev, P. O. and U. R. Fischer (Dec. 2003). "Gibbons-Hawking Effect in the Sonic de Sitter Space-Time of an Expanding Bose-Einstein-Condensed Gas". In: *Phys. Rev. Lett.* 91 (24), p. 240407. DOI: [10.1103/PhysRevLett.91.240407](https://doi.org/10.1103/PhysRevLett.91.240407).

Bibliography III

- Fedichev, P. O. and U. R. Fischer (Mar. 2004a). ““Cosmological” quasiparticle production in harmonically trapped superfluid gases”. In: *Phys. Rev. A* 69 (3), p. 033602. DOI: [10.1103/PhysRevA.69.033602](https://doi.org/10.1103/PhysRevA.69.033602).
- Fedichev, P. O. and U. R. Fischer (Mar. 2004b). “Observer dependence for the phonon content of the sound field living on the effective curved space-time background of a Bose-Einstein condensate”. In: *Phys. Rev. D* 69 (6), p. 064021. DOI: [10.1103/PhysRevD.69.064021](https://doi.org/10.1103/PhysRevD.69.064021).
- Fulling, S. A. (1989). *Aspects of Quantum Field Theory in Curved Spacetime*. London Mathematical Society Student Texts. Cambridge University Press. DOI: [10.1017/CB09781139172073](https://doi.org/10.1017/CB09781139172073).

Bibliography IV

- Gardiner, C. W., J. R. Anglin, and T. I. A. Fudge (Mar. 2002). “The stochastic Gross-Pitaevskii equation”. In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 35.6, p. 1555. DOI: 10.1088/0953-4075/35/6/310.
- Gibbons, G. W. and S. W. Hawking (May 1977). “Cosmological event horizons, thermodynamics, and particle creation”. In: *Phys. Rev. D* 15 (10), pp. 2738–2751. DOI: 10.1103/PhysRevD.15.2738.
- Guth, A. H. (Jan. 1981). “Inflationary universe: A possible solution to the horizon and flatness problems”. In: *Phys. Rev. D* 23 (2), pp. 347–356. DOI: 10.1103/PhysRevD.23.347.
- Jacobson, T. (2004). *Introduction to Quantum Fields in Curved Spacetime and the Hawking Effect*. arXiv: gr-qc/0308048 [gr-qc].

Bibliography V

- Jain, P., S. Weinfurtner, M. Visser, and C. W. Gardiner (Sept. 2007). "Analog model of a Friedmann-Robertson-Walker universe in Bose-Einstein condensates: Application of the classical field method". In: *Phys. Rev. A* 76 (3), p. 033616. DOI: 10.1103/PhysRevA.76.033616.
- Lapedes, A. S. (Aug. 2008). "Bogoliubov transformations, propagators, and the Hawking effect". In: *Journal of Mathematical Physics* 49.8, pp. 082301–082307. ISSN: 0022-2488. DOI: 10.1063/1.2968077. eprint: <https://pubs.aip.org/aip/jmp/article-pdf/49/8/082301/12968077/jmp.49.8.082301.pdf>.
- Liberati, S., M. Visser, and S. Weinfurtner (Apr. 2006a). "Analogue quantum gravity phenomenology from a two-component Bose-Einstein condensate". In: *Classical and Quantum Gravity* 23.9, p. 3129. DOI: 10.1088/0264-9381/23/9/023.

Bibliography VI

- Liberati, S., M. Visser, and S. Weinfurtner (Apr. 2006b). “Naturalness in an Emergent Analogue Spacetime”. In: *Phys. Rev. Lett.* 96 (15), p. 151301. DOI: [10.1103/PhysRevLett.96.151301](https://doi.org/10.1103/PhysRevLett.96.151301).
- Roberts, J. L., N. R. Claussen, J. P. Burke, C. H. Greene, E. A. Cornell, and C. E. Wieman (Dec. 1998). “Resonant Magnetic Field Control of Elastic Scattering in Cold ^{85}Rb ”. In: *Phys. Rev. Lett.* 81 (23), pp. 5109–5112. DOI: [10.1103/PhysRevLett.81.5109](https://doi.org/10.1103/PhysRevLett.81.5109).
- Schützhold, R., G. Plunien, and G. Soff (Jan. 2002). “Dielectric Black Hole Analogs”. In: *Phys. Rev. Lett.* 88 (6), p. 061101. DOI: [10.1103/PhysRevLett.88.061101](https://doi.org/10.1103/PhysRevLett.88.061101).
- Steinhardt, P. J. and N. Turok (May 2002). “Cosmic evolution in a cyclic universe”. In: *Phys. Rev. D* 65 (12), p. 126003. DOI: [10.1103/PhysRevD.65.126003](https://doi.org/10.1103/PhysRevD.65.126003).

Bibliography VII

- Uhlmann, M., Y. Xu, and R. Schützhold (Nov. 2005). "Aspects of cosmic inflation in expanding Bose–Einstein condensates". In: *New Journal of Physics* 7.1, p. 248. DOI: [10.1088/1367-2630/7/1/248](https://doi.org/10.1088/1367-2630/7/1/248).
- Visser, M. (Jan. 1999). "Some general bounds for one-dimensional scattering". In: *Phys. Rev. A* 59 (1), pp. 427–438. DOI: [10.1103/PhysRevA.59.427](https://doi.org/10.1103/PhysRevA.59.427).
- VISSER, M., C. BARCELÓ, and S. LIBERATI (n.d.). "ACOUSTICS IN BOSE–EINSTEIN CONDENSATES AS AN EXAMPLE OF BROKEN LORENTZ SYMMETRY". In: *CPT and Lorentz Symmetry*, pp. 336–340. DOI: [10.1142/9789812778123_0043](https://doi.org/10.1142/9789812778123_0043). eprint: https://www.worldscientific.com/doi/pdf/10.1142/9789812778123_0043.

Bibliography VIII

Visser, M. and S. Weinfurtner (2004). *Massive phonon modes from a BEC-based analog model*. [arXiv: cond-mat/0409639](#) [[cond-mat.stat-mech](#)].

Visser, M. and S. Weinfurtner (Aug. 2005). “Massive Klein-Gordon equation from a Bose-Einstein-condensation-based analogue spacetime”. In: *Phys. Rev. D* 72 (4), p. 044020. DOI: [10.1103/PhysRevD.72.044020](#).

Vogels, J. M., C. C. Tsai, R. S. Freeland, S. J. J. M. F. Kokkelmans, B. J. Verhaar, and D. J. Heinzen (Aug. 1997). “Prediction of Feshbach resonances in collisions of ultracold rubidium atoms”. In: *Phys. Rev. A* 56 (2), R1067–R1070. DOI: [10.1103/PhysRevA.56.R1067](#).

Bibliography IX

- Weinfurtner, S., S. Liberati, and M. Visser (2007). “Analogue Space-time Based on 2-Component Bose-Einstein Condensates”. In: *Quantum Analogues: From Phase Transitions to Black Holes and Cosmology*. Ed. by W. G. Unruh and R. Schützhold. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 115–163. ISBN: 978-3-540-70859-9. DOI: 10.1007/3-540-70859-6_6.
- Weinfurtner, S. (Sept. 2005). “Analogue model for an expanding universe”. In: *General Relativity and Gravitation* 37.9, pp. 1549–1554. DOI: 10.1007/s10714-005-0135-7.
- Weinfurtner, S., S. Liberati, and M. Visser (May 2006a). “Analogue model for quantum gravity phenomenology”. In: *Journal of Physics A: Mathematical and General* 39.21, p. 6807. DOI: 10.1088/0305-4470/39/21/S83.

Bibliography X

Weinfurtner, S., S. Liberati, and M. Visser (Mar. 2006b). "Modelling Planck-scale Lorentz violation via analogue models". In: *Journal of Physics: Conference Series* 33.1, p. 373. DOI: 10.1088/1742-6596/33/1/046.