


胃肠道中的炎症小体：感染，癌症和肠道微生物群稳态

Inflammasomes in the gastrointestinal tract: infection, cancer and gut microbiota homeostasis 【1】



炎症小体复合物

炎症小体传感器 NLRP1b, NLRP3, NLRC4, AIM2 和 pyrin 都能够形成含有衔接蛋白ASC和半胱氨酸蛋白酶 caspase 1 的经典炎症小体复合物。NLRP1b 和 NLRC4 也在没有 ASC 的情况下募集 caspase 1, 因为它们的结构中存在 CARD结构域¹⁻³。NLRP3 和 NLRC4 的激活分别需要 NEK7 激酶和 NLR 家族成员神经元凋亡抑制蛋白 (NAIP)。Caspase 1 裂解前体细胞因子 pro-IL-1 β 和 pro-IL-18 以及成孔蛋白 gasdermin D。Gasdermin D 的活性片段寡聚化导致细胞膜上形成孔洞, 引发细胞焦亡(pyroptosis)⁴⁻¹¹。这些孔洞允许从细胞中被动释放具有生物活性的 IL-1 β 和 IL-18。非经典炎症小体的定义是需要人类 caspase 4 和 caspase 5 或小鼠 caspase 11 激活的 NLRP3 炎症小体复合物¹²。这些半胱天冬酶 (Caspase) 的活化导致 gasdermin D 的裂解和细胞焦亡⁴⁻⁶。gasdermin D 的成孔片段 (pore-forming fragment) 激活 NLRP3 炎症小体以及 IL-1 β 与 IL-18 的 caspase 1 依赖性成熟^{4,5}。

TIPs:

pyroptosis: 细胞焦亡,是一种新的程序性细胞死亡方式, 其特征为依赖于半胱天冬酶-1(caspase-1), 并伴有大量促炎症因子的释放。细胞焦亡的形态学特征、发生及调控机制等均不同于凋亡、坏死等其他细胞死亡方式¹³;

Caspase : 半胱天冬酶 ;

1. Poyet, J. L. et al. Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. *J. Biol. Chem.* 276, 28309–28313 (2001). 64
2. Van Opdenbosch, N. et al. Activation of the NLRP1b inflammasome independently of ASC-mediated caspase-1 autoproteolysis and speck formation. *Nat. Commun.* 5, 3209 (2014). 65
3. Guey, B., Bodnar, M., Manie, S. N., Tardivel, A. & Petrilli, V. Caspase-1 autoproteolysis is differentially required for NLRP1b and NLRP3 inflammasome function. *Proc. Natl Acad. Sci. USA* 111, 17254–17259 66
4. Kayagaki, N. et al. Caspase-11 cleaves gasdermin D for non-canonical inflammasome signaling. *Nature* 526, 666–671 (2015). 73
5. Shi, J. et al. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature* 526, 660–665 (2015). References 73 and 74 demonstrate that gasdermin D is a substrate of inflammatory caspases that upon cleavage converts to a pro-pyroptotic executor mediating cell lysis. 74
6. He, W. T. et al. Gasdermin D is an executor of pyroptosis and required for interleukin-1 β secretion. *Cell Res.* 25, 1285–1298 (2015). 75

7. Ding, J. et al. Pore-forming activity and structural autoinhibition of the gasdermin family. *Nature* 535, 111–116 (2016). 76
8. Aglietti, R. A. et al. GsdmD p30 elicited by caspase-11 during pyroptosis forms pores in membranes. *Proc. Natl Acad. Sci. USA* 113, 7858–7863 (2016). 77
9. Liu, X. et al. Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature* 535, 153–158 (2016). 78
10. Sborgi, L. et al. GSDMD membrane pore formation constitutes the mechanism of pyroptotic cell death. *EMBO J.* 35, 1766–1778 (2016). 79
11. Chen, X. et al. Pyroptosis is driven by non-selective gasdermin-D pore and its morphology is different from MLKL channel-mediated necroptosis. *Cell Res.* 26, 1007–1020 (2016). 80
12. Kayagaki, N. et al. Non-canonical inflammasome activation targets caspase-11. *Nature* 479, 117–121 (2011). This study describes the existence of a non-canonical 16
13. <http://news.bioon.com/article/6710531.html> 01