


胃肠道中的炎症小体：感染，癌症和肠道微生物群稳态

Inflammasomes in the gastrointestinal tract: infection, cancer and gut microbiota homeostasis [3]



炎症小体和其相关的分子在肠道细胞和免疫细胞中促进杀死和清除胃肠道病原菌

a | 炎症小体通过诱导分泌 IL-1 β , IL-18 和细胞焦亡介导保护宿主免受革兰氏阴性菌侵袭¹⁻⁸。NLRP6 和 NLRP12 可以负向调节炎症⁹⁻¹²。NLRP6 在杯状细胞中对 Toll 样受体诱导的自噬作出响应从而介导粘液的分泌¹³。RNA 结合的 DEAH 盒蛋白 9 (DHX9) 与 NLRP9b 相互作用, 诱导炎症小体复合物的组装¹⁴。NLRP6-DHX15 复合物结合病毒 RNA 并且诱导 1 型和 3 型干扰素的产生¹⁵。

b | 肠道巨噬细胞可以区分病原菌和共生菌¹⁶。

c | 激活 caspase 1, caspase 8 或者 caspase 11 导致细胞死亡, 进而促使从上皮清除和排出被感染的肠上皮细胞¹⁷⁻¹⁹。

d | 炎症小体可以通过抑制细菌摄入减少细菌负荷, 这会限制巨噬细胞的运动和刚度 (stiffness) 并且促进活性氧的产生 (ROS)²⁰。焦亡的巨噬细胞释放完整的细菌或者陷入细胞内陷阱的细菌²¹, 这些释放的实体会进一步被中性粒细胞吞噬^{3,22}。

TIPs:

pyroptosis: 细胞焦亡,是一种新的程序性细胞死亡方式,其特征为依赖于半胱天冬酶-1(caspase-1),并伴有大量促炎症因子的释放。细胞焦亡的形态学特征、发生及调控机制等均不同于凋亡、坏死等其他细胞死亡方式⁰¹;

Caspase : 半胱天冬酶;

ASC: apoptosis-associated speck-like protein containing a caspase activation and recruitment domain (CARD) : 含有半胱天冬酶激活和募集结构域的凋亡相关斑点样蛋白
dsRNA , double-stranded RNA: 双链 RNA;

EMCV, encephalomyocarditis virus: 脑心肌炎病毒;

KC, keratinocyte chemoattractant: 角质形成细胞化学引诱物 (也称为CXCL1) ;

MAVS, mitochondrial antiviral-signalling protein: 线粒体抗病毒信号蛋白;

NAIP, neuronal apoptosis inhibitory protein: 神经细胞凋亡抑制蛋白;

NF- κ B, nuclear factor- κ B: 核因子- κ B;

NLRC4, nucleotide-binding domain, leucine-rich repeat-containing protein (NLR) family CARD domain-containing protein 4 : 核苷酸结合结构域, 富含亮氨酸重复序列的蛋白 (NLR) 家族 CARD 结构域蛋白4 (NLRC4)

ssRNA, single-stranded RNA : 单链RNA;

1. Broz, P. et al. Redundant roles for inflammasome receptors NLRP3 and NLRC4 in host defense against *Salmonella*. *J. Exp. Med.* 207, 1745–1755 (2010). 55
2. Franklin, B. S. et al. The adaptor ASC has extracellular and ‘prionoid’ activities that propagate inflammation. *Nat. Immunol.* 15, 727–737 (2014). 59
3. Miao, E. A. et al. Caspase-1-induced pyroptosis is an innate immune effector mechanism against intracellular bacteria. *Nat. Immunol.* 11, 1136–1142 (2010). 84
4. Franchi, L. et al. NLRC4-driven production of IL-1 β discriminates between pathogenic and commensal bacteria and promotes host intestinal defense. *Nat. Immunol.* 13, 449–456 (2012). 85
5. Liu, Z. et al. Role of inflammasomes in host defense against *Citrobacter rodentium* infection. *J. Biol. Chem.* 287, 16955–16964 (2012). 91
6. Nordlander, S., Pott, J. & Maloy, K. J. NLRC4 expression in intestinal epithelial cells mediates protection against an enteric pathogen. *Mucosal Immunol.* 7, 775–785 (2014). 92
7. Lara-Tejero, M. et al. Role of the caspase-1 inflammasome in *Salmonella typhimurium* pathogenesis. *J. Exp. Med.* 203, 1407–1412 (2006). 254
8. Song-Zhao, G. X. et al. Nlrp3 activation in the intestinal epithelium protects against a mucosal pathogen. *Mucosal Immunol.* 7, 763–774 (2014). 255
9. Anand, P. K. et al. NLRP6 negatively regulates innate immunity and host defence against bacterial pathogens. *Nature* 488, 389–393 (2012). 133
10. Zaki, M. H., Man, S. M., Vogel, P., Lamkanfi, M. & Kanneganti, T. D. *Salmonella* exploits NLRP12- dependent innate immune signaling to suppress host defenses during infection. *Proc. Natl Acad. Sci. USA* 111, 385–390 (2014). 134
11. Allen, I. C. et al. NLRP12 suppresses colon inflammation and tumorigenesis through the negative regulation of noncanonical NF- κ B signaling. *Immunity* 36, 742–754 (2012) 256
12. Zaki, M. H. et al. The NOD-like receptor NLRP12 attenuates colon inflammation and tumorigenesis. *Cancer Cell* 20, 649–660 (2011). References 256 and 257 reveal an important role for NLRP12 in colitis and colitis-associated colorectal cancer in mice. 257
13. Wlodarska, M. et al. NLRP6 inflammasome orchestrates the colonic host-microbial interface by regulating goblet cell mucus secretion. *Cell* 156, 1045–1059 (2014). 137
14. Zhu, S. et al. Nlrp9b inflammasome restricts rotavirus infection in intestinal epithelial cells. *Nature* 546, 667–670 (2017). This study reports the existence of an NLRP9b inflammasome complex, which is expressed specifically in mouse intestinal cells and mediates cytosolic recognition of rotavirus. 161
15. Wang, P. et al. Nlrp6 regulates intestinal antiviral innate immunity. *Science* 350, 826–830 (2015). 165

16. Franchi, L. et al. NLRC4-driven production of IL-1 β discriminates between pathogenic and commensal bacteria and promotes host intestinal defense. *Nat. Immunol.* 13, 449–456 (2012). 85
17. Sellin, M. E. et al. Epithelium-Intrinsic NAIP/NLRC4 inflammasome drives infected enterocyte expulsion to restrict *Salmonella* replication in the intestinal mucosa. *Cell Host Microbe* 16, 237–248 (2014) 93
18. Rauch, I. et al. NAIP-NLRC4 inflammasomes coordinate intestinal epithelial cell expulsion with eicosanoid and IL-18 release via activation of caspase-1 and -8. *Immunity* 46, 649–659 (2017). 94
19. Knodler, L. A. et al. Noncanonical inflammasome activation of caspase-4/caspase-11 mediates epithelial defenses against enteric bacterial pathogens. *Cell Host Microbe* 16, 249–256 (2014). References 93, 94 and 104 highlight an expulsion mechanism induced by the inflammasome to remove an entire infected host cell from the intestinal epithelium. 104
20. Man, S. M. et al. Actin polymerization as a key innate immune effector mechanism to control *Salmonella* infection. *Proc. Natl Acad. Sci. USA* 111, 17588–17593 (2014). 105
21. Jorgensen, I., Zhang, Y., Krantz, B. A. & Miao, E. A. Pyroptosis triggers pore-induced intracellular traps (PITs) that capture bacteria and lead to their clearance by efferocytosis. *J. Exp. Med.* 213, 2113–2128 (2016). 103
22. Chen, K. W. et al. The neutrophil NLRC4 inflammasome selectively promotes IL-1 β maturation without pyroptosis during acute *Salmonella* challenge. *Cell Rep.* 8, 570–582 (2014). 106