

Model Selection and Model Averaging

Michael Noonan

February 7, 2021

Biol 520C: Statistical modelling for biological data

1. Housekeeping
2. The ΔAIC Grey Zone
3. The $\Delta AIC = 2$ Threshold
4. AIC Overfitting
5. Model Averaging
6. Model Selection and Averaging Recap

Housekeeping

- Rubric and time slots for the first talk are posted on Canvas.

- Rubric and time slots for the first talk are posted on Canvas.
- If you're still struggling with R have a look at Prof. Schluter's R tips page for a similar course run on the Vancouver campus.
<https://www.zoology.ubc.ca/schluter/R/>

The Δ AIC Grey Zone



Last lecture we covered two approaches for model selection:

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

2. **Information criteria**

IC don't require nested models

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

2. **Information criteria**

IC don't require nested models, well grounded theoretically

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

2. **Information criteria**

IC don't require nested models, well grounded theoretically, easy to apply in practice.

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

2. **Information criteria**

IC don't require nested models, well grounded theoretically, easy to apply in practice.

Just because a model is the best fit out of a pool of candidates doesn't mean it's any good.

Last lecture we covered two approaches for model selection:

1. **Likelihood-ratio tests**

LRTs are quick easy, well grounded theoretically, result in clear outcomes, but become unwieldy for complex models.

If some of the models aren't nested in one another, there is no way to compare them using LRTs.

2. **Information criteria**

IC don't require nested models, well grounded theoretically, easy to apply in practice.

Just because a model is the best fit out of a pool of candidates doesn't mean it's any good.

Sometimes IC approaches result in a clear winner, other times...

Clicker question: i



THE UNIVERSITY OF BRITISH COLUMBIA
Okanagan Campus

```
FIT <- gls(num_sp ~ latitude + elevation, data = data, method = "ML")  
  
dredge(FIT)  
  
Global model call: gls(model = num_sp ~ latitude + elevation, data = data, method = "ML")  
---  
Model selection table  
      (Intrc)      elvtn      lattd df  logLik AICc delta weight  
4  11.120 -0.001373 -0.2018  4   -4.059 18.5   0.00  0.922  
2    2.489 -0.001613          3   -8.672 24.7   6.21  0.041  
3   12.390          -0.2388  3   -8.857 25.0   6.58  0.034  
1    2.114          2  -13.224 31.1  12.61  0.002  
Models ranked by AICc(x)
```

Which parameters would you include in your model?

- A** — latitude and elevation.
- B** — Only elevation.
- C** — Only latitude.
- D** — Only the intercept.

```
FIT <- gls(num_sp ~ latitude + elevation, data = data, method = "ML")  
  
dredge(FIT)  
  
Global model call: gls(model = num_sp ~ latitude + elevation, data = data, method = "ML")  
---  
Model selection table  
      (Intrc)      elvtn    lattd df  logLik AICc  delta weight  
4   11.120 -0.001373 -0.2018  4   -4.059 18.5   0.00  0.922  
2    2.489 -0.001613          3   -8.672 24.7   6.21  0.041  
3   12.390          -0.2388  3   -8.857 25.0   6.58  0.034  
1    2.114          2   -13.224 31.1  12.61  0.002  
Models ranked by AICc(x)
```

Which parameters would you include in your model?

A — latitude and elevation.

C — Only latitude.

B — Only elevation.

D — Only the intercept.

Clicker question: ii



THE UNIVERSITY OF BRITISH COLUMBIA
Okanagan Campus

Model selection table											
	(Int)	β_1	β_2	β_3	β_4	β_5	β_6	df	logLik	AICc	delta weight
33	-103.000					0.05326		4	-10370.46	20748.9	0.00 0.202
49	-103.000					+ 0.05326		4	-10370.46	20748.9	0.00 0.202
34	-104.200	0.25270				0.05385		5	-10369.64	20749.3	0.36 0.168
50	-104.200	0.25270				+ 0.05385		5	-10369.64	20749.3	0.36 0.168
1	4.208							3	-10372.99	20752.0	3.07 0.043
17	4.208					+		3	-10372.99	20752.0	3.07 0.043
2	4.212	-0.05585						4	-10372.23	20752.5	3.55 0.034
18	4.212	-0.05585				+		4	-10372.23	20752.5	3.55 0.034
37	-51.480			-0.028750		0.05640		5	-10372.58	20755.2	6.24 0.009

Which parameters would you include in your model?

A — β_1 , β_6 , and β_5

C — β_6 and β_5

B — Only β_6

D — A, B, and C are good options.

Model selection table											
	(Int)	β_1	β_2	β_3	β_4	β_5	β_6	df	logLik	AICc	delta weight
33	-103.000					0.05326		4	-10370.46	20748.9	0.00 0.202
49	-103.000					+ 0.05326		4	-10370.46	20748.9	0.00 0.202
34	-104.200	0.25270				0.05385		5	-10369.64	20749.3	0.36 0.168
50	-104.200	0.25270				+ 0.05385		5	-10369.64	20749.3	0.36 0.168
1	4.208							3	-10372.99	20752.0	3.07 0.043
17	4.208					+		3	-10372.99	20752.0	3.07 0.043
2	4.212	-0.05585						4	-10372.23	20752.5	3.55 0.034
18	4.212	-0.05585				+		4	-10372.23	20752.5	3.55 0.034
37	-51.480			-0.028750		0.05640		5	-10372.58	20755.2	6.24 0.009

Which parameters would you include in your model?

A — β_1 , β_6 , and β_5

C — β_6 and β_5

B — Only β_6

D — A, B, and C are good options.

No clear winner?



THE UNIVERSITY OF BRITISH COLUMBIA
Okanagan Campus

Sometimes ΔAICc based model selection leads to a clear 'winner'

Sometimes ΔAICc based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

Sometimes ΔAICc based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

As a modeler, you need to use this information to make some decision.
But what do you do?

Sometimes $\Delta AICc$ based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

As a modeler, you need to use this information to make some decision.
But what do you do?

As good scientists, we turn to the literature for an answer:

Sometimes ΔAIC_c based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

As a modeler, you need to use this information to make some decision.
But what do you do?

As good scientists, we turn to the literature for an answer:

Burnham & Anderson (2002)

1. ΔAIC 0–2: Substantial supp.
2. ΔAIC 4–7: Considerably less supp.
3. $\Delta AIC > 10$: Essentially no supp.

Sometimes ΔAIC_c based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

As a modeler, you need to use this information to make some decision.
But what do you do?

As good scientists, we turn to the literature for an answer:

Burnham & Anderson (2002)

1. ΔAIC 0–2: Substantial supp.
2. ΔAIC 4–7: Considerably less supp.
3. $\Delta AIC >10$: Essentially no supp.

Burnham *et al.* (2011)

1. ΔAIC 0–7: Plausible
2. ΔAIC 7–14: Equivocal
3. $\Delta AIC >14$: Implausible

Sometimes ΔAIC_c based model selection leads to a clear 'winner', other times the differences between the top models are miniscule.

As a modeler, you need to use this information to make some decision.
But what do you do?

As good scientists, we turn to the literature for an answer:

Burnham & Anderson (2002)

1. ΔAIC 0–2: Substantial supp.
2. ΔAIC 4–7: Considerably less supp.
3. $\Delta AIC >10$: Essentially no supp.

Burnham *et al.* (2011)

1. ΔAIC 0–7: Plausible
2. ΔAIC 7–14: Equivocal
3. $\Delta AIC >14$: Implausible

Not very helpful...



Because there are no p -values or clear cutoffs tied to ΔAIC values, the literature is confused on what we should do.

Because there are no p -values or clear cutoffs tied to ΔAIC values, the literature is confused on what we should do.

One recurring approach is to use $\Delta AIC < 2$ as a cut-off

Because there are no p -values or clear cutoffs tied to ΔAIC values, the literature is confused on what we should do.

One recurring approach is to use $\Delta AIC < 2$ as a cut-off, which comes from the recommendations of Burnham & Anderson (2002).

Because there are no p -values or clear cutoffs tied to ΔAIC values, the literature is confused on what we should do.

One recurring approach is to use $\Delta AIC < 2$ as a cut-off, which comes from the recommendations of Burnham & Anderson (2002).


Lower AIC values are better (all else being equal), but what's so special about $\Delta AIC < 2$?

The $\Delta\text{AIC} = 2$ Threshold





The ΔAIC values for any given model are linked to the evidence ratio for the best model as $e^{-\left(\frac{\Delta AIC}{2}\right)}$.

The ΔAIC values for any given model are linked to the evidence ratio for the best model as $e^{-\left(\frac{\Delta AIC}{2}\right)}$.



The ΔAIC values for any given model are linked to the evidence ratio for the best model as $e^{-\left(\frac{\Delta AIC}{2}\right)}$.



With $\Delta AIC \lesssim 1.38$ the evidence < 2 , with $\Delta AIC \lesssim 2$, evidence < 3 .

So in the $\Delta AIC < 2$ regime models are only \sim twice as likely as the AIC 'best' model.



The evidence ratio gives us a feel for why this range of values might be important, but it's still a bit murky.

The evidence ratio gives us a feel for why this range of values might be important, but it's still a bit murky.

The equation for AIC is:

$$AIC = -2 \ln(\mathcal{L}) + 2K$$

The evidence ratio gives us a feel for why this range of values might be important, but it's still a bit murky.

The equation for AIC is:

$$AIC = -2 \ln(\mathcal{L}) + 2K$$

The equation for ΔAIC is:

$$\Delta AIC = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

The evidence ratio gives us a feel for why this range of values might be important, but it's still a bit murky.

The equation for AIC is:

$$AIC = -2 \ln(\mathcal{L}) + 2K$$

The equation for ΔAIC is:

$$\Delta AIC = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

What happens if two models have the \sim the same likelihood and only differ by one parameter?



If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

$$\Delta\text{AIC} = \cancel{-2 \log \mathcal{L}(\theta_1)} + \cancel{2 \log \mathcal{L}(\theta_1)}$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

$$\Delta\text{AIC} = \cancel{-2 \log \mathcal{L}(\theta_1)} + \cancel{2 \log \mathcal{L}(\theta_1)} - 2(\cancel{K_2} - \cancel{K_2} - 1)$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

$$\Delta\text{AIC} = \cancel{-2 \log \mathcal{L}(\theta_1)} + \cancel{2 \log \mathcal{L}(\theta_1)} - 2(\cancel{K_2} - \cancel{K_2} - 1)$$

$$\Delta\text{AIC} = -2(-1)$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

$$\Delta\text{AIC} = \cancel{-2 \log \mathcal{L}(\theta_1)} + \cancel{2 \log \mathcal{L}(\theta_1)} - 2(\cancel{K_2} - \cancel{K_2} - 1)$$

$$\Delta\text{AIC} = -2(-1) = 2$$

If $\mathcal{L}(\theta_1) \approx \mathcal{L}(\theta_2)$ and $K_1 = K_2 + 1$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_2) - 2(K_2 - K_1)$$

$$\Delta\text{AIC} = -2 \log \mathcal{L}(\theta_1) + 2 \log \mathcal{L}(\theta_1) - 2(K_2 - (K_2 + 1))$$

$$\Delta\text{AIC} = \cancel{-2 \log \mathcal{L}(\theta_1)} + \cancel{2 \log \mathcal{L}(\theta_1)} - 2(\cancel{K_2} - \cancel{K_2} - 1)$$

$$\Delta\text{AIC} = -2(-1) = 2$$

Because the penalty term is $2K$, models with the \sim the same likelihood that only differ by one parameter will, by definition, have a ΔAIC of 2.

So this sheds some more light on the ΔAIC of 2 threshold

So this sheds some more light on the ΔAIC of 2 threshold and explains why in practice you'll find models that differ from the 'best fit' model by only one parameter within the ΔAIC of 2 threshold.

So this sheds some more light on the ΔAIC of 2 threshold and explains why in practice you'll find models that differ from the 'best fit' model by only one parameter within the ΔAIC of 2 threshold.

Var 1	Var 2	Var 3	Var 4	Var 5	AIC	ΔAIC	Weight
	-0.539 (± 0.244)		-0.602 (± 0.190)		1789.73	0.00	0.26
	-0.674 (± 0.336)		-0.609 (± 0.192)	0.173 (± 0.295)	1791.38	1.65	0.12
	-0.544 (± 0.245)	0.003 (± 0.008)	-0.566 (± 0.214)		1791.60	1.86	0.10
-0.090 (± 0.333)	-0.541 (± 0.244)		-0.574 (± 0.217)		1791.66	1.93	0.10
			-0.641 (± 0.201)		1792.23	2.50	0.08
-0.070 (± 0.335)	-0.670 (± 0.336)		-0.586 (± 0.220)	0.167 (± 0.296)	1793.34	3.61	0.04
			-0.622 (± 0.198)	-0.212 (± 0.222)	1793.34	3.61	0.04
	-0.662 (± 0.344)	0.001 (± 0.008)	-0.591 (± 0.591)	0.155 (± 0.316)	1793.35	3.62	0.04

Source: Mark Brewer

Second best model is $\mathcal{M}_1 + \text{Var}_5$, third best model is $\mathcal{M}_1 + \text{Var}_3$, fourth best is $\mathcal{M}_1 + \text{Var}_1$, all within the ΔAIC of 2 threshold.

ΔAIC of 2 and p -values



THE UNIVERSITY OF BRITISH COLUMBIA
Okanagan Campus

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

$$\Delta\text{AIC} = \lambda - 2(K_2 - K_1)$$

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

$$\Delta\text{AIC} = \lambda - 2(K_2 - K_1)$$


$$P = \Pr(\chi_k^2 > \Delta\text{AIC} + 2(K_2 - K_1))$$

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

$$\Delta\text{AIC} = \lambda - 2(K_2 - K_1)$$

$$P = \Pr(\chi_k^2 > \Delta\text{AIC} + 2(K_2 - K_1))$$




(Murtaugh, 2014)

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

$$\Delta\text{AIC} = \lambda - 2(K_2 - K_1)$$

$$P = \Pr(\chi_k^2 > \Delta\text{AIC} + 2(K_2 - K_1))$$



(Murtaugh, 2014)


There's a one-to-one relationship between ΔAIC and p -values e.g., a ΔAIC of 2 when models differ by 1 parameter corresponds to a p -value of ~ 0.047

The likelihood ratio test allows for estimating p -values for the support between a pair of models.

$$P = \Pr(\chi_k^2 > \lambda)$$

$$\Delta\text{AIC} = \lambda - 2(K_2 - K_1)$$

$$P = \Pr(\chi_k^2 > \Delta\text{AIC} + 2(K_2 - K_1))$$



(Murtaugh, 2014)

There's a one-to-one relationship between ΔAIC and p -values e.g., a ΔAIC of 2 when models differ by 1 parameter corresponds to a p -value of ~ 0.047 (but a p -value of ~ 0.015 when they differ by 10 parameters).

ΔAIC of 2 and p -values





THE UNIVERSITY OF BRITISH COLUMBIA
Okanagan Campus

Putting the pieces together:

Putting the pieces together:

$\Delta AIC = 2$ corresponds to an evidence ratio of ~ 2.7





(Murtaugh, 2014)

Putting the pieces together:

$\Delta AIC = 2$ corresponds to an evidence ratio of ~ 2.7

When two models have identical likelihoods, they can not have $\Delta AIC < 2$ (caps our capacity to distinguish models).





(Murtaugh, 2014)

Putting the pieces together:

$\Delta AIC = 2$ corresponds to an evidence ratio of ~ 2.7

When two models have identical likelihoods, they can not have $\Delta AIC < 2$ (caps our capacity to distinguish models).

A ΔAIC of 2 with a differences of 1 parameter corresponds to a p -value of ~ 0.047 , meaning the complexity is a significant improvement.



(Murtaugh, 2014)



Now we we have a better idea of why $\Delta AIC < 2$ is pervasive in the literature.

Now we we have a better idea of why $\Delta AIC < 2$ is pervasive in the literature.

But are we any closer to knowing what to do when we have a number of top contenders (i.e., $\Delta AIC < 2$)?

AIC Overfitting

Before we decide what to do when multiple models are reasonable contenders we need to explore how AIC model selection performs in practice.

Before we decide what to do when multiple models are reasonable contenders we need to explore how AIC model selection performs in practice.


It's well known in the statistics literature that AIC has a tendency to overfit

Before we decide what to do when multiple models are reasonable contenders we need to explore how AIC model selection performs in practice.


It's well known in the statistics literature that AIC has a tendency to overfit, but what does that look like?




Strong Effects



Tapered Effects




Strong Effects




Randomly used 5 of them


Tapered Effects



Strong Effects




Tapered Effects




Randomly used 5 of them, and added 5 other 'noise' parameters.

Strong Effects




Tapered Effects




Randomly used 5 of them, and added 5 other 'noise' parameters.

Fit models

Strong Effects




Tapered Effects




Randomly used 5 of them, and added 5 other 'noise' parameters.

Fit models, selected the best by AIC and AICc

Strong Effects




Tapered Effects




Randomly used 5 of them, and added 5 other 'noise' parameters.

Fit models, selected the best by AIC and AICc, compared which parameters were selected based on the true system

Strong Effects





Tapered Effects





Randomly used 5 of them, and added 5 other 'noise' parameters.

Fit models, selected the best by AIC and AICc, compared which parameters were selected based on the true system, repeated this 1000s of times









Tapered Effects Results



Strong Effects





Tapered Effects





For systems with strong effect sizes, both AIC and AICc identified true parameters well for high n .

Strong Effects







Tapered Effects







For systems with strong effect sizes, both AIC and AICc identified true parameters well for high n .

For systems with tapered effect sizes, neither AIC nor AICc identified all of the true parameters well



For systems with strong effect sizes, both AIC and AICc identified true parameters well for high n .





For systems with tapered effect sizes, neither AIC nor AICc identified all of the true parameters well even with n was extremely high.



For systems with strong effect sizes, both AIC and AICc identified true parameters well for high n .

For systems with tapered effect sizes, neither AIC nor AICc identified all of the true parameters well even with n was extremely high.

When n was small, AIC missed fewer true parameters than AICc, but at the cost of more false positives.




For systems with strong effect sizes, both AIC and AICc identified true parameters well for high n .

For systems with tapered effect sizes, neither AIC nor AICc identified all of the true parameters well even with n was extremely high.

When n was small, AIC missed fewer true parameters than AICc, but at the cost of more false positives.

For both systems AIC and AICc consistently identified noise parameters as being important.





In practice, AIC/AICc tend to overfit (pick too many parameters)

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC/AICc}$ values are small our ability to distinguish between models gets murky.

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC/AICc}$ values are small our ability to distinguish between models gets murky.

This leaves us with a number of different options:

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC}/\text{AICc}$ values are small our ability to distinguish between models gets murky.

This leaves us with a number of different options:

1. Be cautious and pick the most parsimonious model.

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC}/\text{AICc}$ values are small our ability to distinguish between models gets murky.

This leaves us with a number of different options:

1. Be cautious and pick the most parsimonious model.
2. Be liberal and include all contending parameters in the final model.

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC}/\text{AICc}$ values are small our ability to distinguish between models gets murky.

This leaves us with a number of different options:

1. Be cautious and pick the most parsimonious model.
2. Be liberal and include all contending parameters in the final model.
3. Conduct likelihood-ratio tests on the top models.

In practice, AIC/AICc tend to overfit (pick too many parameters), but when effect sizes are small they can also miss key parameters.

When $\Delta\text{AIC}/\text{AICc}$ values are small our ability to distinguish between models gets murky.

This leaves us with a number of different options:

1. Be cautious and pick the most parsimonious model.
2. Be liberal and include all contending parameters in the final model.
3. Conduct likelihood-ratio tests on the top models.
4. Perform model averaging.


Model Averaging



Model averaging refers to the practice of using several models at once for making predictions or inferring parameters.

Model averaging refers to the practice of using several models at once for making predictions or inferring parameters.

The Idea: If a model is misspecified, the parameters estimates may be too high/low (e.g., noise parameters soaking up effects or some effects getting inflated to make up for missing parameters).




(Dormann *et al.*, 2018)


Model averaging refers to the practice of using several models at once for making predictions or inferring parameters.

The Idea: If a model is misspecified, the parameters estimates may be too high/low (e.g., noise parameters soaking up effects or some effects getting inflated to make up for missing parameters).

Averaging parameter values from different models, with biases in either way, should cancel out and reduce bias in the average.




(Dormann *et al.*, 2018)



(Dormann *et al.*, 2018)


Variance between models is also important!
(Dormann *et al.*, 2018)



(Dormann *et al.*, 2018)

Variance between models is also important!
(Dormann *et al.*, 2018)

If all models make identical, and wrong predictions,
this would cancel any benefit of averaging.




(Dormann *et al.*, 2018)

Variance between models is also important!
(Dormann *et al.*, 2018)

If all models make identical, and wrong predictions, this would cancel any benefit of averaging.

If, however, model predictions are incorrect but vary evenly around the truth, there are substantial benefits to averaging.




(Dormann *et al.*, 2018)

Variance between models is also important!
(Dormann *et al.*, 2018)

If all models make identical, and wrong predictions, this would cancel any benefit of averaging.

If, however, model predictions are incorrect but vary evenly around the truth, there are substantial benefits to averaging.

In other words, you only then get the full benefits of model averaging when models have very different parameter estimates.



(Dormann *et al.*, 2018)



Challenge: If we throw all models we can possibly think of into an averaging procedure, but only a few are actually reasonable, then the junk can ruin the averaged model.

Challenge: If we throw all models we can possibly think of into an averaging procedure, but only a few are actually reasonable, then the junk can ruin the averaged model.

One solution you see in many papers is to only average plausible models (e.g., only models within ΔAICc of 2).

Challenge: If we throw all models we can possibly think of into an averaging procedure, but only a few are actually reasonable, then the junk can ruin the averaged model.

One solution you see in many papers is to only average plausible models (e.g., only models within ΔAICc of 2).

Another solution is to somehow weight all of the different models based on their 'plausibility'.

Challenge: If we throw all models we can possibly think of into an averaging procedure, but only a few are actually reasonable, then the junk can ruin the averaged model.

One solution you see in many papers is to only average plausible models (e.g., only models within ΔAICc of 2).

Another solution is to somehow weight all of the different models based on their 'plausibility'.

A third solution is to take a hybrid approach of averaging a reduced pool of unequally weighted models.

Challenge: If we throw all models we can possibly think of into an averaging procedure, but only a few are actually reasonable, then the junk can ruin the averaged model.

One solution you see in many papers is to only average plausible models (e.g., only models within ΔAICc of 2).

Another solution is to somehow weight all of the different models based on their 'plausibility'.

A third solution is to take a hybrid approach of averaging a reduced pool of unequally weighted models.

The question is how do we assign model weights?



In ecology, model averaging is dominated by the IC framework popularised by Burnham & Anderson (2002).

In ecology, model averaging is dominated by the IC framework popularised by Burnham & Anderson (2002).

We saw earlier that model likelihoods provide a formal measure of evidence for each of the models in the set:

$$\ell_i = e^{(-\frac{\Delta AIC_i}{2})}$$

In ecology, model averaging is dominated by the IC framework popularised by Burnham & Anderson (2002).

We saw earlier that model likelihoods provide a formal measure of evidence for each of the models in the set:

$$\ell_i = e^{(-\frac{\Delta AIC_i}{2})}$$

The ‘probability’ of each model, \mathcal{M}_i , given the data and the N possible models, can be computed as a measure of strength of evidence (Burnham *et al.*, 2011):

$$w_i = \Pr(\mathcal{M}_i | \text{data}) = \frac{\ell_i}{\sum_{i=1}^N \ell_i}$$

In ecology, model averaging is dominated by the IC framework popularised by Burnham & Anderson (2002).

We saw earlier that model likelihoods provide a formal measure of evidence for each of the models in the set:

$$\ell_i = e^{(-\frac{\Delta AIC_i}{2})}$$

The ‘probability’ of each model, \mathcal{M}_i , given the data and the N possible models, can be computed as a measure of strength of evidence (Burnham *et al.*, 2011):

$$w_i = \Pr(\mathcal{M}_i | \text{data}) = \frac{\ell_i}{\sum_{i=1}^N \ell_i} \quad \text{where} \quad \sum_{i=1}^N \ell_i = 1$$




```
library(nlme)
library(MuMIn)

data <- read.csv("Ant_Richness.csv")


FIT <- gls(num_sp ~ latitude + elevation, data = data, method = "ML", na.action = na.fail)

FITS <- dredge(FIT)

FITS

Global model call: gls(model = num_sp ~ latitude + elevation, data = data, method = "ML",
  na.action = na.fail)
```

```
---
Model selection table
  (Intrc)      elvtn  lattd df  logLik AICc delta weight
4  11.120 -0.001373 -0.2018  4   -4.059 18.5  0.00  0.922
2   2.489 -0.001613          3   -8.672 24.7  6.21  0.041
3  12.390          -0.2388  3   -8.857 25.0  6.58  0.034
1   2.114          2  -13.224 31.1 12.61  0.002
Models ranked by AICc(x)
```







```
AVG.FIT <- model.avg(FITS)
```

```
AVG.FIT <- model.avg(FITS)
```

```
summary(AVG.FIT)
```

Component models:

	df	logLik	AICc	delta	weight
12	4	-4.06	18.47	0.00	0.92
1	3	-8.67	24.68	6.21	0.04
2	3	-8.86	25.05	6.58	0.03
(Null)	2	-13.22	31.08	12.61	0.00

Term codes:

elevation	latitude
1	2

Model-averaged coefficients:


(full average)

	Estimate	Std. Error	Adjusted SE	z value	Pr(> z)
(Intercept)	10.7869966	3.2388986	3.3934619	3.179	0.00148 **
elevation	-0.0013333	0.0004967	0.0005211	2.558	0.01051 *
latitude	-0.1943997	0.0758035	0.0794094	2.448	0.01436 *

(conditional average)

	Estimate	Std. Error	Adjusted SE	z value	Pr(> z)
(Intercept)	10.7869966	3.2388986	3.3934619	3.179	0.00148 **
elevation	-0.0013832	0.0004323	0.0004612	2.999	0.00271 **
latitude	-0.2031600	0.0650028	0.0693561	2.929	0.00340 **

Estimated Intercepts






IC based model averaging via MuMIn is one of the many ways you can carry out model averaging

IC based model averaging via MuMIn is one of the many ways you can carry out model averaging, but is any one method better than any other?


IC based model averaging via MuMIn is one of the many ways you can carry out model averaging, but is any one method better than any other? Also is model averaging better than just picking a model?

IC based model averaging via MuMIn is one of the many ways you can carry out model averaging, but is any one method better than any other? Also is model averaging better than just picking a model?



(Dormann *et al.*, 2018)

IC based model averaging via MuMIn is one of the many ways you can carry out model averaging, but is any one method better than any other? Also is model averaging better than just picking a model?



(Dormann *et al.*, 2018)

“We found little in our results to justify the dominance of AIC-based model averaging. And model-averaging did not necessarily outperform single models.”



Model averaging has no super-powers. Like most other statistical methods, model averaging has benefits and costs, and you must weight them to decide which approach is best for your problem.

Model averaging has no super-powers. Like most other statistical methods, model averaging has benefits and costs, and you must weight them to decide which approach is best for your problem.

Benefits include a possible reduction of predictive error and improved parameter estimates.

Model averaging has no super-powers. Like most other statistical methods, model averaging has benefits and costs, and you must weight them to decide which approach is best for your problem.

Benefits include a possible reduction of predictive error and improved parameter estimates.


Costs include extra work/computation time, the fact that it does not always work, and that confidence intervals and p-values are difficult to provide.

Model Selection and Averaging


Recap



Our goal when building models is to identify the fit that optimally balances over- and under-fitting.




Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.


Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.

Know your data


Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.

Know your data, keep your research question in mind


Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.

Know your data, keep your research question in mind, proceed cautiously


Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.


Know your data, keep your research question in mind, proceed cautiously, check model assumptions

Our goal when building models is to identify the fit that optimally balances over- and under-fitting.



In practice, there is no perfect solution for doing this and how you proceed is part science part art.

Know your data, keep your research question in mind, proceed cautiously, check model assumptions, and always check model quality/performance.



References

- Burnham, K.P. & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer New York, New York, NY.
- Burnham, K.P., Anderson, D.R. & Huyvaert, K.P. (2011). AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology*, 65, 23–35.
- Dormann, C.F., Calabrese, J.M., Guillera-Aroita, G., Matechou, E., Bahn, V., Bartoń, K., Beale, C.M., Ciuti, S., Elith, J., Gerstner, K. *et al.* (2018). Model averaging in ecology: A review of bayesian, information-theoretic, and tactical approaches for predictive inference. *Ecological Monographs*, 88, 485–504.
- Murtaugh, P.A. (2014). In defense of p values. *Ecology*, 95, 611–617.