

## 2.2 Linear Equations

- **Definition:** A first-order DE is called a **linear equation** in  $y$  if it can be written in the form

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

When  $g(x) = 0$ , the equation is also called **homogeneous**; otherwise it is **non-homogeneous**.

The **standard form** of a linear equation is obtained by dividing by  $a_1(x)$ , giving

$$\frac{dy}{dx} + P(x)y = f(x).$$

**Claim:** If  $y_c$  is a solution to the homogeneous DE  $y' + P(x)y = 0$ , and  $y_p$  is a solution to the non-homogeneous  $y' + P(x)y = f(x)$ , then  $y = y_c + y_p$  is a solution to the equation  $y' + P(x)y = f(x)$ .

**Proof:** By substitution in the left-hand side of the linear equation

$$\frac{d}{dx}[y_c + y_p] + P(x)[y_c + y_p] = \left(\frac{dy_c}{dx} + P(x)y_c\right) + \left(\frac{dy_p}{dx} + P(x)y_p\right) = 0 + f(x)$$

since  $y_c$  is a solution to the homogeneous equation and  $y_p$  is a solution to the non-homogeneous case. Therefore,  $y = y_c + y_p$  is a solution to the non-homogeneous case as well.

So in order to find a general solution to a linear equation, we need to find a  $y_c$  and a  $y_p$ .

The equation  $y' + P(x)y = 0$  is a separable DE. By separation of variables, we get

$$\int \frac{dy}{y} = - \int P(x)dx \rightarrow y_c = ce^{-\int P(x)dx}.$$

For convenience in future work, we let  $y_c = cy_1$ , where  $y_1 = e^{-\int P(x)dx}$ .

We can now use a technique called **variation of parameters** to find  $y_p$ . For this we want to find a function  $u(x)$  so that  $y_p = u(x)y_1(x)$  is a solution to the non-homogeneous case. By substitution we get

$$\frac{d}{dx}[uy_1] + P(x)uy_1 = uy_1' + y_1u' + P(x)uy_1 = u\left[\frac{dy_1}{dx} + P(x)y_1\right] + y_1\frac{du}{dx} = f(x).$$

Since  $y_1$  is a solution to the homogeneous case, we are left with

$$y_1 \frac{du}{dx} = f(x) \rightarrow u = \int \frac{f(x)}{y_1(x)} dx \rightarrow y_p = e^{-\int P(x)dx} \int e^{\int P(x)dx} f(x) dx.$$

Thus we have a solution of  $y = ce^{-\int P(x)dx} + e^{-\int P(x)dx} \int e^{\int P(x)dx} f(x) dx$ . It is not hard to show that  $y$  represents a **general solution** to the problem.

In general, when solving a linear equation we do not go through all of that work. The key part of the formula is  $e^{\int P(x)dx}$ , which is referred to as the **integrating factor**.

### Solving a Linear First-Order Equation

- i. Put the linear equation into standard form.
- ii. Identify  $P(x)$  and find the integrating factor  $e^{\int P(x)dx}$ .
- iii. Multiply the standard form by the integrating factor. The left-hand side can then be re-written as a derivative, resulting in:

$$\frac{d}{dx} [e^{\int P(x)dx} y] = e^{\int P(x)dx} f(x)$$

- iv. Integrate both sides of the equation and solve for  $y$ .

*\*The constant of integration resulting from  $\int P(x)dx$  can be dropped (or assumed to be 0) when find the integrating factor. The proof of this is left as an exercise.*

- *Example:* Solve the following DE.

$$2x \frac{dy}{dx} - 6xy = 12x$$

Solution: First get the equation in standard form by dividing by  $2x$ .

$$\frac{dy}{dx} - 3y = 6$$

Since  $P(x) = -3$ , the integrating factor would be  $e^{\int -3dx} = e^{-3x}$ . Multiplying both sides by the integrating factor we get

$$\frac{d}{dx} [e^{-3x} y] = 6e^{-3x} \rightarrow e^{-3x} y = \int 6e^{-3x} dx = -2e^{-3x} + c.$$

Dividing by  $e^{-3x}$ , we get a solution of  $y = ce^{3x} - 2$ .

*\*If we wished to break down the solution, we would get that  $y_c = e^{3x}$  and  $y_p = -2$ .*

- *Example:* Solve the following DE.

$$x \frac{dy}{dx} - 5y = x^7 e^x$$

Solution: First get the equation in standard form by dividing by  $x$ .

$$\frac{dy}{dx} - \frac{5}{x}y = x^6 e^x$$

We identify that  $P(x) = -5/x$ , which is continuous on the interval  $(0, \infty)$ . The integrating factor would be  $e^{\int -5/x dx} = e^{-5 \ln |x|} = |x|^{-5} = x^{-5}$ , since  $x > 0$ .

Multiplying both sides by the integrating factor we get

$$\frac{d}{dx}[x^{-5}y] = x e^x \rightarrow x^{-5}y = \int x e^x dx = x e^x - e^x + c.$$

Dividing by  $x^{-5}$ , we get a solution of  $y = x^6 e^x - x^5 e^x + c x^5$ .