

PATIENT FULL NAME

DATE OF BIRTH

DATE OF SERVICE

MEDICATION PRESCRIBED

STARTING DOSE

#### ABOUT THIS TREATMENT

GLP-1 receptor agonists (including semaglutide and tirzepatide) are medications used for chronic weight management and/or type 2 diabetes management. These medications may be commercially available brand-name products or compounded through a licensed 503A pharmacy. I understand that compounded medications are not FDA-approved finished drug products and are prescribed based on my individual clinical need under provider supervision.

**Compounded Medication Notice:** If receiving a compounded GLP-1, I acknowledge it is prepared by a licensed compounding pharmacy, is not FDA-approved, and is prescribed for my individual need. I understand the regulatory status of compounded GLP-1 medications may change and my provider will advise me of any changes affecting my prescription.

#### RISKS & SIDE EFFECTS

- Nausea, vomiting, diarrhea, or constipation — very common
- Fatigue, especially during dose escalation
- Pancreatitis — severe abdominal pain, nausea, vomiting
- Acute kidney injury from dehydration due to GI side effects
- Muscle loss if protein intake is insufficient
- Decreased appetite and food tolerance changes
- Hypoglycemia especially if taking insulin or sulfonylureas
- Gallbladder disease or gallstones (rapid weight loss)
- Hair loss (telogen effluvium) from rapid weight loss
- Possible thyroid C-cell tumors — animal studies, unknown human risk

**Black Box Warning — Thyroid Cancer:** GLP-1 medications caused thyroid C-cell tumors in animal studies. Human risk is unknown. These medications are **contraindicated** in patients with personal or family history of medullary thyroid carcinoma (MTC) or Multiple Endocrine Neoplasia type 2 (MEN 2). I confirm I have no such history unless disclosed below.

#### MEDICAL SCREENING — CIRCLE YES OR NO

☐ Y ☐ N Personal/family history of medullary thyroid carcinoma (MTC)

☐ Y ☐ N Personal/family history of MEN 2 syndrome

☐ Y ☐ N History of pancreatitis

☐ Y ☐ N Pregnant, planning pregnancy, or breastfeeding

☐ Y ☐ N Type 1 or Type 2 diabetes

☐ Y ☐ N History of gallbladder disease

☐ Y ☐ N Severe kidney or liver disease

☐ Y ☐ N Active eating disorder

☐ Y ☐ N Taking insulin or sulfonylureas

☐ Y ☐ N Taking oral contraceptives or thyroid medication

CURRENT MEDICATIONS & DOSES

KNOWN ALLERGIES

#### PATIENT ACKNOWLEDGMENTS

- ☐ I understand GI side effects are most common in the first 4–8 weeks and will not rush my dose titration.
- ☐ I will meet minimum daily protein targets and attend all follow-up appointments.
- ☐ I understand that weight regain is expected if this medication is discontinued.
- ☐ I will contact the practice immediately if I experience severe abdominal pain, persistent vomiting, or signs of pancreatitis.

- ☐ I confirm I have no personal or family history of MTC or MEN 2 unless disclosed above.
- ☐ I am 18 years or older and consent voluntarily to this treatment.

By signing below, I confirm I have read and understand this entire consent form and voluntarily consent to this treatment/service.

PATIENT / CLIENT SIGNATURE

DATE

FOR PROVIDER / TECHNICIAN USE ONLY

|                                    |                           |                     |
|------------------------------------|---------------------------|---------------------|
| PRESCRIBING PROVIDER & CREDENTIALS | MEDICATION / DOSE / LOT # | PHARMACY NAME       |
| BASELINE WEIGHT / BMI              | PROVIDER SIGNATURE        | NEXT FOLLOW-UP DATE |