

Redshift evolution of the underlying type Ia supernova stretch distribution

N. Nicolas^{*}, M. Rigault^{**},

Y. Copin¹, R. Graziani², G. Aldering³, M. Briday¹, Y.-L. Kim¹, J. Nordin⁴, Saul
Perlmutter³, and M. Smith^{1,5}

¹ Univ Lyon, Univ Claude Bernard Lyon 1, CNRS, IP2I Lyon / IN2P3, IMR 5822, F-69622, Villeurbanne, France

² Université Clermont Auvergne, CNRS/IN2P3, Laboratoire de Physique de Clermont, F-63000 Clermont-Ferrand, France.

³ Physics Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA, 94720

⁴ Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489 Berlin

⁵ University of Southampton: Southampton, GB

Submitted to A&A the 19th of May 2020

ABSTRACT

The detailed nature of type Ia supernovae (SNe Ia) remains uncertain, and as survey statistics increase, the question of astrophysical systematic uncertainties arises, notably that of the evolution of SN Ia populations. We study the dependence on redshift of the SN Ia SALT2.4 light-curve stretch, which is a purely intrinsic SN property, to probe its potential redshift drift. The SN stretch has been shown to be strongly correlated with the SN environment, notably with stellar age tracers. We modeled the underlying stretch distribution as a function of redshift, using the evolution of the fraction of young and old SNe Ia as predicted using the SNfactory dataset, and assuming a constant underlying stretch distribution for each age population consisting of Gaussian mixtures. We tested our prediction against published samples that were cut to have marginal magnitude selection effects, so that any observed change is indeed astrophysical and not observational in origin. In this first study, there are indications that the underlying SN Ia stretch distribution evolves as a function of redshift, and that the young/old **[Note 1: the slash commonly denotes "ratio", as in "S/N" for "signal-to-noise ratio". The way you use it here is ambiguous: do you mean "and" or "or", or maybe a range? Please rephrase for clarity]** drifting model is a better description of the data than any time-constant model, including the sample-based asymmetric

^{*}n.nicolas@ip2i.in2p3.fr, equal contribution

^{**}m.rigault@ip2i.in2p3.fr, equal contribution

distributions that are often used to correct Malmquist bias at a significance higher than 5σ . The favored underlying stretch model is a bimodal one, composed of a high-stretch mode shared by both young and old environments, and a low-stretch mode that is exclusive to old environments. The precise effect of the redshift evolution of the intrinsic properties of a SN Ia population on cosmology remains to be studied. The astrophysical drift of the SN stretch distribution does affect current Malmquist bias corrections, however, and thereby the distances that are derived based on SN that are affected by observational selection effects. We highlight that this bias will increase with surveys covering increasingly larger redshift ranges, which is particularly important for the [Note 2: please spell this out here and in the main text, you use the abbreviation only one time each] LSST.

Key words. Cosmology – Type Ia Supernova – Systematic uncertainties

1. Introduction

Type Ia supernovae (SNe Ia) are powerful cosmological distance indicators that enabled the discovery ~~of the acceleration of the Universe's expansion~~ that the Universe expands (Riess et al. 1998; Perlmutter et al. 1999). They remain ~~today~~ a key cosmological probe ~~to understand today for understanding~~ the properties of dark energy (DE) ~~as it~~ because it [Note 3: grammatically, "it" here refers to "DE". Is this correct, or should this be "they" because you still refer to the SNe Ia, or do you mean "the probe"? If you mean the probe, please rephrase to "because this probe" for clarity] is the only tool ~~able to~~ that can precisely map the recent expansion rate ($z < 0.5$) ~~when DE is driving it~~ (e.g., Scolnic et al. 2019). ~~They also are~~ (e.g., Scolnic et al. 2019). Type Ia supernovae are also the key to directly ~~measure the Hubble Constant~~ measuring the Hubble constant (H_0) ~~provided one can calibrate if~~ their absolute magnitude can be calibrated (Riess et al. 2016; Freedman et al. 2019). Interestingly, the value of H_0 ~~that is~~ derived when the SNe Ia are anchored to Cepheids ~~(the SH0ES project, Riess et al. 2009, 2016)~~ (the [Note 4: please provide the spelled-out versions of all instruments and surveys]) $\sim 5\sigma$ higher than what is predicted from cosmic microwave background (CMB) data measured by Planck assuming the standard Λ CDM (Planck Collaboration et al. 2020; Riess et al. 2019; Reid et al. 2019), or when the SN luminosity is anchored at intermediate redshift by the baryon acoustic oscillation (BAO) scale (Feeney et al. 2019). While using the tip of the red giant branch technique in place of the Cepheids ~~seem~~ seems to favor an intermediate value of H_0 (Freedman et al. 2019, 2020), ~~time delay~~ time-delay measurements from strong lensing ~~seem to also~~ also appear to favor high H_0 values (Wong et al. 2020).

The H_0 ~~tension has received a lot of attention~~ as a conundrum has received much attention ~~because~~ it could be a sign of new fundamental physics. ~~Yet, no~~ No simple solution is so far able to accommodate this H_0 ~~tension when accounting for conundrum when~~ all other probes are accounted for, however (Knox & Millea 2020). Alternatively, systematic effects affecting one or several of the aforementioned analyses might explain at least some of this ~~tension~~ discrepancy. Rigault et al.

(2015) suggested that SNe Ia from the Cepheid calibrator sample differ by construction from those in the Hubble flow sample, as the former strongly favors young stellar populations, while the latter does not. This selection effect would ~~impact~~ affect the derivation of H_0 if SNe Ia from young and older environments ~~differ~~ differed in average standardized magnitudes.

The relation between SNe Ia and their host galaxy properties has been studied extensively. The first key finding was that the standardized SNe Ia magnitudes significantly depend on the host galaxy stellar mass, with SNe Ia from high-mass host galaxies being brighter on average

(e.g., Kelly et al. 2010; Sullivan et al. 2010; Childress et al. 2013; Betoule et al. 2014; Kim et al. 2019; Smith et al. 2020) (e.g., Kelly et al. 2010; Sullivan et al. 2010; Childress et al. 2013; Betoule et al. 2014; Kim et al. 2019; Smith et al. 2020).

This mass-step correction is currently used in cosmological analyses (e.g., Betoule et al. 2014; Scolnic et al. 2018);

~~including for deriving~~ (e.g., Betoule et al. 2014; Scolnic et al. 2018), ~~including to derive~~ H_0 (Riess et al. 2016, 2019). ~~Yet, the~~ The underlying connection between the SNe and their host galaxies

remains unclear, ~~particularly when using~~ however, particularly when global properties such as the host stellar mass are used. This in turn raises the question of the accuracy of such corrections ~~since~~

~~host galaxy global properties~~ because the global properties of the host galaxies evolve with redshift. More recently, studies have used the local SN host galaxy environment to probe more direct

connections between the SNe and their host galaxy environments (Rigault et al. 2013), showing

that local age tracers such as the ~~Local specific Star Formation Rate~~ local specific star formation rate (LsSFR) or the local color are more strongly correlated with the standardized SN magnitude

(Roman et al. 2018; Kim et al. 2018; Rigault et al. 2020). ~~Further reinforcing this connection~~

~~is the~~ The identification of SN Ia spectral features that are correlated with LsSFR (Nordin et al. 2018) further support this connection. These results suggest age as the driving parameter underlying

the ~~mass step~~. ~~If true, this would~~ mass step. ~~If this is true, it would~~ have a significant ~~impact for~~ cosmology, ~~since~~ effect for cosmology because the environmental correction that would need to be

applied to ~~apply to~~ SN standardization could strongly vary with redshift (Rigault et al. 2013; Childress et al. 2014; Scolnic et al. 2018). ~~Yet, the~~ The importance of local SN environmental studies

remains highly debated (e.g., Jones et al. 2015, 2019), ~~however~~ (e.g., Jones et al. 2015, 2019), and especially the ~~impact effect~~ of such an astrophysical bias has on the derivation of H_0 (Jones et al. 2015; Riess et al. 2016, 2018; Rose et al. 2019).

The concept of the SN Ia age dichotomy arose with the study of the SN Ia rate. **[Note 5: to properly**

include these references into the main text, please separate them by commas and set off the last reference additionally with "and". This is a LaTeX command error that I cannot fix for you with the program I work with (citet and citep)] Mannucci et al. (2005); Scannapieco &

Bildsten (2005); Sullivan et al. (2006); Smith et al. (2012) have shown that the relative SNe Ia rate in galaxies could be explained if two populations existed, ~~one young, following a young population~~

that follows the host star formation activity, and ~~one old following an old population that followed~~

the host stellar mass (the ~~so-called "prompt and delayed"~~ or "so-called prompt-and-delayed or A+B" model). Rigault et al. (2020) used the LsSFR to ~~classify which are~~ determine the younger (those

with a high LsSFR) and ~~which are~~ the older (those with a low LsSFR) populations. Furthermore,

since the first SNe Ia host analyses, the SN stretch has been known to be strongly correlated with the SN host galaxy properties (Hamuy et al. 1996, 2000). This correlation ~~that~~ has been extensively confirmed since ~~(e.g., Neill et al. 2009; Lampeitl et al. 2010; Gupta et al. 2011; D’Andrea et al. 2011; Pan et al. 2014)~~ then ~~(e.g., Neill et al. 2009; Lampeitl et al. 2010; Gupta et al. 2011; D’Andrea et al. 2011; Pan et al. 2014)~~. Following the “A+B” model and the connection between SN stretch and host galaxy properties, Howell et al. (2007) first discussed the potential redshift drift of the SN stretch distribution. In this paper we revisit this question using the most recent SNe Ia datasets.

~~In this paper, we take a~~ We here step aside from the cosmological analyses to probe the validity of our modeling of the SN population, which we claim ~~to be is~~ constituted of two ~~age-populations~~ age populations (Rigault et al. 2013, 2015, 2020): ~~one old and one younger~~ an older and a younger population, the former having ~~on average lower lightcurve stretches~~ lower light-curve stretches on average and being brighter after standardization. We use the correlation between the SN age ~~;~~ as probed by the LsSFR ~~;~~ and the SN stretch to model the expected evolution of the underlying SN stretch distribution as a function of redshift. This modeling relies on three assumptions: (1) ~~there~~ There are two distinct populations of SNe Ia ~~;~~; (2) the relative fraction of each of these populations as a function of redshift follows the model presented in Rigault et al. (2020), and (3) the underlying distribution of stretch for each age sample is constant. This paper ~~aims at testing tests~~ this specific model with datasets from the literature.

We present ~~in Section 2~~ the sample we ~~are using used~~ for this analysis ~~;~~ in Section 2. It was derived from the Pantheon catalog (Scolnic et al. 2018). We discuss the importance of obtaining a “complete” sample, ~~i.e. that is, a sample that is~~ representative of the true underlying SNe Ia distribution, and how we ~~build built~~ one from the Pantheon sample. We then present ~~in Section 3~~ our modeling of the distribution of stretch in Section 3 and our results ~~are presented~~ in Section 4. In this section, we test whether the SN stretch distribution evolves as a function of redshift and ~~if determine whether~~ the aforementioned age model ~~is in good agreement~~ agrees well with this evolution. We briefly discuss these results in the context of SN cosmology in Section 5, and we conclude in Section 6.

2. Complete ~~Sample Construction~~ sample construction

The ideal SN Ia sample for studying this question would be a very deep, large-area, volume-limited sample. This would capture the true underlying stretch distribution function, and we would then study how it evolves with redshift. No such sample exists, ~~so first we must construct subsamples~~ therefore we must first construct subsamples from existing high-redshift SN Ia samples that are as near to volume-limited as possible ~~from existing high-redshift SN Ia samples~~.

2.1. Applying redshift cuts

We ~~base based~~ our analysis on the most recent comprehensive SNe Ia compilation, the Pantheon catalog from Scolnic et al. (2018). A naive approach to ~~test testing~~ the SN stretch redshift drift

would be to simply compare the observed SN stretch distributions in a few ~~bins of redshift~~ redshift bins. In practice, however, differential observational selection effects will affect the observed SN stretch distributions. ~~Indeed, because~~ Because the observed SN Ia magnitude correlates with the ~~lightcurve~~ light-curve stretch (and color), the first SNe Ia that a magnitude-limited survey will miss are ~~the lowest stretch (and reddest) ones~~ those at the lowest stretch (which are the reddest). Consequently, if magnitude-related observational selection effects are not accounted for, ~~one might confuse~~ one might be confused true population drift with survey properties, and ~~conversely~~ vice versa.

Assuming sufficient (and unbiased) spectroscopic follow-up for acquiring SN types and host galaxy redshifts, the observation selection effects of magnitude-limited surveys should be negligible below a given redshift at which even the faintest normal SNe Ia can be observed. Aiding in the construction of nearly volume-limited subsamples is the fact that the SN Ia population trails off ~~towards~~ toward fainter SNe Ia. A complication is that complete spectroscopic follow-up has not always been the norm, as discussed below. In contrast, targeted surveys have highly complex observational selection functions and so are discarded from our analysis. High-redshift SN cosmology samples, such as ~~Pantheon~~ the Pantheon sample, are predominately assembled from magnitude-limited surveys from which volume-limited SN Ia subsamples can be constructed.

We present in Fig. 1 the ~~lightcurve~~ light-curve stretch and color of SNe Ia from the following surveys: PanStarrs (PS1 [Rest et al. 2014](#)), the Sloan Digital Sky Survey (SDSS [Frieman et al. 2008](#)), and the SuperNovae Legacy Survey (SNLS [Astier et al. 2006](#)). An ellipse in the SALT2 (x_1, c) plane with $x_1 = \pm 3$ and $c = \pm 0.3$ encapsulates the full parent distribution ([Guy et al. 2007](#); [Betoule et al. 2014](#)); see also [Bazin et al. \(2011\)](#) and [Campbell et al. \(2013\)](#) for similar contours, the second using a more conservative $|c| \leq 0.2$ cut. Assuming the SN absolute magnitude with $x_1 = 0$ and $c = 0$ is $M_0 = -19.36$ ([Kessler et al. 2009](#); [Scolnic et al. 2014](#)), we can derive the absolute standardized magnitude at maximum of light $M = M_0 - \alpha x_1 + \beta c$ along within the aforementioned ellipse given the standardization coefficient $\alpha = 0.156$ and $\beta = 3.14$ from [Scolnic et al. \(2018\)](#): ~~the~~ The faintest SN Ia is that with $(x_1 = -1.65, c = 0.25)$ and an absolute standardized magnitude at peak in Bessel B band of $M_{\min}^0 = -18.31$ mag. ~~Since one ought to detect this object typically~~ This object typically ought to be detected 5 days before and a week after peak to build a suitable ~~lightcurve~~ light curve, the effective limiting standardized absolute magnitude is approximately $M_{\lim} = -18.00$ mag. Hence, given the magnitude limit m_{\lim} of a magnitude limited survey, ~~one we~~ one can derive the maximum redshift z_{\lim} above which the faintest SNe Ia will be missed using the relation between apparent magnitude, redshift, and absolute magnitude $\mu(z_{\lim}) = m_{\lim} - M_{\lim}$.

We ~~will thus consider~~ therefore considered a set of cuts that ~~will define~~ defines a first fiducial sample, taking the limits as initially suggested by the previous completeness analysis. However, as this solution might be an overly simplified way to create a complete sample, ~~e.g. for example~~, because it ignores spectroscopic follow-up in efficiency for redshifts below z_{\lim} , we also ~~consider~~ considered another set of cuts to define a so-called ~~“conservative” sample~~ “conservative” sample. ~~The latter conservative sample~~ This is smaller and therefore less statistically constraining, but also even less prone to

Fig. 1. SALT2.4 stretch (x_1) and color (c) ~~lightcurve-light-curve~~ parameters of SNe Ia from the SDSS, PS1, and SNLS samples of the Pantheon catalog. The individual SNe are shown as blue dots. The ellipse ($x_1 = \pm 3, c = \pm 0.3$) is displayed, colored by the corresponding standardized absolute magnitude using the α and β coefficients from Scolnic et al. (2018). The ~~gray~~ diagonal ~~gray~~ lines represent the (x_1, c) evolution for $m = m_{\text{lim}}$ for z between 0.50 and $z = 1.70$ using the SNLS ~~s-~~ m_{lim} of 24.8 mag.

observational selection effects. If the redshift drift is still significant in the conservative sample, it would be even more meaningful in a ~~carefully-tailored~~ ~~carefully tailored~~ selection-free sample. These samples are adequate for the goal of this study, which is to develop a first implementation of a model for drift in SN Ia properties. If fruitful, the sample selection can later be refined ~~,if needed,~~ ~~as required~~ with a more detailed model of the observational selection, ~~e-g-for instance,~~ using the SNANA package (Kessler et al. 2009).

The SNLS typically acquired SNe Ia in the redshift range $0.4 < z < 0.8$; at these redshifts, the rest-frame Bessel B band roughly corresponds to the SNLS i filter, which has a 5σ depth of 24.8 mag¹. This converts ~~to-into~~ a $z_{\text{lim}} = 0.60$, in agreement with Neill et al. (2006), Perrett et al. (2010), and Bazin et al. (2011). Fig. 14 of Perrett et al. (2010, see their Section 5) ~~suggests however,~~ ~~however, suggests~~ a lower limit of $z_{\text{lim}} = 0.55$. We ~~will-therefore-use-therefore used~~ $z = 0.60$ and $z = 0.55$ as redshift limits for the fiducial and conservative samples, respectively, for the SNLS.

Similarly, PS1 observed SNe Ia in the range $0.2 < z < 0.4$, their g -band 5σ depth is 23.1 mag (Rest et al. 2014), which yields $z_{\text{lim}} = 0.31$, in agreement with ~~,e.g.,~~ Fig. 6 of Scolnic et al. (2018), ~~for example~~. If we were to be conservative, this figure would also suggest of a more stringent $z_{\text{lim}} = 0.27$ cut; ~~hence, we will use we therefore used~~ 0.31 and 0.27 for our fiducial and conservative samples, respectively, for PS1.

In a similar redshift range, the SDSS has a limiting magnitude of 22.5 (Dilday et al. 2008; Sako et al. 2008), which would lead to $z_{\text{lim}} = 0.24$. However, the SDSS surveys had to contend with limited spectroscopic resources. As discussed in Kessler et al. (2009, Section 2), during the first year of SDSS, SNe Ia with $r < 20.5$ mag were favored for spectroscopic follow-up, corresponding to a redshift cut at 0.15. For the ~~rest-of-the-remaining~~ SDSS survey, additional spectroscopic resources were available, and Kessler et al. (2009) and Dilday et al. (2008) ~~show-showed~~ a reasonable completeness up to $z_{\text{lim}} = 0.2$. Based on this, we ~~will-use-used~~ $z_{\text{lim}} = 0.20$ and $z_{\text{lim}} = 0.15$ for our fiducial and conservative samples, respectively, for the SDSS.

The sample selection is summarized in Table 1, and the redshift distribution of these three surveys is shown in Fig. 2. As expected, the selected redshift limits are roughly located slightly before the peak of these histograms. In Section 2.2 we ~~validate-confirm~~ that these redshift limits are effective for constructing nearly volume-limited subsamples from samples that were initially more closely ~~magnitude-limited~~ ~~magnitude limited~~ in their search or spectroscopic follow-up.

¹ CFHT final release website.

Table 1. Composition of the SNe Ia dataset used in this analysis. Conservative cuts are indicated in parentheses. The SNf limit is set by Rigault et al. (2020), see text.

Survey	z_{lim}	N_{SN}
SNf	0.08	114
SDSS	0.20 (0.15)	167 (82)
PS1	0.31 (0.27)	160 (122)
SNLS	0.60 (0.55)	102 (78)
HST	–	26
Total	–	569 (422)

Fig. 2. From top to bottom: Redshift histograms of SNe Ia from the SDSS, PS1, and SNLS dataset respectively (data from Pantheon, Scolnic et al. 2018). The colored parts represent the distribution of SNe Ia kept that we retained in our analysis for because they are supposedly assumed to be free from observational selection bias (see Section 2). The darker (resp. lighter) color responds to the conservative (resp. fiducial) selection cut.

In addition, we use-used the SNe Ia from the Nearby Supernova Factory (SNfactory, Aldering et al. 2002) published in Rigault et al. (2020) and that have been discovered from non-targeted nontargeted searches (114 SNe Ia, see their sections 3 and 4.2.2 ; see Aldering et al. 2020). For this dataset, spectroscopic screening was done-performed for candidates with $r \lesssim 19.5$; redshift cuts were then applied when selecting which SN Ia to follow, resulting in a redshift range of $0.02 < z < 0.09$, further reduced to < 0.08 in Rigault et al. (2020) for-extracting-to extract local host properties. These 114 SNfactory SNe Ia are thus in the volume-limited part of the survey (Aldering et al., in prep.), and are therefore assumed to be a random sampling of the underlying SN population. The SNfactory sample is particularly useful for studying SN property drift ,as because it enables us to have a large complete SN Ia sample at $z < 0.1$. Finally, we include the HST sample from Pantheon (Strolger et al. 2004). These high-redshift SNe are of great interest as they provide the greatest leverage for testing evolution. While at these redshifts the supernovae typing is challenging, the target classification was robust enough to include them within-in the cosmological analysis (Scolnic et al. 2018)and-we-do-, and we did not impose further cuts. Section 4 highlights that ,while compatible with it, our results are not dependent on the inclusion of this dataset.

We present the stretch distribution and redshift histogram of these five surveys up to their respective z_{lim} in Fig. 3. We observe here that the fraction of low-stretch SNe (typically $x_1 < -1$) appears to decrease as a function of redshift; this is confirmed in Fig. 6, in which the evolution of the mean stretch is shown, with the data split in redshift bins of regular sample size. We-see-that SNe Ia at higher redshift have on-average-a larger stretch (0.34 ± 0.10 at $z \sim 0.65$) on average than those at lower redshift (-0.17 ± 0.10 at $z \sim 0.05$), suggesting that the underlying stretch distribution is-evolving-evolves with redshift.

Fig. 3. *Bottom:* SALT2.4 ~~lightcurve~~light-curve stretch as a function of redshift for each survey considered in this analysis (see legend). Solid (~~resp.~~open) markers correspond to the conservative (~~resp.~~fiducial) redshift cuts. *Top:* ~~stacked~~Stacked redshift histograms in dark (~~resp.~~light) colors for the conservative (~~resp.~~fiducial) redshift cuts.

2.2. Testing the construction of a volume-limited sample

In section 2.1, we have built volume-limited samples from a set of magnitude-limited ones, using simple redshift cuts. This simplified approach is statistically ~~sub-optimal~~suboptimal, but should ~~be enough~~suffice to test our key question: whether redshift evolution of stretch is compatible with the ~~Rigault et al. (2020) model~~model of Rigault et al. (2020). However, the possibility remains that a complex observational selection function related to spectroscopic follow-up efficiencies below our fiducial (or even conservative) redshift cuts ~~could~~might still affect our sample, making it not fully ~~volume-limited~~volume limited; this would ~~, in turn,~~, in turn bias our conclusion ~~on~~about the astrophysical drift of the SNe Ia population. We now ~~look at~~examine this possibility.

To test for the existence of potential leftover observational selection biases in our sample, we ~~compare~~compared the stretch and color distributions of the SNe Ia originating from different datasets having overlapping redshift ranges: these distributions should be similar if they reflect the underlying parent population. We note that the redshift range has to be narrow enough so that any drift would be negligible.

The two samples that overlap ~~the~~most in redshift are PS1 and SDSS in the redshift range $0.10 < z < 0.20$ (see Fig. 3). This overlapping subsample consists of the 146 SNe Ia at the high-redshift end of SDSS and thus ~~the~~is most likely to be affected by residual observational selection effects (see the corresponding discussion in section 2.1). Over that same redshift range, PS1 has 52 SNe Ia ~~, which~~that are in the lowest redshift bins and thus unlikely to have any observational selection issue. To identify potential inconsistency between the PS1 and SDSS subsamples, Fig. 4 (upper panels) compares the stretch and color distribution of ~~both these~~these two surveys. The resulting Kolmogorov-Smirnov (KS) similarity test p -values ($p > 10\%$) do not support any inconsistency, in agreement with the visual impression from Fig. 4.

We ~~perform~~performed a similar analysis for PS1 and SNLS over the redshift range $0.20 < z < 0.31$ (Fig. 4, lower panels), where the same conclusion can be drawn: ~~there~~There is no substantial sign of discrepancy in the stretch and color distributions between the ~~low-and-high-end~~low and high end of our fiducial SNLS and PS1 samples, respectively. Nonetheless, the small size of the SNLS dataset at $z < 0.31$ (12 SNe Ia vs. 90 for PS1) limits the sensitivity of this test, and only a strong deviation would be noticeable. Extending the redshift range to $0.20 < z < 0.40$ (~~though~~although we have no PS1 data above 0.3) allows ~~to increase~~increasing the SNLS subsample to 31, ~~yet~~but the stretch p -~~values~~value remains high (34%), showing no sign of inconsistency.

We finally highlight that the SNe Ia color is more prone to observational selection effects than stretch, as illustrated in Fig. 1; see also ~~e.g.,~~Fig. 3 of Kessler & Scolnic (2017), for instance.

Fig. 4. x_1 (left) and c (right) distribution histograms of different surveys overlapping in redshift. *Facing up:* SDSS and PS1 within the redshift range $0.10 < z < 0.20$; *facing down:* PS1 and SNLS within the redshift range $0.20 < z < 0.31$. Error bars show the Poisson noise. Our stretch “Base”-base model is illustrated in orange at the mean redshift of the redshift ranges, 0.15 and 0.25, respectively. Kolmogorov-Smirnov test p -values are indicated on-at the top (resp. bottom) of each panel showing-and show no sign that the SDSS and PS1 (resp. PS1 and SNLS) x_1 and c distributions are not drawn from the same underlying distributions.

Hence, since-because the comparison of color distributions shows no significant hint-indication of leftover observational selection effect, it-this further supports our claim that our simple redshift-based selection criteria are sufficient to build the complete SNe Ia samples required to test the redshift evolution of the stretch distribution.

3. Modeling the redshift drift

Rigault et al. (2020) presented a model for the evolution of the fraction of younger and older SNe Ia as a function of redshift following former-previous work on rates and delay-time-delay-time distributions (e.g., Mannucci et al. 2005; Scannapieco & Bildsten 2005; Sullivan et al. 2006; Smith et al. 2012; Childress et al. 2014; Maoz et al. 2014). In short, it was assumed that the number of “young”-young SNe Ia follows the star formation rate (SFR) in the Universe, while the number of “old”-old SNe Ia follows the number of Gyr-old-billion-old stars in the Universe, i.e.-that is, the stellar mass (M^*). Hence, if we denote $\delta(z)$ (resp. $\psi(z) = 1 - \delta(z)$) the fraction of young (resp. old) SNe Ia in the Universe as a function of redshift, then the ratio δ/ψ is expected to follow the evolution of the specific star formation rate (SFR/M^*), which goes as $(1+z)^{-2.8}$ until $z \sim 2$ (e.g., Tasca et al. 2015). Since $\delta(0.05) \sim \psi(0.05)$ (Rigault et al. 2013, 2020; Wiseman et al. 2020), in agreement with rate expectations (Mannucci et al. 2006; Rodney et al. 2014), Rigault et al. (2020) concluded that

$$\delta(z) = \left(K^{-1} \times (1+z)^{-2.8} + 1 \right)^{-1} \quad (1)$$

with $K = 0.87$. This model is comparable to the evolution subsequently predicted by Childress et al. (2014) based on SN rates in galaxies depending on their quenching time as a function of their stellar mass.

3.1. “Base”-underlying stretch distribution

To model the evolution of the full SN stretch distribution as a function of redshift, we need to model the SN stretch distribution for each age subsample given our aforementioned model of the

Fig. 5. *Main:* SALT2 .4 ~~lightcurve-light-curve~~ stretch (x_1) as a function of the ~~local-specific-star formation-rate (LsSFR)~~ for SNfactory SNe ~~used in this analysis~~. The color corresponds to the probability, p_y , for the SNe Ia to be young, i.e., to have $\log \text{LsSFR} \geq -10.82$ (see Rigault et al. 2020). *Right:* p_y -weighted histogram of the SN stretches, as well as the adjusted ~~Base-base~~ model; ~~contributions of~~ the younger and older population ~~contributions~~ are shown in purple and yellow, respectively.

evolution of the fraction of younger and older SNe Ia with cosmic time, ~~we need to model the SN stretch distribution for each age subsample.~~

Rigault et al. (2020) presented the relation between SN stretch and LsSFR measurement, a progenitor age tracer, using the SNfactory sample. This relation is shown in Fig. 5 for the SNfactory SNe used in the current analysis. Given the structure of the stretch-LsSFR scatter plot, our model of the underlying SN Ia stretch distribution is defined as follows [Note 6: please reformat so that this becomes proper text, there is no need for a two-item bulleted list] :

- for the younger population (i.e., $\log(\text{LsSFR}) \geq -10.82$), the stretch distribution is modeled as a single normal distribution $\mathcal{N}(\mu_1, \sigma_1^2)$;
- the older population (i.e., $\log(\text{LsSFR}) < -10.82$) stretch distribution is modeled as a bimodal Gaussian mixture $a \times \mathcal{N}(\mu_1, \sigma_1^2) + (1 - a) \times \mathcal{N}(\mu_2, \sigma_2^2)$, where one mode is the same as for the young population, a representing the relative ~~influence-effect~~ of the two modes.

The stretch probability distribution function (pdf) of a given SN will be the linear combination of the stretch distributions of these two population weighted by its probability y^i to be young (see Section 3.2). ~~But generally,~~ ~~In general, however,~~ the fraction of young SNe Ia as a function of redshift is given by $\delta(z)$ (see Eq. 1) ~~and therefore,~~ ~~and therefore~~ our redshift drift model of the underlying stretch distribution of SNe Ia as a function of redshift $X_1(z)$ is given by ÷

$$X_1(z) = \delta(z) \times \mathcal{N}(\mu_1, \sigma_1^2) + (1 - \delta(z)) \times [a \times \mathcal{N}(\mu_1, \sigma_1^2) + (1 - a) \times \mathcal{N}(\mu_2, \sigma_2^2)] \quad (2)$$

This is our ~~Base-base~~ drifting model.

3.2. Comparison to data

Given the probability y^i that a given SN is young, ~~and assuming our~~ ~~Base-base~~ model (see Section 3.1), the probability ~~to measure of measuring~~ a SALT2 .4 stretch x_1^i with an error dx_1^i is given

Table 2. ~~Best-fit~~ Best-fit values of the parameters for the ~~Base-base~~ stretch distribution model when applied to the SNfactory dataset only (114 SNe Ia), the fiducial 569 SN Ia sample, or the conservative ~~one-model~~ (422).

Sample	μ_1	σ_1	μ_2	σ_2	a
SNfactory	0.41 ± 0.08	0.55 ± 0.06	-1.38 ± 0.10	0.44 ± 0.08	0.48 ± 0.08
Fiducial	0.37 ± 0.05	0.61 ± 0.04	-1.22 ± 0.16	0.56 ± 0.10	0.51 ± 0.09
Conservative	0.38 ± 0.05	0.60 ± 0.04	-1.26 ± 0.13	0.53 ± 0.08	0.47 ± 0.09

by \div

$$\begin{aligned} \mathcal{P}(x_1^i | \theta; dx_1^i, y^i) = & y^i \times \mathcal{N}(x_1^i | \mu_1, \sigma_1^2 + dx_1^{i2}) + \\ & (1 - y^i) \times \left[a \times \mathcal{N}(x_1^i | \mu_1, \sigma_1^2 + dx_1^{i2}) + \right. \\ & \left. (1 - a) \times \mathcal{N}(x_1^i | \mu_2, \sigma_2^2 + dx_1^{i2}) \right]. \end{aligned} \quad (3)$$

The maximum-likelihood estimate of the ~~five~~ free parameters $\theta \equiv (\mu_1, \mu_2, \sigma_1, \sigma_2, a)$ of the model is obtained by minimizing the following:

$$-2 \ln(L) = -2 \sum_i \ln \mathcal{P}(x_1^i | \theta; dx_1^i, y^i). \quad (4)$$

Depending on whether y^i can be estimated directly from LsSFR measurements ~~or not~~, there are two ways to proceed, ~~which we now discuss~~. We discuss them below.

3.2.1. With LsSFR measurements

For the SNfactory sample, we can readily set $y^i = p_y^i$, the probability ~~to have of having~~ log(LsSFR) ≥ -10.82 (see Fig. 5), to minimize Eq. 4 with respect to θ . Results ~~on of~~ fitting the SNf SNe with this model are shown Table 2 and illustrated in Fig. 6.

3.2.2. Without LsSFR measurements

When ~~lacking~~ direct LsSFR measurements are lacking (i.e., p_y^i), we can extend the analysis to non-SNfactory samples by using the ~~redshift-evolution~~ redshift evolution of the fraction $\delta(z)$ of young SNe Ia (Eq. 1) as a proxy for the probability of a SN to be young. This still corresponds to minimizing Eq. 4 with respect to the parameters $\theta \equiv (\mu_1, \mu_2, \sigma_1, \sigma_2, a)$ of the stretch distribution X_1 (Eq. 2), but this time, assuming $y^i = \delta(z^i)$ for any given SN i .

For the ~~rest of the remaining~~ analysis, we ~~will therefore minimize~~ therefore minimized Eq. 4 using p_y^i ~~—, the probability for the SN i to be young—, when available (i.e., for SNfactory dataset), and~~ $\delta(z^i)$ — $\delta(z^i)$, the expected fraction of young SNe Ia at the SN redshift z^i —, otherwise.

Results of fitting this model to all the 569 (resp. 422) SNe from the fiducial (~~resp.~~ conservative) sample are given Table 2, and the predicted redshift evolution of mean stretch (expected x_1 given the distribution of Eq. 2) illustrated as a blue band in Fig. 6 accounting for ~~parameters~~ parameter errors and their covariances. ~~We see in this figure~~ This figure shows that the mea-

Fig. 6. Evolution of the mean SN SALT2.4 stretch (x_1) as a function of redshift. Markers show the stretch plain mean (~~with the~~ error ~~is~~ estimated from ~~the~~ scatter) measured in redshift bins of equal sample size, indicated in light gray at the bottom of each redshift bin. Full and light markers are used when ~~considering~~ the fiducial or ~~the~~ conservative samples ~~are considered~~, respectively. The ~~orange~~-horizontal ~~orange~~ line represents the mean stretch for the ~~non-evolving~~ ~~nonevolving~~ Gaussian model (last line of Table 3) on the fiducial sample. Best fits of our ~~Base~~-~~base~~ drifting model are shown as blue, ~~dashed-blue~~ ~~dashed blue~~, and gray ~~;~~ when fitted on the fiducial sample, the conservative ~~one~~ ~~sample~~, or the SNfactory dataset only, respectively; all are compatible. The ~~light-blue~~ ~~light blue~~ band illustrates the amplitude of the error (~~incl.~~ ~~including~~ covariance) of the ~~best-fit~~ ~~best-fit~~ model when considering the fiducial dataset.

sured mean SN Ia stretch per redshift ~~bins~~ ~~bin~~ of equal sample size closely follows our redshift drift modeling. This is indeed what is expected if old environments favor low SN stretches (~~e.g.~~ ~~Howell et al. 2007~~) (~~e.g.~~ ~~Howell et al. 2007~~) ~~and~~ ~~and~~ if the fraction of old SNe Ia declines as a function of redshift. See Section 4 for a more quantitative discussion.

3.3. Alternative models

In Section 3.1 ~~;~~ we have modeled the underlying stretch distribution following Rigault et al. (2020), ~~i.e. that is~~, as a single Gaussian for the “~~young~~” ~~young~~ SNe Ia and a mixture of two Gaussians for the “~~old~~” ~~old~~ SNe Ia population, one being the same as for the young population, plus another one for the fast-declining SNe Ia that ~~seem~~ ~~appear~~ to only exist in old local environments. This is our so-called “~~Base~~” ~~base~~ model. However, to test different modeling choices, we ~~have~~ implemented a suite of alternative parameterizations that we also ~~adjust~~ ~~adjusted~~ to the data following the procedure described in Section 3.2.2.

Howell et al. (2007) used a simpler unimodal model per age category, assuming a single normal distribution for each of the young and old populations. We thus ~~consider a~~ “~~considered a~~ Howell+drift” model, with one single Gaussian per age group and the $\delta(z)$ drift from Eq. 1.

Alternatively, ~~since we aim at probing because we aim to probe~~ the existence of an evolution with redshift, we also ~~test~~ ~~tested~~ constant models by restricting the “~~Base~~” and “~~Howell~~” ~~base and~~ ~~Howell~~ models to use ~~a supposedly an assumed~~ redshift-independent fraction $\delta(z) \equiv f$ of young SNe; these models are hereafter labeled “~~Base~~base+constant” and “~~and~~ ~~and~~ Howell+constant”.

We also ~~consider another intrinsically non-drifting~~ ~~considered another intrinsically nondrifting~~ model, the functional form developed for ~~Beams with Bias Correction~~ ~~beams with bias correction~~ (BBC, Scolnic & Kessler 2016; Kessler & Scolnic 2017), used in recent SN cosmological analyses

(~~e.g.~~ ~~Scolnic et al. 2018; Abbott et al. 2019; Riess et al. 2016, 2019~~) (~~e.g.~~ ~~Scolnic et al. 2018; Abbott et al. 2019; Riess et al. 2016, 2019~~) to

account for Malmquist biases. The BBC formalism assumes sample-based (hence intrinsically ~~non-drifting~~ ~~nondrifting~~) asymmetric Gaussian stretch distributions: $\mathcal{N}(\mu, \sigma_-^2 \text{ if } x_1 < \mu, \text{ else } \sigma_+^2)$.

The idea behind this sample-based approach is twofold: (1) Malmquist biases are driven by survey

Table 3. Comparison of the relative ability of each model to describe the data. For each considered model, we report ~~if whether~~ the model is drifting ~~or not~~, its number of free parameters ~~and~~, ~~and~~ for both the fiducial and the conservative cuts, $-2 \ln(L)$ (see Eq. 4), the AIC and the AIC difference (ΔAIC) between this model and the ~~Base-base~~ model used as reference ~~for because~~ it has the lowest AIC.

Name	drift	k	Fiducial sample (569 SNe)			Conservative sample (422 SNe)		
			$-2 \ln(L)$	AIC	ΔAIC	$-2 \ln(L)$	AIC	ΔAIC
Base	$\delta(z)$	5	1456.7	1466.7	–	1079.5	1089.5	–
Howell+drift	$\delta(z)$	4	1463.3	1471.3	–4.6	1088.2	1096.2	–6.7
Asymmetric	–	3	1485.2	1491.2	–24.5	1101.3	1107.3	–17.8
Howell+const	f	5	1484.2	1494.2	–27.5	1101.2	1111.2	–21.7
Base+const	f	6	1484.2	1496.2	–29.5	1101.2	1113.2	–23.7
Per sample Asym.	per sample	3×5	1468.2	1498.2	–31.5	1083.6	1113.6	–24.1
Gaussian	–	2	1521.8	1525.8	–59.1	1142.6	1146.6	–57.1

Fig. 7. ΔAIC between “~~Base~~”-~~base~~ model (reference) and other models (see Table 3). Full and open blue markers correspond to models with and without redshift drift, respectively. Light markers show the results when the analysis is performed on the conservative sample rather than the fiducial one. ~~Color-bands~~ ~~Color bands~~ illustrate the validity of the models, from ~~Aacceptable~~ ~~acceptable~~ ($\Delta AIC > -5$) to ~~Excluded~~ ~~excluded~~ ($\Delta AIC < -20$), see text. According to the AIC, all ~~non-drifting~~ ~~nondrifting~~ models (open symbols) are excluded ~~to be~~ as ~~good~~ a ~~poorer~~ representation of the data ~~as than~~ the ~~Base-base~~ (drifting) model.

properties, and (2) ~~since because~~ current surveys cover limited redshift ranges, having a sample-based approach covers some potential redshift evolution information (Scolnic & Kessler 2016; Scolnic et al. 2018). See ~~further discussion concerning a more detailed discussion of~~ BBC in Section 5.

Finally, for the sake of completeness, we also ~~consider~~ ~~considered~~ redshift-independent pure and asymmetric Gaussian models.

4. Results

We adjusted each of the models described above on both the fiducial and conservative samples (cf. Section 2); ~~The~~ results are gathered in Table 3, ~~and and are~~ illustrated in Fig. 7.

~~Since Because~~ the various models have different degrees of freedom, we ~~use the Akaike Information Criterion (AIC, e.g. Burnham & Anderson 2004)~~ ~~used the Akaike information criterion (AIC, e.g., Burnham & Anderson 2004)~~ to compare their ability to properly describe the observations. This estimator penalizes ~~extra additional~~ degrees of freedom to avoid ~~over-fitting the data~~, ~~overfitting the data~~ and is defined as ~~follow~~: ~~follows~~:

$$AIC = -2 \ln(L) + 2k, \quad (5)$$

where $-2 \ln(L)$ is derived by minimizing Eq. (4), and k is the number of free parameters to be adjusted. The reference model ~~is the one with the~~ ~~has the~~ smallest AIC; in comparison to this model, the models with $\Delta AIC < 5$ are coined acceptable, ~~the ones those~~ with $5 < \Delta AIC < 20$ are

~~unfavored~~not favored, and those with $\Delta\text{AIC} > 20$ are deemed excluded. This roughly corresponds to 2, 3, and 5 σ limits for a Gaussian probability distribution.

The best model (with the smallest AIC) is the so-called ~~Base-base~~ model and thus is our reference model; this is true ~~both-on-for~~ the fiducial and conservative samples. The ~~Base-base~~ model also has the smallest $-2\ln(L)$, making it the most likely model even ~~ignoring the over-fitting issue~~ when the overfitting issue that is accounted for by the AIC formalism is ignored.

Furthermore, we find that redshift-independent stretch distributions are all excluded as suitable descriptions of the data relative to the ~~Base-model. In fact, the best non-drifting base model. The best nondrifting~~ model (the ~~Asymmetric-one~~asymmetric model) has a very marginal chance ($p \equiv \exp(\Delta\text{AIC}/2) = 5 \times 10^{-6}$) to describe the data as well as the ~~Base-base~~ model. This result is just a quantitative assessment of qualitative facts that are clearly visible in Fig. 6: ~~the~~ The mean SN stretch per bin of redshift strongly suggests a significant redshift evolution rather than a constant value, and this evolution is well described by Eq. 1.

Surprisingly, the sample-based Gaussian asymmetric modeling used by current implementations of the BBC technique (Scolnic & Kessler 2016; Kessler & Scolnic 2017) has one of the highest AIC ~~value-values~~ in our analysis (see Section 4). While its $-2\ln(L)$ is the smallest of all redshift-independent models (but still -11.5 ~~worse-lower~~ than the reference ~~Base-base~~ model), it is strongly penalized for requiring 15 free parameters (μ_0, σ_{\pm} for each of the ~~5~~five samples of the analysis). Hence, its $\Delta\text{AIC} < -20$, which could be interpreted as a probability $p = 2 \times 10^{-7}$ of being as good a representation of the data as the ~~Base-base~~ model.

We note that ~~when-comparing-models-when models are compared that were~~ adjusted on individual subsamples rather than globally, the Bayesian ~~Information-Criterion-information criterion~~ (BIC = $-2\ln(L) + k\ln(n)$, with n the number of data points) might be better suited than AIC ~~since~~ because it explicitly accounts for the fact that each subsample is fitted separately: the sample-based model BIC is rightfully the sum of the BIC for each sample. We find $\Delta\text{BIC} = -48$, again refuting the sample-based asymmetric Gaussian model as being as pertinent as the ~~Base-base~~ model.

In order to ensure that our results are not driven by the ~~incompletely-modeled-incompletely modeled~~ HST subsample, we ~~recompute-recomputed~~ ΔAIC for each model excluding this dataset; ~~we find that it does~~ this did not change ΔAIC by more than few tenths. The consistency of these values with those in Table 3 ~~show-shows~~ that the HST subsample does not drive our conclusions.

We report in Table 4 our determination of μ_0 and σ_{\pm} for each sample when ~~implementing-an~~ asymmetric Gaussian model ~~was implemented, and~~ adjusted on the nominally selection-free samples using our fiducial cuts (see Section 2). ~~We find our results~~ Our results are in close agreement with ~~Scolnic & Kessler (2016) for those of Scolnic & Kessler (2016) for the~~ SNLS and SDSS and with the results reported by Scolnic et al. (2018) for PS1, who derived these model parameters using the full BBC formalism, using numerous simulations to model the observational selection effects (see details, e.g., Section 3 of Kessler & Scolnic 2017). The agreement between our fit of the asymmetric Gaussians on the supposedly selection-free part of the samples and the results de-

Table 4. Best-fit parameters for our sample-based asymmetric modeling of the underlying stretch distribution.

Asymmetric	σ_-	σ_+	μ_0
SNfactory	1.34 ± 0.13	0.41 ± 0.10	0.68 ± 0.15
SDSS	1.31 ± 0.11	0.42 ± 0.09	0.72 ± 0.13
PS1	1.01 ± 0.11	0.52 ± 0.12	0.38 ± 0.16
SNLS	1.41 ± 0.13	0.15 ± 0.13	1.22 ± 0.15
HST	0.76 ± 0.36	0.79 ± 0.35	0.11 ± 0.44

rived using the BBC formalism supports our approach to constructing a sample with negligible observational selection effects. If we were to use [the Scolnic & Kessler \(2016\)](#) and [Scolnic et al. \(2018\)](#) ~~best-fit~~ [best-fit](#) values of the μ_0, σ_{\pm} asymmetric parameters for [the](#) SNLS, SDSS and PS1, respectively, the ΔAIC between our ~~Base-base~~ drifting model and the BBC modeling would go even deeper from -32 to -47 . We further discuss the consequence of this result for cosmology in Section 5.

We also performed tests allowing the high-stretch mode of the old population to differ from the young population mode, hence adding two degrees of freedom. The corresponding fit is not significantly better, with a ΔAIC of -0.4 . This reinforces our assumption that the young and old populations indeed appear to share the same underlying high-stretch mode. Furthermore, ~~one-we~~ might wonder whether a low-stretch mode might also exist in the ~~young-population~~ [young population](#) (see Fig. 5). We ~~test-tested~~ for this by allowing this population to also be bimodal, finding the amplitude of such a low-stretch mode to be compatible with 0 ($< 2\%$) in this young population. More generally, this raises the question of how well a given environmental tracer (here LsSFR) traces the age. An analysis dedicated to this question will be presented in Briday et al. in prep.

Finally, ignoring the LsSFR measurements ~~—available only—, which are only available~~ for the SNfactory dataset (see Section 3) ~~—~~, reduces the significance of the results presented in this section, as expected. Even so, ~~non-drifting-nondrifting~~ models remain strongly disfavored. For instance, the ~~best-fitting-best-fitting~~ sample-based Gaussian asymmetric model is still ~~$\Delta\text{AIC} < -10$~~ [AIC < -10, which is](#) less representative of the data than our ~~Base-base~~ drifting model.

5. Discussion

To the best of our knowledge, a SN Ia stretch redshift drift modeling has never been explicitly used in cosmological analyses, ~~though-although~~ a Bayesian hierarchy formalism such as UNITY ([Rubin et al. 2015](#)), BAHAMAS ([Shariff et al. 2016](#)), or Steve ([Hinton et al. 2019](#)) can easily allow it (see, e.g., sections 1.3 and 2.5 of [Rubin et al. \(2015\)](#)). Not doing so is a ~~second-order-second-order~~ issue for SN cosmology ~~—as-because~~ it only affects the way ~~one-accounts-for~~ [Malmquist bias](#). ~~Indeed,~~ ~~as-in which the Malmquist bias is accounted for.~~ As long as the Phillips relation ([Phillips 1993](#)) standardization parameter α is not redshift dependent (a study [that is](#) beyond the scope of this paper, but see, e.g., [Scolnic et al. 2018](#)), the stretch-corrected SNe Ia magnitudes used for cosmology are [indeed](#) blind to the underlying stretch distribution for complete samples. However, surveys usually do have significant Malmquist bias for the upper half of their SN redshift distribution. As

Fig. 8. Distribution of the PS1 SN Ia SALT2.4 stretch (x_1) after the fiducial redshift limit cut (gray histogram). This distribution is supposed to be a random draw from the underlying stretch distribution. The green lines show the BBC model of this underlying distribution (asymmetric Gaussian). The full line (band) is our best fit (its error); the dashed line shows the Scolnic et al. (2018) result. The black line (band) shows our ~~best-fitted Base-modeling~~ best-fit base modeling (its error, see Table 2) that includes redshift drift. For illustration, we show ~~as-colored-~~ colored from blue to red with increasing redshifts) the evolution of the underlying stretch distribution as a function of redshift for the redshift range covered by PS1 data.

a consequence, mismodeling of the underlying stretch distribution will bias the SN magnitudes derived from such surveys.

Commonly used Malmquist bias correction techniques, such as the BBC-formalism, assume ~~per-sample-per-sample~~ asymmetric Gaussian functions to model the underlying stretch and color distributions. ~~Yet, as~~ As shown in Section 4, however, such a sample-based distribution is excluded in comparison to our drifting model. ~~Then, contrary~~ In contrast to what Scolnic & Kessler (2016, Section 2) and Scolnic et al. (2018, Section 5.4) have suggested—~~i.e., that, that is, that~~ traditional surveys span sufficiently limited redshift ranges such that the per-sample approach accounts for implicit redshift drifts—, a direct modeling of the redshift drift is therefore more appropriate than a sample-based approach. We add here that ~~, as~~ measurements of modern surveys try to cover increasingly larger redshift ranges in order to reduce calibration systematic uncertainties, this sample-based approach becomes less valid, notably for PS1, DES and, soon, the LSST.

We illustrate in Fig. 8 the prediction difference in the underlying stretch distribution between the per-sample asymmetric modeling and our ~~Base-base~~ drifting model for the PS1 sample. Our model is bimodal, and the relative amplitude of each mode depends on the redshift-dependent fraction of old and young SNe Ia in the sample: the higher the fraction of old SNe Ia (at lower redshift), the higher the amplitude of the old-specific low-stretch mode. This redshift ~~dependency~~ of dependence on the underlying stretch distributions is shown ~~running with colors~~ from blue to red in Fig. 8 for the redshift range covered by PS1. The observed x_1 histogram follows the model we defined using the sum of individual underlying SN-redshift distributions. As expected, the two modeling approaches differ mostly in the negative part of the SN stretch distribution. The asymmetric Gaussian distribution goes through the middle of the bimodal distribution, ~~over-estimating~~ overestimating the number of SNe Ia at $x_1 \sim -0.7$ and ~~under-estimating-underestimating~~ it at $x_1 \sim -1.7$ in comparison to our ~~Base-base~~ drifting model for typical PS1 SN redshifts. This means that the SN bias-corrected standardized magnitude estimated at a redshift ~~suffering from~~ plagued by observational selection effects would be biased by a mismodeling of the true underlying stretch distribution.

Assessing the amplitude of this magnitude bias for cosmology is beyond the scope of this paper given the complexity of the BBC analysis. It would require a full study using our ~~Base-base~~ drifting model

(Eq. 2) in place of the sample-based asymmetric modeling as part of the BBC simulations. However, we already ~~highlight~~ highlighted that even if a ~~non-drifting~~ nondrifting sample-based model could provide comparable result in the volume-limited part of the various samples, these models would differ when ~~extrapolating~~ extrapolated to higher redshifts, precisely where the underlying distribution will matter for correcting Malmquist biases.

In the era of modern cosmology, where we aim to measure w_0 at a ~~sub-percent~~ subpercent level and w_a with ~~ten-percent~~ 10% precision (e.g., Ivezić et al. 2019), we stress that correct modeling of potential SN redshift drift should be further studied and care should be taken when ~~using samples~~ samples are used that are affected by observational selection effects.

6. Conclusion

We have presented an initial study of the drift of the underlying SNe Ia stretch distribution as a function of redshift. We built effectively volume-limited SN Ia subsamples from the Pantheon dataset (~~Scolnic et al. 2018, SDSS, PS1 and SNLS~~) (Scolnic et al. 2018, SDSS, PS1, and SNLS), to which we added HST and SNfactory data from Rigault et al. (2020) for the high- and low-redshift bins. We only considered the SNe that have been discovered in the redshift range of each survey ~~where~~ in which observational selection effects are negligible, so that the observed SNe Ia stretches are a random sampling of the true underlying distribution. This resulted in a fiducial sample of 569 SNe Ia (422 SNe when more conservative cuts were applied).

Following predictions made in Rigault et al. (2020), we introduced a redshift drift model ~~which~~ that depends on the expected fraction of ~~“young” and “old”~~ young and old SNe Ia as a function of redshift, with each age population having its own underlying stretch distribution.

In addition to this ~~“base”~~ base model, we ~~have~~ have studied various distributions, including ~~redshift independent~~ redshift-independent models; we also studied the prediction from a per-sample asymmetric Gaussian stretch distribution used, for instance, by the ~~Beams with Bias Correction~~ beams with bias correction Malmquist bias correction algorithm (Scolnic & Kessler 2016; Kessler & Scolnic 2017).

Our conclusions are ~~the following:~~ listed below.

1. The underlying SN Ia stretch distribution is significantly redshift dependent, as previously suggested by ~~e.g.~~ Howell et al. (2007), ~~for example,~~ in a way that observational selection effects alone cannot explain. This result is largely independent of ~~details on the~~ details of each age-population model.
2. Redshift-independent models are quantitatively excluded as suitable descriptions of the data relative to our ~~Base~~ base model. This model assumes that ~~÷~~ (1) the younger population has a unimodal Gaussian stretch distribution ~~÷~~ while the older population stretch distribution is bimodal, one mode being the same as the young one ~~÷~~ and (2) the evolution of the relative fraction of younger and older SNe Ia follows the prediction made in Rigault et al. (2020). This second result further supports the existence of both young and old SN Ia populations, in

agreement with rate studies (Mannucci et al. 2005; Scannapieco & Bildsten 2005; Sullivan et al. 2006; Aubourg et al. 2008).

3. Models using survey-based asymmetric Gaussian distributions, ~~e.g. for instance~~, as employed in the current implementation of BBC, are excluded as a good description of the data relative to our drifting model. ~~Hence, This means that~~ the sample-based approach does not accurately account for redshift drift, a problem that will be exacerbated as surveys span increasingly larger redshift ranges. As a result, even if the necessary extra degrees of freedom might be acceptable given the large number of SNe Ia in cosmological studies, extrapolating the SN property distributions from the volume-limited part of a survey to its Malmquist-biased magnitude-limited part would still be inaccurate because of the redshift evolution.
4. Given the current dataset, we suggest the use of the following stretch population model as a function of redshift:

$$X_1(z) = \delta(z) \times \mathcal{N}(\mu_1, \sigma_1^2) + (1 - \delta(z)) \times [a \times \mathcal{N}(\mu_1, \sigma_1^2) + (1 - a) \times \mathcal{N}(\mu_2, \sigma_2^2)], \quad (2)$$

with $a = 0.51$, $\mu_1 = 0.37$, $\mu_2 = -1.22$, $\sigma_1 = 0.61$, and $\sigma_2 = 0.56$ (see Table 2), and using the age-population drift model,

$$\delta(z) = (K^{-1} \times (1 + z)^{-2.8} + 1)^{-1} \quad (1)$$

with $K = 0.87$.

~~In this paper, we~~ We considered a simple two-population Gaussian mixture modeling. Additional data free from significant Malmquist bias would enable us to refine it, ~~as necessary~~ as required. We note that samples at the low- and high-redshift ends of the Hubble diagram would be particularly helpful for this drifting analysis; fortunately, this will soon be provided by the Zwicky Transient Facility (low- z , Bellm et al. 2019; Graham et al. 2019) ~~, and~~ and the Subaru and SeeChange SNe Ia programs (high- z), respectively.

The next step in this line of analysis will incorporate our model into the SNANA framework (Kessler et al. 2009), both to more accurately account for observational selection functions and to test the ~~impact-effect~~ of our model on the derivation of cosmological parameters; ~~this study is underway~~. This study is currently being conducted.

Acknowledgements. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 Research and Innovation program (grant agreement no 759194 - USNAC). This work was supported in part by the Director, Office of Science, Office of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC025CH11231. This project is partly financially supported by Région Rhône-Alpes-Auvergne.

References

- Abbott, T. M. C., Allam, S., Andersen, P., et al. 2019, ApJ, 872, L30
- Aldering, G., Adam, G., Antilogus, P., et al. 2002, Proc. SPIE, 61

- Aldering, G., Antilogus, P., Aragon, C., et al. 2020, *Research Notes of the American Astronomical Society*, 4, 63
- Astier, P., Guy, J., Regnault, N., et al. 2006, *A&A*, 447, 31
- Aubourg, É., Tojeiro, R., Jimenez, R., et al. 2008, *A&A*, 492, 631
- Bazin, G., Ruhlmann-Kleider, V., Palanque-Delabrouille, N., et al. 2011, *A&A*, 534, A43
- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
- Betoule, M., Kessler, R., Guy, J., et al. 2014, *A&A*, 568, A22
- Brout, D., Scolnic, D., Kessler, R., et al. 2019, *ApJ*, 874, 150
- Burnham, K., Anderson, D., 2004, *Sociological Methods & Research*, 33, 2
- Campbell, H., D’Andrea, C. B., Nichol, R. C., et al. 2013, *ApJ*, 763, 88
- Childress, M., Aldering, G., Antilogus, P., et al. 2013, *ApJ*, 770, 108
- Childress, M. J., Wolf, C., & Zahid, H. J. 2014, *MNRAS*, 445, 1898
- D’Andrea, C. B., Gupta, R. R., Sako, M., et al. 2011, *ApJ*, 743, 172
- Dilday, B., Kessler, R., Frieman, J. A., et al. 2008, *ApJ*, 682, 262
- Feeney, S. M., Peiris, H. V., Williamson, A. R., et al. 2019, *Phys. Rev. Lett.*, 122, 061105
- Freedman, W. L., Madore, B. F., Hatt, D., et al. 2019, *ApJ*, 882, 34
- Freedman, W. L., Madore, B. F., Hoyt, T., et al. 2020, *ApJ*, 891, 57. doi:10.3847/1538-4357/ab7339
- Frieman, J. A., Bassett, B., Becker, A., et al. 2008, *AJ*, 135, 338
- Graham, M. J., Kulkarni, S. R., Bellm, E. C., et al. 2019, *PASP*, 131, 078001
- Gupta, R. R., D’Andrea, C. B., Sako, M., et al. 2011, *ApJ*, 740, 92
- Guy, J., Astier, P., Baumont, S., et al. 2007, *A&A*, 466, 11
- Hamuy, M., Phillips, M. M., Suntzeff, N. B., et al. 1996, *AJ*, 112, 2391
- Hamuy, M., Trager, S. C., Pinto, P. A., et al. 2000, *AJ*, 120, 1479
- Hinton, S. R., Davis, T. M., Kim, A. G., et al. 2019, *ApJ*, 876, 15
- Howell, D. A., Sullivan, M., Conley, A., et al. 2007, *ApJ*, 667, L37
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., et al. 2019, *ApJ*, 873, 111
- Jones, D. O., Riess, A. G., & Scolnic, D. M. 2015, *ApJ*, 812, 3 1
- Jones, D. O., Riess, A. G., Scolnic, D. M., et al. 2018, *ApJ*, 867, 108
- Jones, D. O., Scolnic, D. M., Riess, A. G., et al. 2018, *ApJ*, 857, 51
- Jones, D. O., Scolnic, D. M., Foley, R. J., et al. 2019, *ApJ*, 881, 19
- Kelly, P. L., Hicken, M., Burke, D. L., et al. 2010, *ApJ*, 715, 743
- Kessler, R., Becker, A. C., Cinabro, D., et al. 2009, *ApJS*, 185, 32
- Kessler, R., Bernstein, J. P., Cinabro, D., et al. 2009, *PASP*, 121, 1028
- Kessler, R., & Scolnic, D. 2017, *ApJ*, 836, 56
- Kim, Y.-L., Smith, M., Sullivan, M., et al. 2018, *ApJ*, 854, 24
- Kim, Y.-L., Kang, Y., & Lee, Y.-W. 2019, *Journal of Korean Astronomical Society*, 52, 181
- Knox, L. & Millea, M. 2020, *Phys. Rev. D*, 101, 043533. doi:10.1103/PhysRevD.101.043533
- Lampeitl, H., Smith, M., Nichol, R. C., et al. 2010, *ApJ*, 722, 566
- Mannucci, F., Della Valle, M., Panagia, N., et al. 2005, *A&A*, 433, 807
- Mannucci, F., Della Valle, M., & Panagia, N. 2006, *MNRAS*, 370, 773
- Maoz, D., Mannucci, F., & Nelemans, G. 2014, *ARA&A*, 52, 107
- Neill, J. D., Sullivan, M., Balam, D., et al. 2006, *AJ*, 132, 1126
- Neill, J. D., Sullivan, M., Howell, D. A., et al. 2009, *ApJ*, 707, 1449
- Nordin, J., Aldering, G., Antilogus, P., et al. 2018, *A&A*, 614, A71
- Pan, Y.-C., Sullivan, M., Maguire, K., et al. 2014, *MNRAS*, 438, 1391
- Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, *ApJ*, 517, 565
- Perrett, K., Balam, D., Sullivan, M., et al. 2010, *AJ*, 140, 518
- Phillips, M. M. 1993, *ApJ*, 413, L105
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, *A&A*, 641, A6. doi:10.1051/0004-6361/201833910

- Reid, M. J., Pesce, D. W., & Riess, A. G. 2019, *ApJ*, 886, L27. doi:10.3847/2041-8213/ab552d
- Rest, A., Scolnic, D., Foley, R. J., et al. 2014, *ApJ*, 795, 44
- Riess, A. G., Filippenko, A. V., Challis, P., et al. 1998, *AJ*, 116, 1009
- Riess, A. G., Macri, L., Casertano, S., et al. 2009, *ApJ*, 699, 539
- Riess, A. G., Macri, L. M., Hoffmann, S. L., et al. 2016, *ApJ*, 826, 56
- Riess, A. G., Casertano, S., Yuan, W., et al. 2018, *ApJ*, 861, 126
- Riess, A. G., Casertano, S., Yuan, W., et al. 2019, *ApJ*, 876, 85
- Rigault, M., Copin, Y., Aldering, G., et al. 2013, *A&A*, 560, A66
- Rigault, M., Aldering, G., Kowalski, M., et al. 2015, *ApJ*, 802, 20
- Rigault, M., Brinnel, V., Aldering, G., et al. 2029, *A&A*, 644, A176
- Rodney, S. A., Riess, A. G., Strolger, L.-G., et al. 2014, *AJ*, 148, 13
- Roman, M., Hardin, D., Betoule, M., et al. 2018, *A&A*, 615, A68
- Rose, B. M., Garnavich, P. M., & Berg, M. A. 2019, *ApJ*, 874, 32
- Rubin, D., Aldering, G., Barbary, K., et al. 2015, *ApJ*, 813, 137
- Rubin, D., & Hayden, B. 2016, *ApJ*, 833, L30
- Sako, M., Bassett, B., Becker, A., et al. 2008, *AJ*, 135, 348
- Saunders, C., Aldering, G., Antilogus, P., et al. 2018, *ApJ*, 869, 167. doi:10.3847/1538-4357/aaec7e
- Scannapieco, E., & Bildsten, L. 2005, *ApJ*, 629, L85
- Scolnic, D., Rest, A., Riess, A., et al. 2014, *ApJ*, 795, 45
- Scolnic, D., & Kessler, R. 2016, *ApJ*, 822, L35
- Scolnic, D. M., Jones, D. O., Rest, A., et al. 2018a, *ApJ*, 859, 101
- Scolnic, D., Perlmutter, S., Aldering, G., et al. 2019, *Astro2020: Decadal Survey on Astronomy and Astrophysics*, 2020, 270
- Shariff, H., Jiao, X., Trotta, R., et al. 2016, *ApJ*, 827, 1
- Smith, M., Nichol, R. C., Dilday, B., et al. 2012, *ApJ*, 755, 61
- Smith, M., Sullivan, M., Wiseman, P., et al. 2020, *MNRAS*, 494, 4426
- Strolger, L.-G., Riess, A. G.,
- Sullivan, M., Le Borgne, D., Pritchett, C. J., et al. 2006, *ApJ*, 648, 868
- Sullivan, M., Conley, A., Howell, D. A., et al. 2010, *MNRAS*, 406, 782
- Tasca, L. A. M., Le Fèvre, O., Hathi, N. P., et al. 2015, *A&A*, 581, A54
- Wiseman, P., Smith, M., Childress, M., et al. 2020, *MNRAS*, 495, 4040. doi:10.1093/mnras/staa1302
- Wong, K. C., Suyu, S. H., Chen, G. C.-F., et al. 2020, *MNRAS*, 498, 1420. doi:10.1093/mnras/stz3094

List of Objects