

www.wuxiapptec.com





© 2012 ABGENT. All Rights Reserved 10239 Flanders Court San Diego, CA 92121, USA Tel: 858.622.0099 www.abgent.com info@abgent.com





K.Gramatikoff<sup>2</sup>PhD, J.Mountzouris<sup>1</sup>PhD, T.Gilliam<sup>1</sup> & C.Wu<sup>1</sup>PhD (1) Abgent Inc., 10239 Flanders Ct, San Diego, CA 92121

(2) Burnham Institute for Medical Research, 10901 North Torrey Pines Road, La Jolla, CA 92037









cells types. The gene symbols of SCs specific and common proteins are indicated on the right. The three SC types (ESCs, MSCs, and NSCs) shared nine proteins identified in proteomics screens, including proteins involved in energy production and metabolisms.

**Distribution** of stem cell proteins in cell subtypes and across cellular function **Associations** of stem cell proteins in major tissue types and among each other **Profiling** of stem cell markers in cancer tissues via antibodies

**Associations** 

## **Abbreviations & References**

plug-in to the Agilent Literature Search (http://www.agilent.com, Ref. 10), and color coded in Adobe Illistrator CS2 (http://www.adobe.com).

HSC, Haematopoietic stem cells; HSC-LT, long-term HSCs; HSC-ST, short-term HSCs; MPP, multipotential progenitor; hESC, human ESC; mESC, mouse ESC; SP, side-population ability; Rho, rhodamine 123; Lin, lineage-negative, Lin; SCA1, stem-cell antigen 1; FLT3, fms-related tyrosine kinase 3; N-cad, N-cadherin; SCF, membrane-bound stem-cell factor also known as KIT ligand; TIE2, tyrosine kinase receptor 2; Atp5b, ATP synthase chain; Eno 1, Enolase 1; Tpi 1, Triosephosphate isomerase, Stip 1, stress-induced-phosphoprotein 1; Prdx 1, Peroxiredoxin 1; Cct 2, chaperonin-containing TCP1-subunit 2; Hspa5, 78-kDa glucose-regulated protein precursor; Hspa8; subunit and Heat shock cognate 71-kDa protein, an unclassifie protein; Tpt 1, Translationally controlled tumor protein or TCTP; Oct4, POU domain-containing transcription factor encoded by Pou5f1 gene; FRAP1, Mammalian target of rapamycin mTOR, FKBP12-rapamycin complex-associated protein, SUMO1, Small ubiquitin-related modifier 1, Sentrin, Ubiquitin-like protein SMT3C, GAP-modifying protein 1, UBL1, PIC1, GMP1; MCL1, nduced myeloid leukemia cell differentiation protein Mcl-1, Bcl-2-related protein EAT/mcl1; SLK, Proto-oncogene tyrosine-protein kinase Fyn; SYK, spleen tyrosine kinase; PTEN, phosphatase and tensin homologue deleted from chromosome 10, DDX3X, DEAD (Asp-Glu-Ala-Asp) box polypeptide 3X; SOX9, SRY (sex determining region Y)-box 9; SOX2, SRY (sex determining region Y)-box 2; BMP4, Bone morphogenic protein 4, AFP, Alpha-fetoprotein; TfR, Transferrin Receptor, CD71.

1. Hoffrogge R, Mikkat S, Scharf C et al. (2006) 2-DE proteome analysis of a proliferating and differentiating human neuronal stem cell line. Proteomics 6, pp.1833-1847

2. Elliott ST, Crider DG, Garnham CP et al. (2004) Two-dimensional gel electrophoresis database of murine r1 embryonic stem cells. Proteomics 4, pp.3813-3832.

3. Nagano K, Taoka M, Yamauchi Y et al. (2005) Large-scale identification of proteins expressed in mouse embryonic stem cells. Proteomics 5, pp.1346–1361.

4. van Hoof D, Passier R, Ward-Van Oostwaard D et al. (2006) A quest for human and mouse embryonic stem cell-specific proteins. Mol Cell Proteomics 5, pp.1261-1273.

5. Baharvand H, Hajheidari M, Ashtiani SK et al. (2006) Proteomic signature of human embryonic stem cells. Proteomics 6, pp.3544-3549.

6. Feldmann RE Jr, Bieback K, Maurer MH et al. (2005) Stem cell proteomes: A profile of human mesenchymal stem cells derived from umbilical cord blood. Electrophoresis 26, pp.2749-2758.

7. Maurer MH, Feldmann RE Jr, Futterer CD et al. (2003) The proteome of neural stem cells from adult rat hip-

8. DeLany JP, Floyd ZE, Zvonic S et al. (2005) Proteomic analysis of primary cultures of human adipose-derived

9. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B. and Ideker, T. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome

stem cells: Modulation by adipogenesis. Mol Cell Proteomics 4, pp.731-740.

Res 13, pp.2498-2504.

10. Vailaya, A., Bluvas, P., Kincaid, R., Kuchinsky, A., Creech, M. and Adler, A. (2005) An architecture for biological information extraction and representation. Bioinformatics, 21, pp.430-438. 11. Natalia B. Ivanova, John T. Dimos, Christoph Schaniel, Jason A. Hackney, Kateri A. Moore, Ihor R. Lemischka

(2002) A Stem Cell Molecular Signature. Science 298, pp.601-604.

12. Anne Wilson and Andreas Trumpp (2006) Bone-marrow haematopoieticstem-cell niches. Nature Rev Immun