


Backpropagation and chain rule

Chain rule is just simple math:

$$\frac{\partial L}{\partial x} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial x}$$


Backprop is just way to use it in NN training.



Backpropagation example

$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

A magenta arrow points from this box to the z input node in the computational graph above.

Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2$, $y = 5$, $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$, $\frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

$$\frac{\partial f}{\partial y}$$

Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Backpropagation example


$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

$$\frac{\partial f}{\partial x}$$

Practice time: interactive playground

DATA


Which dataset do you want to use?



Ratio of training to test data: 50%



Noise: 0



Batch size: 10



REGENERATE

FEATURES

Which properties do you want to feed in?



+ - 1 HIDDEN LAYER

+ -


2 neurons

This is the output from one **neuron**.
Hover to see it larger.

OUTPUT

Test loss 0.208

Training loss 0.207



Colors shows data, neuron and weight values.



☐ Show test data ☐ Discretize output


<https://playground.tensorflow.org/>

Backpropagation and chain rule

Chain rule is just simple math:


$$\frac{\partial L}{\partial x} = \frac{\partial L}{\partial z} \frac{\partial z}{\partial x}$$

Backprop is just way to use it in NN training.




Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$$\begin{array}{lcl}
 f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\
 f_a(x) = ax & \rightarrow & \frac{df}{dx} = a
 \end{array}
 \quad \Bigg| \quad
 \begin{array}{lcl}
 f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\
 f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1
 \end{array}$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$




$$\begin{array}{lcl} f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\ f_a(x) = ax & \rightarrow & \frac{df}{dx} = a \end{array}$$

$$\begin{array}{lcl} f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\ f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1 \end{array}$$

Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		<div style="border: 1px solid red; padding: 5px;">$f_c(x) = c + x$</div>	\rightarrow	<div style="border: 1px solid red; padding: 5px;">$\frac{df}{dx} = 1$</div>

Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$f(x) = e^x$	\rightarrow	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	\rightarrow	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	\rightarrow	$\frac{df}{dx} = a$		$f_c(x) = c + x$	\rightarrow	$\frac{df}{dx} = 1$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$$\boxed{f(x) = e^x \rightarrow \frac{df}{dx} = e^x}$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

$$f_c(x) = c + x$$

\rightarrow


$$\frac{df}{dx} = -1/x^2$$

\rightarrow

$$\frac{df}{dx} = 1$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$

$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

$$f_c(x) = c + x$$

→


$$\frac{df}{dx} = -1/x^2$$

→

$$\frac{df}{dx} = 1$$

Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$$\begin{array}{lcl}
 f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\
 f_a(x) = ax & \rightarrow & \frac{df}{dx} = a
 \end{array}
 \quad \Bigg| \quad
 \begin{array}{lcl}
 f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\
 f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1
 \end{array}$$

Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$



$$\begin{array}{lcl}
 f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\
 f_a(x) = ax & \rightarrow & \frac{df}{dx} = a
 \end{array}
 \quad \Bigg| \quad
 \begin{array}{lcl}
 f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\
 f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1
 \end{array}$$

Backpropagation example


Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



$$\begin{array}{lcl}
 f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\
 f_a(x) = ax & \rightarrow & \frac{df}{dx} = a
 \end{array}
 \quad \Bigg| \quad
 \begin{array}{lcl}
 f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\
 f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1
 \end{array}$$

Backpropagation example

Another example: $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$



[local gradient] x [its gradient]

$$x_0: [2] \times [0.2] = 0.4$$

$$w_0: [-1] \times [0.2] = -0.2$$


$$\begin{array}{lcl} f(x) = e^x & \rightarrow & \frac{df}{dx} = e^x \\ f_a(x) = ax & \rightarrow & \frac{df}{dx} = a \end{array} \quad \Bigg| \quad \begin{array}{lcl} f(x) = \frac{1}{x} & \rightarrow & \frac{df}{dx} = -1/x^2 \\ f_c(x) = c + x & \rightarrow & \frac{df}{dx} = 1 \end{array}$$

Backpropagation example

$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

$$\sigma(x) = \frac{1}{1 + e^{-x}} \quad \text{sigmoid function}$$



$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left(\frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left(\frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$



Gradient optimization

Stochastic gradient descent (and variations)
is used to optimize NN parameters.

$$x_{t+1} = x_t - \text{learning rate} \cdot dx$$




Once more: nonlinearities

$$f(a) = \frac{1}{1 + e^a}$$


$$f(a) = \tanh(a)$$

$$f(a) = \max(0, a)$$

$$f(a) = \log(1 + e^a)$$



Activation functions



Sigmoid


$$f(a) = \frac{1}{1 + e^a}$$

- Squashes numbers to range [0,1]
- Historically popular since they have nice interpretation as a saturating “firing rate” of a neuron

3 problems:

1. Saturated neurons “kill” the gradients
2. Sigmoid outputs are not zero-centered
3. $\exp()$ is a bit compute expensive

Activation functions




tanh(x)

- Squashes numbers to range [-1,1]
- zero centered (nice)
- still kills gradients when saturated :(

$$f(a) = \tanh(a)$$

Activation functions



ReLU


(Rectified Linear Unit)

$$f(a) = \max(0, a)$$

- Does not saturate (in +region)
- Very computationally efficient
- Converges much faster than sigmoid/tanh in practice (e.g. 6x)
- Not zero-centered output
- An annoyance:

hint: what is the gradient when $x < 0$?

Activation functions




- Does not saturate
- Computationally efficient
- Converges much faster than sigmoid/tanh in practice! (e.g. 6x)
- **will not “die”.**

Leaky ReLU

$$f(x) = \max(0.01x, x)$$

Activation functions

- Does not saturate
- Computationally efficient
- Converges much faster than sigmoid/tanh in practice! (e.g. 6x)
- **will not “die”.**



Leaky ReLU


$$f(x) = \max(0.01x, x)$$

Parametric Rectifier (PReLU)

$$f(x) = \max(\alpha x, x)$$

backprop into α
(parameter)

Exponential Linear Units (ELU)



$$f(x) = \begin{cases} x & \text{if } x > 0 \\ \alpha (\exp(x) - 1) & \text{if } x \leq 0 \end{cases}$$

- All benefits of ReLU
- Does not die
- Closer to zero mean outputs
- Computation requires $\exp()$

Activation functions: sum up

- Use **ReLU** as baseline approach
- Be careful with the learning rates
- Try out **Leaky ReLU** or **ELU**
- Try out **tanh** but do not expect much from it
- Do not use **Sigmoid**