

# **Fundamentals of Artificial Intelligence and Knowledge Representation (Module 3)**

Last update: 13 October 2023

Academic Year 2023 – 2024  
Alma Mater Studiorum · University of Bologna

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Uncertainty . . . . .	1
1.1.1	Handling uncertainty . . . . .	1
1.1.2	Probability . . . . .	1

# 1 Introduction

## 1.1 Uncertainty

**Uncertainty** A task is uncertain if we have:

Uncertainty

- Partial observations
- Noisy or wrong information
- Uncertain action outcomes
- Complex models

A purely logic approach leads to:

- Risks falsehood: unreasonable conclusion when applied in practice.
- Weak decisions: too many conditions required to make a conclusion.

### 1.1.1 Handling uncertainty

**Default/nonmonotonic logic** Works on assumptions. An assumption can be contradicted by an evidence.

Default/nonmonotonic logic

**Rule-based systems with fudge factors** Formulated as premise  $\rightarrow_{\text{prob.}}$  effect. Have the following issues:

Rule-based systems with fudge factors

- Locality: how can the probability account all the evidence.
- Combination: chaining of unrelated concepts.

**Probability** Assign a probability given the available known evidence.

Probability

Note: fuzzy logic handles the degree of truth and not the uncertainty.

**Decision theory** Defined as:

Decision theory

Decision theory = Utility theory + Probability theory

where the utility theory depends on one's preferences.

### 1.1.2 Probability

**Sample space** Set  $\Omega$  of all possible worlds.

Sample space

**Event** Subset  $A \subseteq \Omega$ .

Event

**Sample point/Possible world/Atomic event** Element  $\omega \in \Omega$ .

Sample point

**Probability space** A probability space/model is a function  $\mathcal{P}(\cdot) : \Omega \rightarrow [0, 1]$  assigned to a sample space such that:

Probability space

- $0 \leq \mathcal{P}(\omega) \leq 1$
- $\sum_{\omega \in \Omega} \mathcal{P}(\omega) = 1$

- $\mathcal{P}(A) = \sum_{\omega \in A} \mathcal{P}(\omega)$

**Random variable** A function from an event to some range (e.g. reals, booleans, ...).

Random variable

**Probability distribution** For any random variable  $X$ :

Probability distribution

$$\mathcal{P}(X = x_i) = \sum_{\omega \text{ st } X(\omega) = x_i} \mathcal{P}(\omega)$$

**Proposition** Event where a random variable has a certain value.

Proposition

$$a = \{\omega \mid A(\omega) = \text{true}\}$$

$$\neg a = \{\omega \mid A(\omega) = \text{false}\}$$

$$(\text{Weather} = \text{rain}) = \{\omega \mid B(\omega) = \text{rain}\}$$

**Prior probability** Prior/unconditional probability of a proposition based on known evidence.

Prior probability

**Probability distribution (all)** Gives all the probabilities of a random variable.

Probability distribution (all)

$$\mathbf{P}(A) = \langle \mathcal{P}(A = a_1), \dots, \mathcal{P}(A = a_n) \rangle$$

**Joint probability distribution** The joint probability distribution of a set of random variables gives the probability of all the different combinations of their atomic events.

Joint probability distribution

Note: Every question on a domain can, in theory, be answered using the joint distribution. In practice, it is hard to apply.

**Example.**  $\mathbf{P}(\text{Weather}, \text{Cavity}) =$

	Weather=sunny	Weather=rain	Weather=cloudy	Weather=snow
Cavity=true	0.144	0.02	0.016	0.02
Cavity=false	0.576	0.08	0.064	0.08

**Probability density function** The probability density function (PDF) of a random variable  $X$  is a function  $p : \mathbb{R} \rightarrow \mathbb{R}$  such that:

Probability density function

$$\int_{\mathcal{T}_X} p(x) dx = 1$$

**Uniform distribution**

Uniform distribution

$$p(x) = \text{Unif}[a, b](x) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

**Gaussian (normal) distribution**

Gaussian (normal) distribution

$$\mathcal{N}(\mu, \sigma^2) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

$\mathcal{N}(0, 1)$  is the standard gaussian.

**Conditional probability** Probability of a prior knowledge with new evidence:

Conditional probability

$$\mathcal{P}(a|b) = \frac{\mathcal{P}(a \wedge b)}{\mathcal{P}(b)}$$