

## Lecture Outline

This lecture will cover the roles of insulin and glucagon in normal and pathophysiological states, especially diabetes. It will focus on pharmacological interventions for both of these pathologies. This lecture will cover the following four topics.

- Physiological regulation of blood glucose
  - Insulin Signaling
  - Glucagon Signaling
- Pathophysiology related to glucose control
  - Type I Diabetes
  - Type II Diabetes
- Common Pharmacological Interventions for Diabetes

## Diabetes

### Diabetes in the United States.




Figure 1: County Level Prevalence in Type II Diabetes

As of 2011, diabetes affects 25.8 million people or 8.3% of the population. This includes approximately 7 million undiagnosed individuals. In addition to this, 79 million Americans over the age of 20 are estimated to be pre-diabetic. As of 2007, diabetes is estimated to cost \$116 billion in direct and \$58 billion in indirect costs [CDC2011].

Diabetes is especially prevalent in this region. As shown in the map, there is a higher percentage of diabetics in the mid and deep south regions of the United States.

## Complications of Diabetes

The major complications of diabetes include retinopathy, nephropathy, neuropathy and cardiovascular disease including heart attacks and stroke. These events are primarily due to excessive glycation of extracellular and intracellular targets preventing their function.

## The Discovery of Insulin




CHART III.—*Effect of one injection of extract on Blood Sugar (mg. per c.c.=tenth per cent)*

Figure 2: Effects of a dog pancreas isolate on blood glucose levels in a 14 year old diabetic child (F.G. Banting et al. 1922).

Insulin was discovered by Frederick Banting and his colleagues at the University of Toronto in 1921. They performed experiments in which they injected extracts from pancreas fractions into dogs which had their pancreas' surgically removed. They showed that a secreted substance from the pancreas lowered blood glucose in these dogs (Frederick G. Banting and Best 1922). They were then able to confirm that this treatment was also effective in children with diabetes (F.G.

Banting et al. 1922). This work led to Banting and John Macleod winning the Nobel Prize in Medicine and Physiology in 1923.

## **Normal regulation of circulating glucose**

Glucose is maintained in a very narrow range, between 4.4 to 6.1 mmol/L. These levels need to be re-established after changes in feeding status, or energy utilization. In general, when glucose levels decrease, glucagon is released from alpha cells of the pancreas to promote glucose production, either from glycogen breakdown or gluconeogenesis. Alternately, after a meal when glucose levels increase, insulin is secreted from beta cells of the pancreas causing glucose levels to decrease.

## **Mechanisms of glucose control**

For the purposes of the acute maintenance of glucose homeostasis, four organs are the most important; the pancreas, liver, muscle and adipose tissue. The pancreas senses changes in glucose levels and responds by releasing either glucagon or insulin.

### **Glucose Uptake**

Glucose is taken up down a concentration gradient into most tissues including liver, pancreas, kidneys and the brain. However, for glucose to enter into muscle and fat tissue, insulin is required.

### **Glycolysis**

The major route for energy production is glycolysis, the TCA cycle and oxidative respiration in the mitochondria. This pathway converts glucose and other carbohydrates first into pyruvate, then into components used in the electron transport chain to generate ATP in the mitochondria.

Glycolysis is regulated both by allosteric activators and inhibitors, as well as by protein phosphorylation in response to extracellular and intracellular signals.

### **Gluconeogenesis**

Gluconeogenesis is the generation of glucose from non-carbohydrate precursor molecules. These typically include amino acids, lactate and the products of fatty acid oxidation. The vast majority of gluconeogenesis occurs in the liver. This

process is similar to reverse glycolysis though in several cases different enzymes are used.

The rate limiting enzymes in gluconeogenesis are Phosphoenolpyruvate Carboxykinase, Fructose-1,6-bisphosphatase and Glucose-6-Phosphatase. These enzymes are under both transcriptional and post-translational control as described below.

### **Glycogenesis**

As an alternative to glycolysis, glucose can also be stored in esterified form as glycogen. To form glycogen, glucose must first be converted through Glucose-1-Phosphate into UDP-Glucose. This activated form of glucose is then added onto existing glycogen chains through the activity of an enzyme named Glycogen Synthase. In addition to being regulated by protein phosphorylation and sub-cellular location, Glycogen synthase is also allosterically activated by Glucose-6-Phosphate, promoting increased glycogen synthesis when glucose levels in the cell are high.

### **Glycogenolysis**

To liberate glucose from stored glycogen, glycogen phosphorylase is activated. This enzyme hydrolyses glycogen releasing glucose-1-phosphate, which can then be dephosphorylated into glucose for glycolysis or release into the blood stream. In addition to post-translational modifications and recruitment to the glycogen pellet by accessory proteins, glycogen phosphorylase is allosterically activated by energy stress such as increases in AMP, or negatively by increases in Glucose-6-Phosphate levels.

## **Endocrine Control of Blood Glucose**

In order to balance the energy requirements of all tissues, blood glucose is primarily controlled via endocrine and neuroendocrine mechanisms. The primary mediators are insulin and glucagon which are secreted from the pancreas during times of hyper and hypoglycemia respectively.

### **Insulin Functions**

When glucose levels are raised, such as after a meal, insulin has four main functions, all of which serve to reduce blood glucose levels:

1. Promotes the uptake of glucose from the blood into muscle and adipose tissue.

2. Enhances the synthesis of glycogen and triglycerides in liver, adipose and muscle.
3. Insulin inhibits gluconeogenesis, or the production of glucose from non-glucose precursors such as amino acids and lipids.
4. Promote the breakdown of glucose via glycolysis.

### Insulin Secretion




Figure 3: Regulation of insulin secretion by beta cells in pancreatic islets. (Dean and McEntyre 2004).

Beta cells in the pancreas generate insulin and store it in pre-formed secretory granules. After glucose mediated depolarization of beta cells in response to insulin, these secretory granules are exocytosed and their contentes are released into the blood.

## Insulin Signal Transduction

Insulin functions by binding to and activating a receptor tyrosine kinase. This receptor transautophosphorylates itself generating binding sites for phosphotyrosine binding proteins known as insulin receptor substrates.

These proteins are also phosphorylated by the insulin receptor, which creates binding sites for a phosphatidylinositol-3-kinase (PI3K). This kinase generates the key second messenger in insulin signaling, phosphatidylinositol-(3,4,5)-triphosphate (PIP3). Most known functions of insulin are blocked when PI3K is inhibited (Kanai et al. 1993).

Once PIP3 is generated by insulin stimulation, it can diffuse along the internal membranes of the cells. This lipid second messenger recruits two important protein kinases, Akt (also referred to as PKB) and PDK1. Both of these proteins have domains called pleckstrin homology domains which recruit the kinases together to the plasma membrane. Once there, PDK1 and another protein kinase called mTORC2 are able to phosphorylate and activate Akt. Once activated, Akt is the most important protein kinase in mediating insulin function.

## Insulin Stimulated Glucose Uptake

In fat and muscle tissue, insulin promotes the movement of a facilitative glucose transporter named GLUT4. Normally GLUT4 resides in intracellular compartments, but in response to insulin vesicles from these compartments fuse with the plasma membrane, inserting GLUT4 into the extracellular surface. This allows for glucose to enter fat and muscle cells (Leto and Saltiel 2012).

In both fat and muscle, the PI3K/Akt dependent signaling pathways are absolutely required for insulin stimulated glucose uptake (Okada et al. 1994). The major targets of Akt in this signaling pathway are AS160 and RGC1/2, two proteins which regulate the activity of small GTPases involved in GLUT4 translocation (Leto and Saltiel 2012). These mechanisms are not yet fully understood.

## How Does Insulin Stimulate Glycogenesis

Allosterically, the other functions of insulin, including promoting glucose uptake and glycolysis will provide more UDP-Glucose for glycogen synthase, and will also allosterically activate the enzyme by generating large amounts of Glucose-6-Phosphate.

In addition to these allosteric activators, both glycogen synthase and glycogen phosphorylase are regulated by protein phosphorylation. In the case of glycogen synthase, the phosphorylated form is relatively inactive, and is resistant

to allosteric activation by glucose-6-phosphate (Friedman and Lerner 1963). Glycogen synthase is phosphorylated by several protein kinases including AMPK and GSK-3 (Parker et al. 1982). In addition to inactivating the upstream kinases, insulin also activates a protein phosphatase, which removes the phosphate groups.




Figure 4: Regulation of glycogen synthesis by insulin and protein phosphorylation.

In a co-ordinated way, protein phosphorylation activates glycogen phosphorylase (Krebs and Fischer 1964). This means that when these enzymes are phosphorylated, the balance tips towards glycogenolysis, and when they are dephosphorylated glycogen is synthesized.

## How Does Insulin Suppress Gluconeogenesis

The activation of glucose uptake and glycolysis leads to increased levels of several glycolytic intermediates which themselves will regulate gluconeogenesis. The most important of these is Fructose-2,6-bisphosphate which is raised during glycolysis and inhibits FBPase, one of the key rate limiting steps in gluconeogenesis.

In addition to these effects, both G6Pase and PEPCK, two other rate limiting enzymes are regulated transcriptionally. Akt phosphorylates and inactivates the transcription factor FOXO which would normally drive the expression of these enzymes. Therefore when insulin activate the PI3K/Akt cascade, FOXO mediated transcription of G6Pase and PEPCK is decreased and the levels of these enzymes are reduced, decreasing gluconeogenesis.

## How Is Blood Glucose Increased?

### Physiological effects of glucagon

When glucose levels are low, glucagon is released from alpha cells in the pancreas. This promotes the breakdown of glycogen stores in liver and muscle, and the generation of glucose from gluconeogenic precursors. Glucagon receptors exist mainly in the liver, so glucagon does not exert its catabolic effects on either adipose or muscle tissue [Drucker2003].

### Regulation of glucagon release

The mechanisms which underlie hypoglycemia induced glucagon are incompletely understood. What is clear however, is that when blood glucose levels decrease, glucagon is released from the alpha cells of the pancreas into the portal vein.

### Glucagon signaling

Adrenergic-receptor coupled mediated cAMP synthesis was the first example of a hormonal second messenger. Earl Sutherland was interested in the regulation of glycogenolysis and he noticed that if he added adrenaline to intact cells, he could accelerate glycogen breakdown, but if he added it to lysed cells he could not. In his key experiment he treated one set of livers with adrenaline, then lysed them. He then added that lysate to a second set of livers which had already been broken. He found that there was an internal factor (later identified by cAMP) in the stimulated tissues, that could accelerate glycogenolysis in the other tissues (Rall, Sutherland, and Wosilait 1956). For this work, Sutherland won the Nobel Prize in Medicine and Physiology in 1971.

In metabolism, the main effector of cAMP in cells is Protein Kinase A (PKA). This protein kinase is allosterically activated by cAMP and phosphorylates



a wide variety of important metabolic substrates. The identification of PKA and its role in carbohydrate homeostasis led to Fisher and Krebs winning the Nobel Prize in Medicine and Physiology in 1992. The major role of glucagon is to stimulate glucose release, both by mobilizing glycogen stores and inducing gluconeogenesis.

### **Effects of glucagon on glycolysis.**

As described above, glucagon stimulates the breakdown of glycogen. This proceeds via protein phosphorylation of both glycogen phosphorylase (which activates the enzyme) and glycogen synthase (which inactivates the enzyme). In combination, this leads to a breakdown of glycogen into glucose.

PKA is the primary mediator of the activation of glycogen phosphorylase. Once activated by adrenergic signaling, PKA phosphorylates and activates glycogen phosphorylase kinase. This kinase in turn, phosphorylates and activates glycogen phosphorylase (Krebs and Fischer 1956). PKA also directly phosphorylates glycogen synthase, which in concert with the activation of the other glycogen synthase kinases (notably GSK3 and AMPK) leads to increased phosphorylation and inactivation of glycogen synthase.

In addition to the activation of these protein kinases, there is a reduction of glycogen associated protein phosphatase activity. As a balance, this leads to more highly phosphorylated and therefore more glycogenolytic activities.

### **How does glucagon promote gluconeogenesis?**

In addition to the decreased flux of glycolytic intermediates which allosterically activate gluconeogenesis, there are both post-translational and transcriptional mechanisms by which adrenergic signaling promotes gluconeogenesis.

Post-translationally, the best studied route by which PKA activates gluconeogenesis is through inactivation of phosphofructokinase-2. PFK-2 normally generates the carbohydrate Fructose-2,6-bisphosphate which is a positive regulator of glycolysis and a negative regulator of gluconeogenesis. The alleviation of this inhibition allows for promotion of the gluconeogenic metabolism.

Transcriptionally, the transcription factor CREB is phosphorylated by PKA where it plays a role in transcriptionally activating the rate limiting gluconeogenic enzymes PEPCK, FBPase and G6Pase.

## **Exercise and Sympathetic Tone**

### **Adrenaline and the Sympathetic Nervous System**

Since glucagon works primarily on liver tissue, different hormonal messengers function to stimulate catabolism in muscle and fat tissue. The activation of PKA by GPCR and cAMP signaling pathways leads to glycogen breakdown in muscle via similar mechanisms as those in liver. Adrenaline also leads to enhanced lipid and glucose oxidation in muscle primarily as an energy source.

In adipose tissue, these pathways induce lipolysis, via phosphorylation and activation of Hormone Sensitive Lipase (HSL), Perilipin and Adipocyte Triglyceride Lipase (ATGL). These proteins function to mobilize triglycerides into free fatty acids for use in other tissues, especially muscle. For more information on the regulation of lipolysis, see (Young and Zechner 2013). At an acute level, these do not contribute much to glucose homeostasis.

### **The AMPK Activated Protein Kinase**

Intracellularly, when levels of ATP decrease and correspondingly, the levels of AMP rise, a protein kinase called the AMP Activated Protein Kinase is turned on. AMPK is like a cellular fuel gauge, so that when ATP levels are decreased, energy producing pathways are activated and energy using pathways are reduced (Hardie, Ross, and Hawley 2012).

## **Pathophysiology related to glucose control**

### **Type I Diabetes Mellitus**

Type I Diabetes is typically caused by autoimmune destruction of pancreatic beta cells. Without these cells, the pancreas is unable to produce insulin and without careful monitoring and exogenous insulin, blood glucose levels will rise.

### **Insulin Resistance and Type II Diabetes Mellitus**

Type II diabetes occurs as a result of a multi-step process starting with negative feedback loops on insulin signaling. As more nutrients are stored, metabolic tissues become resistant to the effects of insulin, likely as a way to protect against excessive lipid storage.

As tissues become more insulin resistant, more insulin must be secreted by the pancreas to maintain normoglycemia. If insulin resistance proceeds, more and more insulin will need to be produced and secreted by beta cells. Eventually the

beta cells will be unable to keep up with this demand and glucose levels will rise as the amount of endogenous or exogenous insulin is less and less effective.

### **Why Insulin Resistance**

Insulin resistance is a standard negative feedback loop, common in most endocrine pathways. Normally, it serves to ensure that anabolic pathways are not over-active. Evolutionarily, it has been hypothesized that our carbohydrate and lipid homeostatic mechanisms are wired to promote lipid storage since during the majority of human existence, excessive nutrient storage was not problematic (Neel 1999).

There is both an adaptive and a genetic component to the extent of insulin resistance. Some obese individuals are able to maintain insulin sensitivity to a greater extent, while others progress from moderate insulin resistance to pancreatic failure rather quickly. These variations, which are largely consistent between ethnic groups underlie how rapidly and severely individuals may progress towards Type II diabetes.

### **Molecular mechanisms of insulin resistance**

Obese individuals respond poorly to insulin, including less insulin stimulated glucose uptake, and less insulin suppression of glucose production in the liver. Biochemical measurements were done to dissect these processes. These studies showed the activity of the PI3K signaling pathways were reduced in obese states (L. J. Goodyear et al. 1995)(Saad et al. 1992). This is thought to occur via serine phosphorylation of insulin receptor substrate proteins (Takayama et al. 1984)(White, Takayama, and Kahn 1985). This prevents these substrates from being phosphorylated by the insulin receptor tyrosine kinase and prevents the activation of the PI3K signal transduction cascade.

### **Mechanisms of Insulin Resistance**

#### **Nutrient overload induced insulin resistance**

#### **Lipid mediated insulin resistance**

#### **Inflammation associated insulin resistance**

## **How Does Insulin Resistance Lead to Hyperglycemia and Diabetes?**

### **Adaptations to Insulin Resistance**

- Hyperinsulinemia -> Pancreatic Failure
- Regulation of food intake

## **Common Pharmacological Interventions for Diabetes**

- Primary intervention is diet and exercise alteration

### **Treatments for Type I Diabetes**

#### **Insulin Administration**

The amino acid sequence of insulin was determined by Frederick Sanger, which led to him winning the Nobel Prize in Chemistry in 1958 [Sanger1951]. This eventually allowed for recombinant production and manipulation of insulin, rather than using purified porcine, ovine or bovine insulin.

#### **Short and Long Acting Insulin**

#### **Insulin Pumps**

#### **Islet Transplantation**

### **Treatments for Type II Diabetes**

#### **Excretion of circulating glucose**

#### **Storage of circulating glucose**

#### **Insulin administration and insulin secretagogues**

- Sulfonylureas

#### **Drugs that promote insulin sensitivity**

- Thiazolidinediones
- Anti-inflammatories

## Glucose Utilization

One class of drugs called biguanides is a common therapeutic for Type II Diabetes. These drugs, such as phenformin and metformin (marketed as Glucophage) function by disrupting mitochondrial function. These leads to energy stress and activation of AMPK due to the increases in intracellular AMP (Zhou et al. 2001). The activation of AMPK leads to activation of energy producing pathways, including glucose uptake and oxidation and lipid oxidation. The effects of this are to reduce circulating glucose, while also burning fat.




Figure 5: Mechanism of Action of Biguanides.

## Further Reading

Banting, F. G., C. H. Best, J. B. Collip, W. R. Campbell, and a Fletcher. 1922. "Pancreatic Extracts in the Treatment of Diabetes Mellitus." *Canadian Medical Association Journal* 12 (3) (mar): 141–6. <http://www.ncbi.nlm.nih.gov/pubmed/17580419>.

- Banting, Frederick G., and Charles H. Best. 1922. "The Internal Secretion of the Pancreas." *Journal of Laboratory and Clinical Medicine* 7: 251–266.
- Dean, Laura, and Jo McEntyre. 2004. *The Genetic Landscape of Diabetes*. Bethesda, MD: National Center for Biotechnology Information (US). <http://www.ncbi.nlm.nih.gov/books/NBK1671>.
- Friedman, D. L., and J. Lerner. 1963. "STUDIES ON UDPG-ALPHA-GLUCAN TRANSGLUCOSYLASE. III. INTERCONVERSION OF TWO FORMS OF MUSCLE UDPG-ALPHA-GLUCAN TRANSGLUCOSYLASE BY A PHOSPHORYLATION-DEPHOSPHORYLATION REACTION SEQUENCE." *Biochemistry* 2: 669–75. <http://www.ncbi.nlm.nih.gov/pubmed/14075096>.
- Goodyear, L. J., F. Giorgino, L. a Sherman, J. Carey, R. J. Smith, and G. L. Dohm. 1995. "Insulin Receptor Phosphorylation, Insulin Receptor Substrate-1 Phosphorylation, and Phosphatidylinositol 3-kinase Activity Are Decreased in Intact Skeletal Muscle Strips from Obese Subjects." *The Journal of Clinical Investigation* 95 (5) (may): 2195–204. doi:10.1172/JCI117909. <http://dx.doi.org/10.1172/JCI117909>.
- Hardie, D. Grahame, Fiona a Ross, and Simon a Hawley. 2012. "AMPK: a Nutrient and Energy Sensor That Maintains Energy Homeostasis." *Nature Reviews. Molecular Cell Biology* 13 (4) (jan): 251–62. doi:10.1038/nrm3311. <http://dx.doi.org/10.1038/nrm3311>.
- Kanai, F., K. Ito, M. Todaka, H. Hayashi, S. Kamohara, K. Ishii, T. Okada, O. Hazeki, M. Ui, and Y. Ebina. 1993. "Insulin-stimulated GLUT4 Translocation Is Relevant to the Phosphorylation of IRS-1 and the Activity of PI3-kinase." *Biochemical and Biophysical Research Communications* 195 (2) (sep): 762–8. doi:10.1006/bbrc.1993.2111. <http://www.ncbi.nlm.nih.gov/pubmed/8396927>.
- Krebs, E. G., and E. H. Fischer. 1956. "The Phosphorylase B to a Converting Enzyme of Rabbit Skeletal Muscle." *Biochimica Et Biophysica Acta* 20 (1) (apr): 150–7. <http://www.ncbi.nlm.nih.gov/pubmed/13315361>.
- . 1964. "Phosphorylase and Related Enzymes of Glycogen Metabolism." *Vitamins and Hormones* 22 (jan): 399–410. doi:10.1016/S0083-6729(08)60345-3. <http://www.ncbi.nlm.nih.gov/pubmed/14284112>.
- Leto, Dara, and Alan R. Saltiel. 2012. "Regulation of Glucose Transport by Insulin: Traffic Control of GLUT4." *Nature Reviews. Molecular Cell Biology* 13 (6) (jun): 383–96. doi:10.1038/nrm3351. <http://www.ncbi.nlm.nih.gov/pubmed/22617471>.
- Neel, James V. 1999. "The 'Thrifty Genotype' in 1998." *Nutrition Reviews* 57 (5) (apr): 2–9. doi:10.1111/j.1753-4887.1999.tb01782.x. <http://dx.doi.org/10.1111/j.1753-4887.1999.tb01782.x>.
- Okada, T., Y. Kawano, T. Sakakibara, O. Hazeki, and M. Ui. 1994. "Essential Role of Phosphatidylinositol 3-kinase in Insulin-induced Glucose Transport and Antilipolysis in Rat Adipocytes. Studies with a Selective Inhibitor Wortmannin."

*The Journal of Biological Chemistry* 269 (5) (feb): 3568–73. <http://www.ncbi.nlm.nih.gov/pubmed/8106400>.

Parker, Peter J., Noor Embi, F. Barry Caudwell, and Philip Cohen. 1982. “Glycogen Synthase from Rabbit Skeletal Muscle. State of Phosphorylation of the Seven Phosphoserine Residues in Vivo in the Presence and Absence of Adrenaline.” *European Journal of Biochemistry / FEBS* 124 (1) (may): 47–55. <http://www.ncbi.nlm.nih.gov/pubmed/6211353>.

Rall, T. W., E. W. Sutherland, and W. D. Wosilait. 1956. “The Relationship of Epinephrine and Glucagon to Liver Phosphorylase. III. Reactivation of Liver Phosphorylase in Slices and in Extracts.” *The Journal of Biological Chemistry* 218 (1) (jan): 483–95. <http://www.ncbi.nlm.nih.gov/pubmed/13278355>.

Saad, M. J., E. Araki, M. Miralpeix, P. L. Rothenberg, M. F. White, and C. Ronald Kahn. 1992. “Regulation of Insulin Receptor Substrate-1 in Liver and Muscle of Animal Models of Insulin Resistance.” *The Journal of Clinical Investigation* 90 (5) (nov): 1839–49. doi:10.1172/JCI116060. <http://dx.doi.org/10.1172/JCI116060>.

Takayama, S., M. F. White, V. Lauris, and C. Ronald Kahn. 1984. “Phorbol Esters Modulate Insulin Receptor Phosphorylation and Insulin Action in Cultured Hepatoma Cells.” *Proceedings of the National Academy of Sciences of the United States of America* 81 (24) (dec): 7797–801. <http://www.pubmed.org/6393128>.

White, M. F., S. Takayama, and C. Ronald Kahn. 1985. “Differences in the Sites of Phosphorylation of the Insulin Receptor in Vivo and in Vitro.” *The Journal of Biological Chemistry* 260 (16) (aug): 9470–8. <http://www/pubmed.org/3848433>.

Young, Stephen G., and Rudolf Zechner. 2013. “Biochemistry and Pathophysiology of Intravascular and Intracellular Lipolysis.” *Genes & Development* 27 (5) (mar): 459–84. doi:10.1101/gad.209296.112. <http://dx.doi.org/10.1101/gad.209296.112>.

Zhou, Gaochao, Robert Myers, Ying Li, Yuli Chen, Xiaolan Shen, Judy Fenyk-Melody, Margaret Wu, et al. 2001. “Role of AMP-activated Protein Kinase in Mechanism of Metformin Action.” *Journal of Clinical Investigation* 108 (8) (oct): 1167–1174. doi:10.1172/JCI200113505. <http://dx.doi.org/10.1172/JCI200113505> <http://www.jci.org/articles/view/13505>.