Outcome	Time	Studies (n)	Total Participants	:s								BMD	Change	95% CI	I Squared
Transwomen vs cisgender men femoral neck BMD	24m	3	258						_				0.02	[-0.12 ; 0.16]	0.96
Transwomen vs cisgender men femoral neck BMD	12m	1	35					+					-0.07	[-0.08; -0.06]	n/a
Transwomen vs cisgender men lumbar spine BMD	24m	4	327				_	-	-				-0.01	[-0.13; 0.10]	0.95
Transwomen vs cisgender men lumbar spine BMD	12m	1	35					-					-0.06	[-0.08; -0.04]	n/a
Transwomen vs cisgender men total femur BMD	24m	3	274					$\overline{}$					-0.08	[-0.23; 0.06]	0.97
Transwomen vs cisgender men total femur BMD	12m	1	35	- 23									-0.74	[-0.77; -0.71]	n/a
Transgender men vs cisgender women femoral neck BMD	24m	2	170					$+\cdot$					0.06	[-0.09; 0.20]	0.95
Transgender men vs cisgender women lumbar spine BMD	24m	2	170					-					-0.01	[-0.06; 0.03]	0.19
Transgender men vs cisgender women total femur BMD	24m	1	100										0.01	[-0.03; 0.05]	n/a
						-	ı	1	ı						
				-0.8	-0.6	-0.4	-0.2	0	0.2	0.4	0.6	8.0			