

Table S1: Data sources for each ecoregion, including literature references for mined data sources and the names of contributed data sets. Contributed data sets are described in more detail in Table S2. Also included are the species represented by each data set in each ecoregion.

Ecoregion	Data source	Species
Agulhas Bank	<i>Contributed:</i> South Africa Gansbaai	<i>Ecklonia maxima</i>
Aleutian Islands	<i>Contributed:</i> Estes NPRB	<i>Agarum clathrus</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Laminaria yezoensis</i> , <i>Laminaria longipes</i> , <i>Eualaria fistulosa</i> , <i>Cymathaere triplicata</i> , <i>Agarum clathratum</i>
Bassian	<i>Contributed:</i> Kelp Cover Australia Victoria, Kelp Cover Temperate Australia	<i>Undaria pinnatifida</i> , <i>Macrocystis pyrifera</i> , <i>Lessonia corrugata</i> , <i>Ecklonia radiata</i>
Beaufort Sea	<i>Contributed:</i> Konar Beaufort Sea	<i>Laminaria solidungula</i> , <i>Saccharina latissima</i> , <i>Alaria esculenta</i>
Cape Howe	<i>Contributed:</i> Kelp Cover Australia Victoria, Kelp Cover Temperate Australia	<i>Ecklonia radiata</i>
Celtic Seas	<i>Mined:</i> 1, 2 <i>Contributed:</i> The Brittany Rebet Monitoring, Northern Ireland Summary	<i>Undaria pinnatifida</i> , <i>Saccharina latissima</i> , <i>Laminaria ochroleuca</i> , <i>Laminaria hyperborea</i> , <i>Laminaria digitata</i> , <i>Alaria esculenta</i>
Central Chile	<i>Contributed:</i> Northern Chile Fondecyt	<i>Macrocystis pyrifera</i> , <i>Lessonia trabeculata</i>
Chiloense	<i>Contributed:</i> Southern Chile	<i>Macrocystis pyrifera</i>
Gulf of Alaska	<i>Mined:</i> 3 <i>Contributed:</i> Konar Gulf of Alaska	<i>Saccharina latissima</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Laminaria yezoensis</i> , <i>Laminaria longipes</i> , <i>Laminaria bongardiana</i> , <i>Eualaria fistulosa</i> , <i>Cymathaere triplicata</i> , <i>Costaria costata</i> , <i>Alaria marginata</i> , <i>Agarum clathratum</i>
Gulf of Maine	<i>Mined:</i> 4 <i>Contributed:</i> Steneck Gulf of Maine, Witman Gulf of Maine, Normandeau Associates	<i>Saccharina latissima</i> , <i>Laminaria digitata</i> , <i>Alaria esculenta</i> , <i>Agarum clathratum</i>
Gulf of St. Lawrence	<i>Mined:</i> 5, 6	<i>Saccharina latissima</i> , <i>Alaria esculenta</i> , <i>Agarum clathratum</i>
Houtman	<i>Mined:</i> 7,8,9 <i>Contributed:</i> Kelp Cover Temperate Australia, Australia Western Australia Wernberg	<i>Ecklonia radiata</i>
Humboldtian	<i>Mined:</i> 10 <i>Contributed:</i> Northern Chile Fondecyt	<i>Macrocystis pyrifera</i> , <i>Lessonia trabeculata</i>
Leeuwin	<i>Contributed:</i> Australia Western Australia Wernberg	<i>Ecklonia radiata</i>
Manning-Hawkesbury	<i>Mined:</i> 11,12,13	<i>Ecklonia radiata</i>
Namaqua	<i>Mined:</i> 14,15	<i>Laminaria pallida</i> , <i>Ecklonia maxima</i>
North American Pacific Fjordland	<i>Mined:</i> 16, 17 <i>Contributed:</i> Haida Gwaii	<i>Saccharina latissima</i> , <i>Saccharina groenlandica</i> , <i>Pterygophora californica</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Macrocystis pyrifera</i> , <i>Laminaria yezoensis</i> , <i>Laminaria setchellii</i> , <i>Laminaria longipes</i> , <i>Saccharina groenlandica</i> , <i>Eualaria fistulosa</i> , <i>Egregia menziesii</i> , <i>Cymathaere triplicata</i> , <i>Costaria costata</i> , <i>Alaria marginata</i> , <i>Agarum clathrus</i> , <i>Agarum clathratum</i>
North Sea	<i>Contributed:</i> Norway Inner Coast (monitoring), Norway Inner Cast (assessment), Norway Monitoring	<i>Saccharina latissima</i> , <i>Laminaria hyperborea</i> , <i>Laminaria digitata</i>
Northeastern New Zealand	<i>Contributed:</i> New Zealand Hakurai Gulf, Northern New Zealand	<i>Lessonia variegata</i> , <i>Ecklonia radiata</i>
Northern and Central California	<i>Mined:</i> 18,19, 20 <i>Contributed:</i> Edwards Baja, San Nicholas Island, Channel Islands National Park, PISCO Subtidal	<i>Undaria pinnatifida</i> , <i>Pterygophora californica</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Macrocystis pyrifera</i> , <i>Laminaria setchellii</i> , <i>Saccharina latissima</i> , <i>Laminaria farlowii</i> , <i>Ecklonia arborea</i> , <i>Egregia menziesii</i> , <i>Dictyonium californicum</i> , <i>Costaria costata</i> , <i>Alaria marginata</i> , <i>Agarum fimbriatum</i>
Oregon, Washington, Vancouver Is.	<i>Mined:</i> 21, 22, 23, 24, 25 <i>Contributed:</i> PISCO Oregon, Vancouver Island	<i>Undaria pinnatifida</i> , <i>Saccharina latissima</i> , <i>Pterygophora californica</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Macrocystis</i>

		<i>pyrifer</i> , <i>Lessoniopsis littoralis</i> , <i>Laminaria setchellii</i> , <i>Laminaria farlowii</i> , <i>Ecklonia arborea</i> , <i>Egregia menziesii</i> , <i>Costaria costata</i> , <i>Alaria marginata</i> , <i>Agarum fimbriatum</i> , <i>Saccharina groenlandica</i>
Scotian Shelf	<i>Mined</i> : 26, 27, 28, 29, 30, 31, 32, 33,34, 35, 36, 37, 38, 39, 40	<i>Saccharina latissima</i> , <i>Laminaria digitata</i> , <i>Agarum clathratum</i>
South Australian Gulfs	<i>Contributed</i> : Kelp Cover Temperate Australia, Australia Connell Canopy Loss	<i>Ecklonia radiata</i>
South European Atlantic Shelf	<i>Contributed</i> : The Brittany Rebet Monitoring	<i>Undaria pinnatifida</i> , <i>Saccharina latissima</i> , <i>Laminaria ochroleuca</i> , <i>Laminaria hyperborea</i>
Southern California Bight	<i>Contributed</i> : Edwards Baja, San Nicholas Island, Channel Islands National Park, PISCO Subtidal, Santa Barbara Coastal Long Term Ecological Research, Isla Natividad	<i>Undaria pinnatifida</i> , <i>Pterygophora californica</i> , <i>Pleurophycus gardneri</i> , <i>Nereocystis luetkeana</i> , <i>Macrocystis pyrifera</i> , <i>Laminaria setchellii</i> , <i>Saccharina latissima</i> , <i>Laminaria farlowii</i> , <i>Ecklonia arborea</i> , <i>Egregia menziesii</i> , <i>Dictyoneurum californicum</i> , <i>Costaria costata</i> , <i>Alaria marginata</i> , <i>Agarum fimbriatum</i>
Southern Norway	<i>Contributed</i> : Norway Inner Coast (monitoring), Norway Inner Coast (assessment), Norway Monitoring	<i>Saccharina latissima</i> , <i>Laminaria hyperborea</i> , <i>Laminaria digitata</i>
Western Bassian	<i>Contributed</i> : Kelp Cover Temperate Australia,	<i>Ecklonia radiata</i> , <i>Macrocystis pyrifera</i>

Table S2. Contributed data set information, including the corresponding ecoregion, and the years covered, number of sites, contact person, and associated publications.

Data Set Name	Ecoregion	Years	# Sites	Contact	Associated Publications
Vancouver Island	Oregon, Washington Vancouver Coast and Shelf	1987-2011	8	Jane Watson	44
San Nicolas Island	Northern California and Southern California Bight	1980-2011	7	Mike Kenner	45
Witman Gulf of Maine	Gulf of Maine	1979-2014	4	Jon Witman	46, 47
Konar Beaufort Sea	Beaufort Sea	2003-2007	4	Brenda Konar	
Channel Islands National Park	Northern California and Southern California Bight	1986-2011	16	David Kushner	48
South Africa Gansbaai	Agulhas Bank	1992-1994	2	Rob Anderson	
Central Chile Fondecyt # 11110351 & 1151094	Central Chile	2013-2013	5	Alejandro Perez Matus	49
PISCO Subtidal	Central California and Southern California Bight	1999-2012	146	Jenn Caselle Mark Carr	
Santa Barbara Coastal Long Term Ecological Research	Southern California Bight	2001-2014	8	Dan Reed	51
Kelp Cover Temperate Australia	Bassian, Cape Howe, Houtman, South Australian Gulfs, Western Bassian	1992-2012	187	Graham Edgar Neville Barrett	52, 53, 54
Estes NPRB	Aleutian Islands	1987-2010	147	Mike Kenner, Jim Estes	
Isla Natividad, Baja California, Mexico	Southern California Bight	2006-2014	5	Fiorenza Micheli, Jorge Torre (Comunidad y Biodiversidad, Asociacion Civil)	55, 56
New Zealand Hauraki Gulf	Northeastern New Zealand	2007-2011	1	Nick Shears	57
Northern New Zealand	Northeastern New Zealand	1999-2011	12	Nick Shears	58, 59
The Brittany Rebent Monitoring The European Water Framework Directive Monitoring	Celtic Seas, South European Atlantic Shelf	2004-2013	25	Sandrine Derrien- Courtel	60, 61, 62
Australia Western Australia Wernberg	Houtman and Leeuwin	2006-2011	6	Thomas Wernberg	63
PISCO Oregon	Oregon, Washington Vancouver Coast and Shelf	2001-2004	8	Mark Carr	
Norway Monitoring	North Sea and Southern Norway	1990-2013	31	Kjell Magnus Norderhaug	64, 65
Northern Chile Fondecyt # 1040425	Central Chile and Humboldtian	2004-2007	4	Alejandro Perez Matus	66
Northern Ireland Summary	Celtic Seas	1982-2012	226	Julia Nunn and Claire Goodwin	
Haida Gwaii	North American Pacific Fjordland	2010-2013	8	Anne Salomon	
Southern Chile	Chiloense	2012	1	Alejandro Buschmann	67

Edwards Baja	Northern California and Southern California Bight	1997-2006	62	Matt Edwards	
Kelp Cover Australia Victoria	Bassian and Cape Howe	1998-2011	45	Scott Ling	68
Australia Connell Canopy Loss	South Australian Gulfs	1968-2008	2	Sean Connell	69
Steneck Gulf of Maine	Gulf of Maine	1996-2003	16	Bob Steneck	
Konar Gulf of Alaska	Gulf of Alaska	2003-2012	4	Brenda Konar	
Norway Inner Coast (monitoring)	North Sea/Southern Norway	1976-2012	1	Kjell Magnus Norderhaug	70
Norway Inner Coast (assessment)	North Sea and Southern Norway	1975-2006	9	Frithjof E Moy	71, 72
Normandeau Associates	Gulf of Maine	1978-2011	4	Paul Geoghegan	Error! Reference source not found.




Figure S1. Black circles show site level estimates of mean slope with bars denoting 90% credible intervals (data from 1-72). Sites are grouped by ecoregion and realm, with solid black vertical lines showing ecoregion level mean slopes and shaded bars indicating 90% credible intervals. Dashed lines denote 0. Bars are colored red where there is a high probability (>95%) that the slope does not overlap with zero.




Figure S1 continued




Figure S1 continued




Figure S2. Raw data by ecoregion with site and ecoregion level modeled slopes (data from 1-72). Data and slope estimates are colored and symbolized by study (i.e., literature reference or contributed data set), while black lines show the ecoregion mean slope. Solid lines are used where site and ecoregion level slopes have a high probability ($>95\%$) of being different from zero, and dashed lines are used where slopes do not meet this criterion. Shaded bands show the 90% credible interval around the mean rate of change (the black line).




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S2 continued




Figure S3. The number of sites in the global data set by year (data from 1-72).




Figure S4. A) Site level slopes and associated standard error, plotted against study duration and colored according to sign (positive, negative, or no change) (data from 1-72). B) Distribution of site level slope estimates colored by sign (positive, negative, or no change).

References

1. Davoult D, Engel CR, Arzel P, Knoch D, Laurans M. 2011. Environmental factors and commercial harvesting: exploring possible links between the decline of the kelp *Laminaria digitata* in Brittany, France. *Cah Mar Biol* 52:429-435
2. Trowbridge CD, Little C, Dlouhy-Massengale B, Stirling P, Pilling GM. 2013. Changes in brown seaweed distributions in Lough Hyne, SW Ireland: a long-term perspective. *Botanica Marina* 56:323-338
3. Dean TA, Bodkin JL, Jewett SC, Monson DH, Jung D. 2000. Changes in sea urchins and kelp following a reduction in sea otter density as a result of the Exxon Valdez oil spill. *Mar Ecol Prog Ser.* 199:281-291
4. Lambert WJ, Levin PS, Berman J. 1992. Changes in the structure of a New England (USA) kelp bed: the effects of an introduced species? *Mar Ecol Prog Ser* 88: 303-307
5. Gagnon P, Himmelman JH, Johnson LE. 2004. Temporal variation in community interfaces: kelp-bed boundary dynamics adjacent to persistent urchin barrens. *Mar Biol* 144:1191-1203
6. Gagnon P, Johnson LE, Himmelman JH. 2005. Kelp patch dynamics in the face of intense herbivory: stability of *Agarum clathratum* (Phaeophyta) stands and associated flora on urchin barrens. *J Phycol* 41:498-505
7. de Bettignies T, Wernberg T, Lavery P, Vanderklift MA, Mohring MB. 2013. Contrasting mechanisms of dislodgement and erosion contribute to production of kelp detritus. *Limnol Oceanogr* 58:1680-1688
8. Kirkman H. 1984. Standing stock and production of *Ecklonia radiata*. *J Exp Mar Biol Ecol.* 76:119-130
9. Kirkman H. 1989. Growth, density, and biomass of *Ecklonia radiata* at different depths and growth under artificial shading off Perth, Western Australia. *Aust. J. Mar. Freshwater Res.* 40:169-177
10. Vásquez JA, Vega JMA, Buschmann AH. 2006. Long term variability in the structure of kelp communities in Northern Chile and then 1997-1998 ENSO. *J App Phycol* 18:505-519
11. Andrew NL. 1991. Changes in subtidal habitat following mass mortality of sea urchins in Botany Bay, New South Wales. *Australian Journal of Ecology* 16:353-362

12. Andrew NL, Jones GP. 1990. Patch formation by herbivorous fish in a temperature Australian kelp forest. *Oecologia* 85:57-68
13. Ettinger-Epstein P., Kingsford MJ. 2008. Effects of the El Nino southern oscillation on *Turbo torquatus* (Gastropoda) and their kelp habitat. *Austral Ecology* 33:594-606
14. Anderson RJ, Rothman MD, Share A, Drummond. 2006. Harvesting of the kelp *Ecklonia maxima* in South Africa affects its three obligate, red algal epiphytes. *J App Phycol* 18: 343-349
15. Pulfrich A, Parkins CA, Branch GM. 2003. The effects of shore-based diamond-diving on intertidal and subtidal biological communities and rock lobsters in southern Namibia. *Aquatic Conserv Mar Freshw Ecosyst* 13: 233-255
16. Duggins DO. 1980. Kelp beds and sea otters: An experimental approach. *Ecology* 61: 447-453
17. Estes JA, Duggins DO. 1995. Sea otters and kelp forests in Alaska: generality and variation in a community ecological paradigm. *Ecol Monogr* 65: 75-100
18. Gerard VA. 1976. Some aspects of material dynamics and energy flow in a kelp forest in Monterey Bay, California. University of California.
19. Schiel DR, Steinbeck JR, Foster MS. 2004. Ten years of induced ocean warming causes comprehensive changes in marine benthic communities. *Ecology* 85: 1833-1839
20. Harrold C, Reed D. 1985. Food availability, sea urchin grazing, and kelp forest community structure. *Ecology* 66:1160-1169
21. Fox D, Amend M, Merems A, Miller B, Golden J. 1998. Nearshore rocky reef assessment. Oregon Department of Fish and Wildlife Marine Program.
22. Fox D, Amend M, Merems A, Appy M. 2000. Nearshore rocky reef assessment. Oregon Department of Fish and Wildlife Marine Program.
23. Fox D, Merems A, Amend M, Weeks H, Romsos C, Appy M. 2004. Nearshore rocky reef assessment. Oregon Department of Fish and Wildlife Marine Program.
24. Kvitek RG, Iampietro PJ, Thomas K. 2001. Quantitative assessment of sea otter benthic prey communities within the Olympic Coast National Marine Sanctuary: 1999 re-survey of 1995 and 1985 monitoring stations. Puget Sound Research

25. Shaffer JA. 2000. Seasonal variation in understory kelp bed habitats of the Strait of Juan de Fuca. *J Coast Res* 16: 768-775
26. Feehan C, Scheibling RE, Lauzon-Guay JS. 2012. Aggregative feeding behavior in sea urchins leads to destructive grazing in a Nova Scotian kelp bed. *Mar Ecol Prog Ser* 444: 69-83
27. Gerard VA, Mann KH. 1979. Growth and production of *Laminaria longicruris* (Phaeophyta) populations exposed to different intensities of water movement. *J Phycol* 15: 33-41
28. Kelly JR, Scheibling RE, Balch T. 2011. Invasion-mediated shifts in the macrobenthic assemblage of a rocky subtidal ecosystem. *Mar Ecol Prog Ser* 437:69-78
29. Krumhansl KA, Scheibling RE. 2012. Detrital production in Nova Scotian kelp beds: patterns and processes. *Mar Ecol Prog Ser* 421:67-82
30. Lauzon-Guay JS, Scheibling RE. 2007. Behaviour of sea urchin *Strongylocentrotus droebachiensis* grazing fronts: food-mediated aggregation and density-dependent facilitation. *Mar Ecol Prog Ser* 329: 181-204
31. Mann KH. 1972. Ecological energetics of the seaweed zone in a marine bay on the Atlantic Coast of Canada. I. Zonation and biomass of seaweeds. *Mar Biol* 12:1-10
32. MacFarlane C. 1952. A survey of certain seaweeds of commercial importance in southwest Nova Scotia. *Can J Bot* 30: 78-97
33. Miller RJ. 1985. Succession in sea urchin and seaweed abundance in Nova Scotia, Canada. *Mar Biol* 84:275-286
34. O'Brien JM, Scheibling RE, Krumhansl KA. 2015. Positive feedback between large-scale disturbance and density-dependent grazing decreases resilience of a kelp bed ecosystem. *Mar Ecol Prog Ser* 522:1-13
35. Saunders MI, Metaxas A. 2009. Population dynamics of a nonindigenous epiphytic bryozoan *Membranipora membranacea* in the western North Atlantic: effects of kelp substrate. *Aquat Biol* 8:83-94
36. Scheibling R. 1986. Increased macroalgal abundance following mass mortalities of sea urchins (*Strongylocentrotus droebachiensis*) along the Atlantic coast of Nova Scotia. *Oecologia* 68:186-198.
37. Scheibling RE, Gagnon P. 2009. Temperature-mediated outbreak dynamics of the invasive bryozoan *Membranipora membranacea*. *Mar Ecol Prog Ser* 390:1-13

38. Smith BD. 1985. Recovery following experimental harvesting of *Laminaria longicruris* and *L. digitata* in southwestern Nova Scotia. *Helgo Meer* 39:83-101
39. Smith BD. 1988. Comparison of productivity estimates for *Laminaria* in Nova Scotia. *Can J Fish Aquat Sci* 45:557- 562
40. Yorke AF, Metaxas A. 2012. Relative importance of kelps and fucoids as substrata of the invasive epiphytic bryozoan *Membranipora membranacea* in Nova Scotia, Canada. *Aquat Biol* 16:17-30
41. Dayton PK, Tegner MJ, Parnell PE, Edwards PB. 1992. Temporal and spatial patterns of disturbance and recovery in a kelp forest community. *Ecol Monogr* 62:421-445
42. Dayton PK, Tegner MJ, Edwards PB, Riser KL. 1999. Temporal and spatial scales of kelp demography: the role of oceanographic climate. *Ecol Monogr* 69:219-250
43. Leinaas HP, Christie H. 1996. Effects of removing sea urchins (*Strongylocentrotus droebachiensis*): stability of the barren state and succession of kelp forest recover in the East Atlantic. *Oecologia* 105:524-536
44. Watson J, Estes JA (2011). Stability, resilience, and phase shifts in rocky subtidal communities along the west coast of Vancouver Island, Canada. *Ecol. Monogr.* 81:215-239.
45. Kenner MC et al. (2013) A multi-decade time series of kelp forest community structure at San Nicolas Island, California (USA). *Ecology* 94:2654–2654.
46. Witman JD, Genovese SJ, Bruno JF, McLaughlin JW, Pavlin BI (2003) Massive prey recruitment and the control of rocky subtidal communities on large spatial scales. *Ecol Monogr* 73:441-462.
47. Witman JD (1987) Subtidal coexistence: storms, grazing, mutualism, and the zonation of kelps and mussels. *Ecol Monogr* 57:167-187.
48. Kushner DJ, Rassweiler A, McLaughlin JP, Lafferty KD (2013) A multi-decade time series of kelp forest community structure at the California Channel Islands. *Ecology* 94:2655
49. Pérez-Matus A, Carrasco S, Gelcich S, Fernandez M, Wieters E. Upwelling and anthropogenic factors influence functional community structure of subtidal kelp forests of Central Chile. *Ecosphere* (In Review)

50. Levitt GJ, Anderson RJ, Boothroyd CJT, and Kemp FA (2002) The effects of kelp harvesting on its regrowth and the understory benthic community at danger point, South Africa, and a new method of harvesting kelp fronds. *S Afr J Mar Sci* 24: 71-95
51. Reed DC (2014) SBC LTER: Reef: Kelp Forest Community Dynamics: Abundance and size of giant kelp (*Macrocystis pyrifera*), ongoing since 2000. Santa Barbara Coastal LTER. doi:10.6073/pasta/d90872297e30026b263a119d4f5bca9f
52. Edgar GJ, Barrett NS (1997) Short term monitoring of biotic change in Tasmanian marine reserves. *Journal of Experimental Marine Biology and Ecology* 213:261-279
53. Edgar GJ, Barrett NS (1999) Effects of the declaration of marine reserves on Tasmanian reef fishes, invertebrates and plants. *Journal of Experimental Marine Biology and Ecology* 242:107-144
54. Barrett NS, Buxton CD, Edgar GJ (2009) Changes in invertebrate and macroalgal populations in Tasmanian marine reserves in the decade following protection. *Journal of Experimental Marine Biology and Ecology* 370:104-119
55. Micheli F et al. (2012) Evidence that marine reserves enhance resilience to climatic impacts. *PLoS ONE* 7(7): e40832
56. Sáenz-Arroyo A. et al. (2012) Proyecto de reservas marinas Isla Natividad, BCS, Mexico. Final Report, 48 pp.
57. Shears NT (2010) Long Bay Marine Monitoring Program: 1999-2010. Prepared by Auckland UniServices for Auckland Regional Council. Auckland Regional Council Technical Report
58. Shears NT, Babcock RC (2007) Quantitative description of mainland New Zealand's shallow subtidal reef communities. *Science for Conservation* 280: 126p.
59. Shears NT, Babcock RC (2003) Continuing trophic cascade effects after 25 years of no-take marine reserve protection. *Mar Ecol Prog Ser* 246: 1-16
60. Derrien-Courtél S (2008) L'étude des peuplements subtidaux rocheux (flore et faune) du littoral breton permet-elle de contribuer à l'évaluation de la qualité écologique du littoral et d'en mesurer les changements dans le temps ? Thèse de l'Ecole doctorale du Muséum « Sciences de la Nature et de l'Homme », Concarneau, France, p. 222.
61. Derrien-Courtél S, Le Gal A, Grall J (2013) Regional-scale analysis of subtidal rocky shore community. *Helgol Mar Res* 67:697-712

62. Le Gal A, Derrien-Courtet S (2015) Quality Index of Subtidal Macroalgae (QISubMac), a suitable tool for ecological quality status assessment under the scope of the European Water Framework Directive. *Mar Poll Bull* 101:334-349
63. Wernberg T. et al. (2013) An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nature Climate Change*, 3:78–82
64. Norderhaug KM et al. (2015) Combined effects from climate variation and eutrophication on the diversity in hard bottom communities on the Skagerrak coast 1990-2010. *Mar Ecol Prog Ser* 530: 29–46
65. Norderhaug KM et al. (2011) Long-term monitoring of environmental quality in the coastal regions of Norway. Klif report TA-2777. 115 pp.
66. Perez Matus A, Ferry-Graham L , Cea A, Vásquez JA (2007) Community structure of temperate reef fishes in kelp-dominated subtidal habitats of northern Chile. *Mar Fresh Res* 58: 1069–1085.
67. Buschmann AH et al. (2014) Ecophysiological plasticity of annual populations of giant Kelp (*Macrocystis pyrifera*) in a seasonally variable Coastal environmental in the Northern Patagonian Inner Seas of Southern Chile. *J Appl Phycol* 26:837-847
68. Edmunds M, Hart S (2003) Parks Victoria Standard Operating Procedure. Biological Monitoring of Subtidal Reefs. *Parks Victoria Technical Series* 9:1-48
69. Connell SD et al. (2008) Recovering a lost baseline: missing kelp forests from a metropolitan coast. *Mar Ecol Prog Ser* 360: 63–72
70. Norderhaug KM et al. (2012) Miljøovervåking av sukkertare lang kysten. Sugar kelp monitoring programme. Yearly KLIF report. 47 pp.
71. Moy FE, Christie H (2012) Large-scale shift from sugar kelp (*Saccharina latissima*) to ephemeral algae along the south and west coast of Norway. *Mar Biol Res* 8:309-321
72. Åsen PA (2006) Marine macroalgae vegetation from Kristiansandsfjorden to Jøssingfjorden _with special reference to *Laminaria saccharina* and *Alaria esculenta*. Agder naturmuseums rapportserie 2006-4. 35 pages.