

Proof Theory: Logical and Philosophical Aspects

Class 3: Beyond Sequents

Greg Restall and Shawn Standefer



THE UNIVERSITY OF
MELBOURNE

NASSLLI · JULY 2016 · RUTGERS

To introduce *proof theory*, with a focus in its applications in philosophy, linguistics and computer science.

Introduce extensions of sequent systems to naturally deal with modal logics.

Today's Plan

Basic Modal Logic

Modal Sequent Systems

Display Logic

Labelled Sequents

Tree Hypersequents

BASIC MODAL LOGIC


Possibility and Necessity

Modal logic adds propositional logic the notions of *possibility* and *necessity*.


Add to the language of propositional logic the ' \Box ' and ' \Diamond .'

- If A is a formula, so are $\Box A$ and $\Diamond A$.


Example Interpretation




Example Interpretation



Example Interpretation



Example Interpretation



Modal Logic: Interpretations

An *interpretation* for the language is a triple: $\langle W, R, v \rangle$.

W is a non-empty set of *states* (or *possible worlds*).

R is a two-place relation on W , of *relative possibility*. uRw means that from the point of view of u , w is possible.

Finally, v assigns a truth value to a propositional parameter *at a state*.

That is, for each world w and propositional parameter p , we will have either $v_w(p) = 1$ (if p is “true at w ”) or $v_w(p) = 0$ (if p is “false at w ”).

Interpreting the Language

We keep the rules for the classical connectives, with state subscripts on v :

- ▶ $v_w(\neg A) = 1$ if and only if $v_w(A) = 0$.
- ▶ $v_w(A \wedge B) = 1$ if and only if $v_w(A) = 1$ and $v_w(B) = 1$.
- ▶ $v_w(A \vee B) = 1$ if and only if $v_w(A) = 1$ or $v_w(B) = 1$.
- ▶ $v_w(A \supset B) = 1$ if and only if $v_w(A) = 0$ or $v_w(B) = 1$.

No novelty there.

Interpreting the Language

We keep the rules for the classical connectives, with state subscripts on v :

- ▶ $v_w(\neg A) = 1$ if and only if $v_w(A) = 0$.
- ▶ $v_w(A \wedge B) = 1$ if and only if $v_w(A) = 1$ and $v_w(B) = 1$.
- ▶ $v_w(A \vee B) = 1$ if and only if $v_w(A) = 1$ or $v_w(B) = 1$.
- ▶ $v_w(A \supset B) = 1$ if and only if $v_w(A) = 0$ or $v_w(B) = 1$.

No novelty there.

The innovation is found with \Box and \Diamond :

- ▶ $v_w(\Box A) = 1$ if and only if $v_u(A) = 1$ for each u where wRu .
- ▶ $v_w(\Diamond A) = 1$ if and only if $v_u(A) = 1$ for some u where wRu .

Interpretations can be used to define validity, as with classical propositional logic.

Interpretations can be used to define validity, as with classical propositional logic.

The argument from X to Y is *valid* (written ' $X \vdash Y$ ' as before) if and only if for every interpretation $\langle W, R, v \rangle$ for any state $w \in W$, if $v_w(B) = 1$ for each $B \in X$ then for some $C \in Y$, $v_w(C) = 1$ too.

Modal Validity

Interpretations can be used to define validity, as with classical propositional logic.

The argument from X to Y is *valid* (written ' $X \vdash Y$ ' as before) if and only if for every interpretation $\langle W, R, v \rangle$ for any state $w \in W$, if $v_w(B) = 1$ for each $B \in X$ then for some $C \in Y$, $v_w(C) = 1$ too.

... or equivalently, there is no state $w \in W$ at which every member of X is true and every member of Y is false.

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

$$\frac{X, \Box A, \Box B \vdash Y}{X, \Box(A \wedge B) \vdash Y}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

$$\frac{X, \Box A, \Box B \vdash Y}{X, \Box(A \wedge B) \vdash Y}$$

$$\frac{X \vdash \Diamond A, \Diamond B, Y}{X \vdash \Diamond(A \vee B), Y}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

$$\frac{X, \Box A, \Box B \vdash Y}{X, \Box(A \wedge B) \vdash Y}$$

$$\frac{X \vdash \Diamond A, \Diamond B, Y}{X \vdash \Diamond(A \vee B), Y}$$

$$\frac{X, \Box A \vdash Y}{X, \neg \Diamond \neg A \vdash Y}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

$$\frac{X, \Box A, \Box B \vdash Y}{X, \Box(A \wedge B) \vdash Y}$$

$$\frac{X \vdash \Diamond A, \Diamond B, Y}{X \vdash \Diamond(A \vee B), Y}$$

$$\frac{X, \Box A \vdash Y}{X, \neg \Diamond \neg A \vdash Y}$$

$$\frac{X \vdash \Box A, Y}{X \vdash \neg \Diamond \neg A, Y}$$

Some Basic Validity Facts

$$\frac{\vdash A}{\vdash \Box A}$$

$$\frac{A \vdash}{\Diamond A \vdash}$$

$$\frac{A \vdash B}{\Box A \vdash \Box B}$$

$$\frac{A \vdash B}{\Diamond A \vdash \Diamond B}$$

$$\frac{X, \Box A, \Box B \vdash Y}{X, \Box(A \wedge B) \vdash Y}$$

$$\frac{X \vdash \Diamond A, \Diamond B, Y}{X \vdash \Diamond(A \vee B), Y}$$

$$\frac{X, \Box A \vdash Y}{X, \neg \Diamond \neg A \vdash Y}$$

$$\frac{X \vdash \Box A, Y}{X \vdash \neg \Diamond \neg A, Y}$$

None of these are much like good L/R rules for \Box or \Diamond .

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$
<i>transitivity</i>	$wRv \wedge vRu \supset wRu$	$\Box A \vdash \Box \Box A$	$\Diamond \Diamond A \vdash \Diamond A.$

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$
<i>transitivity</i>	$wRv \wedge vRu \supset wRu$	$\Box A \vdash \Box \Box A$	$\Diamond \Diamond A \vdash \Diamond A.$
<i>symmetry</i>	$wRv \supset vRw$	$A \vdash \Box \Diamond A$	$\Diamond \Box A \vdash A.$

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$
<i>transitivity</i>	$wRv \wedge vRu \supset wRu$	$\Box A \vdash \Box \Box A$	$\Diamond \Diamond A \vdash \Diamond A.$
<i>symmetry</i>	$wRv \supset vRw$	$A \vdash \Box \Diamond A$	$\Diamond \Box A \vdash A.$
<i>directedness</i>	$(\exists v)wRv$	$\Box \perp \vdash$	$\vdash \Diamond \top$

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$
<i>transitivity</i>	$wRv \wedge vRu \supset wRu$	$\Box A \vdash \Box \Box A$	$\Diamond \Diamond A \vdash \Diamond A.$
<i>symmetry</i>	$wRv \supset vRw$	$A \vdash \Box \Diamond A$	$\Diamond \Box A \vdash A.$
<i>directedness</i>	$(\exists v)wRv$	$\Box \perp \vdash$	$\vdash \Diamond \top$
\vdots		\vdots	

Moving Beyond Basic Modal Logic

Restrictions on the accessibility relation lead to properties for \Box and \Diamond .

CONDITION		PROPERTY	
<i>reflexivity</i>	wRw	$\Box A \vdash A$	$A \vdash \Diamond A.$
<i>transitivity</i>	$wRv \wedge vRu \supset wRu$	$\Box A \vdash \Box \Box A$	$\Diamond \Diamond A \vdash \Diamond A.$
<i>symmetry</i>	$wRv \supset vRw$	$A \vdash \Box \Diamond A$	$\Diamond \Box A \vdash A.$
<i>directedness</i>	$(\exists v)wRv$	$\Box \perp \vdash$	$\vdash \Diamond \top$
\vdots		\vdots	

K: *all models* T: *reflexive models* S4: *reflexive transitive models*
 S5: *reflexive symmetric transitive models.*

MODAL SEQUENT SYSTEMS

What could L/R rules for \Box and \Diamond look like?

$$\frac{??? \vdash ???}{X \vdash \Box A, Y}$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{X \vdash A}{\Box X \vdash \Box A}$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{\Box X \vdash A}{\Box X \vdash \Box A}$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

$$\frac{\Box X, A \vdash \Diamond Y}{\Box X, \Diamond A \vdash \Diamond Y} [\Diamond L]$$

$$\frac{X \vdash A, Y}{X \vdash \Diamond A, Y} [\Diamond R]$$

What could L/R rules for \Box and \Diamond look like?

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

$$\frac{\Box X, A \vdash \Diamond Y}{\Box X, \Diamond A \vdash \Diamond Y} [\Diamond L]$$

$$\frac{X \vdash A, Y}{X \vdash \Diamond A, Y} [\Diamond R]$$

These rules characterise the modal logic S4.

Example Derivations

What about S5?

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

What about S5?

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

$$\frac{\frac{\frac{\Box p \vdash \Box p}{\vdash \Box p, \neg \Box p} [\neg R]}{\vdash \Box p, \Box \neg \Box p} [\Box R'] \quad \frac{p \vdash p}{\Box p \vdash p} [\Box L]}{\vdash p, \Box \neg \Box p} [Cut]$$

What about S5?

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

$$\frac{\frac{\Box p \vdash \Box p}{\vdash \Box p, \neg \Box p} [\neg R] \quad \frac{p \vdash p}{\Box p \vdash p} [\Box L]}{\vdash \Box p, \Box \neg \Box p} [\Box R']$$

$$\frac{\vdash \Box p, \Box \neg \Box p \quad \Box p \vdash p}{\vdash p, \Box \neg \Box p} [Cut]$$

The sequent $\vdash p, \Box \neg \Box p$ has *no* cut-free proof.
(How could you apply a \Box rule?)

Problems with these \Box and \Diamond rules

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Box X, A \vdash \Diamond Y}{\Box X, \Diamond A \vdash \Diamond Y} [\Diamond L]$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

$$\frac{X \vdash A, Y}{X \vdash \Diamond A, Y} [\Diamond R]$$

Entanglement between \Box and \Diamond .

Problems with these \Box and \Diamond rules

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Box X, A \vdash \Diamond Y}{\Box X, \Diamond A \vdash \Diamond Y} [\Diamond L]$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

$$\frac{X \vdash A, Y}{X \vdash \Diamond A, Y} [\Diamond R]$$

Entanglement between \Box and \Diamond .

*$\Box L$ and $\Diamond R$ are weak
— all the work is done by
the left \Diamond rules and right \Box rules.*

Problems with these \Box and \Diamond rules

$$\frac{X, A \vdash Y}{X, \Box A \vdash Y} [\Box L]$$

$$\frac{\Box X \vdash A, \Diamond Y}{\Box X \vdash \Box A, \Diamond Y} [\Box R]$$

$$\frac{\Box X \vdash A, \Box Y}{\Box X \vdash \Box A, \Box Y} [\Box R']$$

$$\frac{\Box X, A \vdash \Diamond Y}{\Box X, \Diamond A \vdash \Diamond Y} [\Diamond L]$$

$$\frac{\Diamond X, A \vdash \Diamond Y}{\Diamond X, \Diamond A \vdash \Diamond Y} [\Diamond L']$$

$$\frac{X \vdash A, Y}{X \vdash \Diamond A, Y} [\Diamond R]$$

Entanglement between \Box and \Diamond .

*$\Box L$ and $\Diamond R$ are weak
— all the work is done by
the left \Diamond rules and right \Box rules.*

Hard/impossible to generalise.

From Modal to *Temporal* Logic

- ▶ $v_w(\Box A) = 1$ if and only if $v_u(A) = 1$ for each u where wRu .
- ▶ $v_w(\Diamond A) = 1$ if and only if $v_u(A) = 1$ for some u where wRu .

- ▶ $v_w(\blacksquare A) = 1$ if and only if $v_u(A) = 1$ for each u where uRw .
- ▶ $v_w(\blacklozenge A) = 1$ if and only if $v_u(A) = 1$ for some u where uRw .

From Modal to *Temporal* Logic

- ▶ $v_w(\Box A) = 1$ if and only if $v_u(A) = 1$ for each u where wRu .
- ▶ $v_w(\Diamond A) = 1$ if and only if $v_u(A) = 1$ for some u where wRu .
- ▶ $v_w(\blacksquare A) = 1$ if and only if $v_u(A) = 1$ for each u where uRw .
- ▶ $v_w(\blacklozenge A) = 1$ if and only if $v_u(A) = 1$ for some u where uRw .

$$\frac{A \vdash \Box B}{\blacklozenge A \vdash B}$$

$$\frac{\Diamond A \vdash B}{A \vdash \blacksquare B}$$

Going Forward and Back in a Derivation

$$\begin{array}{c}
 \frac{\frac{\square A, \square B \vdash \square A}{\square A \wedge \square B \vdash \square A} [\wedge L]}{\diamond(\square A \wedge \square B) \vdash \textcolor{yellow}{B}} [\square \diamond] \\
 \frac{\frac{\frac{\square A, \square B \vdash \square B}{\square A \wedge \square B \vdash \square B} [\wedge L]}{\diamond(\square A \wedge \square B) \vdash B} [\square \diamond]}{\diamond(\square A \wedge \square B) \vdash A \wedge B} [\wedge R] \\
 \frac{\diamond(\square A \wedge \square B) \vdash A \wedge B}{\square A \wedge \square B \vdash \square(A \wedge B)} [\diamond \square]
 \end{array}$$

Generalised Sequents

How do we establish $X \vdash \Box A, Y$?

Generalised Sequents

How do we establish $X \vdash \Box A, Y$?

It should have *something* to do with some $X' \vdash A, Y'$
but the A is evaluated in a different *state*.

Generalised Sequents

How do we establish $X \vdash \Box A, Y$?

It should have *something* to do with some $X' \vdash A, Y'$

but the A is evaluated in a different *state*.

We need to record state shifts in sequents.

Generalised Sequents

How do we establish $X \vdash \Box A, Y$?


It should have *something* to do with some $X' \vdash A, Y'$

but the A is evaluated in a different *state*.

We need to record state shifts in sequents.

DISPLAY LOGIC • LABELLED SEQUENTS • TREE HYPERSEQUENTS

DISPLAY LOGIC



Nuel Belnap

Sequents

Sequents are of the form $X \vdash Y$, where X and Y are *structures*

Structures are built up out of formulas and the structural connectives $*$, \bullet (both unary), and \circ (binary)

For example, $*(p \circ q) \vdash \bullet(r \circ *s)$

Display equivalences

Certain sequents are stipulated to be equivalent via *display equivalences*

$$X \vdash Y \circ Z \iff X \circ *Y \vdash Z \iff X \vdash Z \circ Y$$

$$X \vdash Y \iff *Y \vdash *X \iff X \vdash **Y$$

$$\bullet X \vdash Y \iff X \vdash \bullet Y$$

Display equivalences

Certain sequents are stipulated to be equivalent via *display equivalences*

$$X \vdash Y \circ Z \iff X \circ *Y \vdash Z \iff X \vdash Z \circ Y$$

$$X \vdash Y \iff *Y \vdash *X \iff X \vdash **Y$$

$$\bullet X \vdash Y \iff X \vdash \bullet Y$$

(These rules ensure that $*$ acts like *negation*,
 \circ is *conjunctive* on the left and *disjunctive* on the right,
and \bullet acts like a *necessity* on the right
and its *converse possibility* the left.)

Displaying

By means of the display equivalences, one can display a formula or structure on one side of the turnstile in isolation

This permits the left and right rules to deal with only the displayed formulas and structures

$$\frac{A \circ B \vdash X}{A \wedge B \vdash X} [\wedge L]$$

$$\frac{X \vdash A \quad Y \vdash B}{X \circ Y \vdash A \wedge B} [\wedge R]$$

Generality

The connectives rules are formulated so that each connective is paired with a structural connective

Different logical behaviour is obtained by imposing different rules on the structural connectives

A single form of conjunction rule can be used for, say, classical conjunction and relevant fusion, the difference coming out in the structural rules in force

Cut

Because formulas can always be displayed,
a simple form of *Cut* can be used for a range of logics

$$\frac{X \vdash A \quad A \vdash Y}{X \vdash Y} [Cut]$$

Eliminating Cut

The *Elimination Theorem* is proved via a general argument that depends on eight conditions on the rules.

If these conditions are satisfied, then it follows that *Cut* is admissible

This argument is due to *Haskell Curry* and *Nuel Belnap*.

The Structure of the Curry–Belnap Cut Elimination Proof

- ▶ It's a *Cut* elimination argument (it doesn't appeal to a Mix rule).
- ▶ It's an induction on *grade* (complexity of the *Cut* formula), as usual.
- ▶ To eliminate a *Cut* on a formula A , trace the *parametric* occurrences of a formula in the premises of the cut inference upward to where they first appear. Replace the cut at those instances (either with cuts on subformulas, or by weakening, or the cuts evaporate into identities) and then replay the substitution downward.

The Crucial Step

$$\begin{array}{c}
 \vdots \\
 \hline
 \dots A, A \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots A, A \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots A, A \dots \\
 \hline
 \dots A \dots
 \end{array}$$

$$\begin{array}{c}
 \dots A \dots \quad \dots A \dots \\
 \hline
 \dots A \dots
 \end{array}$$

$$\begin{array}{c}
 \dots A \dots \\
 \hline
 A \vdash Y
 \end{array}$$

$$\frac{X \vdash A \quad A \vdash Y}{X \vdash Y} [Cut]$$

The Crucial Step

$$\begin{array}{c}
 \vdots \\
 \hline
 \dots A, A \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots A, A \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots A \dots
 \end{array}$$

$$\begin{array}{c}
 \dots A \dots \quad \dots A \dots \\
 \hline
 \dots A \dots
 \end{array}$$

$$\begin{array}{c}
 \dots A \dots \\
 \hline
 A \vdash Y
 \end{array}$$

The Crucial Step

$$\begin{array}{c}
 \vdots \\
 \hline
 \dots X, X \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots X, X \dots \\
 \hline
 \dots A \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots X \dots \\
 \hline
 \dots A \dots
 \end{array}$$

$$\begin{array}{c}
 \dots A \dots \quad \dots A \dots \\
 \hline
 \dots A \dots \\
 \hline
 A \vdash Y
 \end{array}$$

The Crucial Step

$$\begin{array}{c}
 \vdots \\
 \hline
 \dots X, X \dots \\
 \hline
 \dots X \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots X, X \dots \\
 \hline
 \dots X \dots
 \end{array}
 \quad
 \begin{array}{c}
 \vdots \\
 \hline
 \dots X, X \dots \\
 \hline
 \dots X \dots
 \end{array}$$

$$\begin{array}{c}
 \dots X \dots \quad \dots X \dots \\
 \hline
 \dots X \dots
 \end{array}$$

$$\begin{array}{c}
 \dots X \dots \\
 \hline
 X \vdash Y
 \end{array}$$

The Eight Conditions

- ▶ c1: *Preservation of formulas.*
- ▶ c2: *Shape-alikeness of parameters.*
- ▶ c3: *Non-proliferation of parameters.*
- ▶ c4: *Position-alikeness of parameters.*
- ▶ c5: *Display of principal constituents.*
- ▶ c6: *Closure under substitution for consequent parameters.*
- ▶ c7: *Closure under substitution for antecedent parameters.*
- ▶ c8: *Eliminability of matching principal constituents.*

Modal Rules

To give rules for modal operators, you use the *modal structure*.

$$\frac{A \vdash Y}{\Box A \vdash \bullet Y} [\Box L]$$

$$\frac{X \vdash \bullet B}{X \vdash \Box B} [\Box R]$$

Example Display Logic Derivation

$$\begin{array}{c}
 \frac{A \vdash A}{\square A \vdash \bullet A} [\square L] \\
 \frac{\square A \vdash \bullet A}{\square A \circ \square B \vdash \bullet A} [K] \\
 \frac{\square A \circ \square B \vdash \bullet A}{\bullet(\square A \circ \square B) \vdash A} [display] \\
 \frac{\bullet(\square A \circ \square B) \vdash A}{\bullet(\square A \circ \square B) \circ \bullet(\square A \circ \square B) \vdash A \wedge B} [\wedge R] \\
 \frac{\bullet(\square A \circ \square B) \circ \bullet(\square A \circ \square B) \vdash A \wedge B}{\bullet(\square A \circ \square B) \vdash A \wedge B} [W] \\
 \frac{\bullet(\square A \circ \square B) \vdash A \wedge B}{\square A \circ \square B \vdash \bullet A \wedge B} [display] \\
 \frac{\square A \circ \square B \vdash \bullet A \wedge B}{\square A \wedge \square B \vdash \bullet A \wedge B} [\wedge L] \\
 \frac{\square A \wedge \square B \vdash \bullet A \wedge B}{\square A \wedge \square B \vdash \square(A \wedge B)} [\square R]
 \end{array}$$

Structural Rules

$$\frac{X \vdash \bullet Y}{X \vdash Y} \text{ [refl]}$$

$$\frac{\frac{A \vdash A}{\Box A \vdash \bullet A} \text{ [\Box L]}}{\Box A \vdash A} \text{ [refl]}$$

Structural Rules

$$\frac{X \vdash \bullet Y}{X \vdash Y} \text{ [refl]}$$

$$\frac{X \vdash \bullet \bullet Y}{X \vdash \bullet Y} \text{ [trans]}$$

$$\begin{array}{c} \frac{A \vdash A}{\bullet A \vdash \Box A} \text{ [\Box R]} \\ \hline \frac{\bullet A \vdash \Box A}{A \vdash \bullet \Box A} \text{ [display]} \\ \hline \frac{A \vdash \bullet \Box A}{\Box A \vdash \bullet \bullet \Box A} \text{ [\Box L]} \\ \hline \frac{\Box A \vdash \bullet \bullet \Box A}{\Box A \vdash \bullet \Box A} \text{ [trans]} \\ \hline \frac{\Box A \vdash \bullet \Box A}{\Box A \vdash \Box \Box A} \text{ [\Box R]} \end{array}$$

Structural Rules

$$\frac{X \vdash \bullet Y}{X \vdash Y} \text{ [refl]}$$

$$\frac{X \vdash \bullet \bullet Y}{X \vdash \bullet Y} \text{ [trans]}$$

$$\frac{X \vdash \bullet * Y}{X \vdash * \bullet Y} \text{ [sym]}$$

$$\begin{array}{c} \frac{A \vdash A}{*A \vdash *A} \text{ [display]} \\ \frac{*A \vdash *A}{\neg A \vdash *A} \text{ [\neg L]} \\ \frac{\neg A \vdash *A}{\Box \neg A \vdash \bullet * A} \text{ [\Box L]} \\ \frac{\Box \neg A \vdash \bullet * A}{\Box \neg A \vdash * \bullet A} \text{ [sym]} \\ \frac{\Box \neg A \vdash * \bullet A}{\bullet A \vdash * \Box \neg A} \text{ [display]} \\ \frac{\bullet A \vdash * \Box \neg A}{\bullet A \vdash \neg \Box \neg A} \text{ [\neg R]} \\ \frac{\bullet A \vdash \neg \Box \neg A}{A \vdash \bullet \neg \Box \neg A} \text{ [display]} \\ \frac{A \vdash \bullet \neg \Box \neg A}{A \vdash \Box \neg \Box \neg A} \text{ [\Box R]} \end{array}$$

Structural Rules

$$\frac{X \vdash \bullet Y}{X \vdash Y} \text{ [refl]}$$

$$\frac{X \vdash \bullet \bullet Y}{X \vdash \bullet Y} \text{ [trans]}$$

$$\frac{X \vdash \bullet * Y}{X \vdash * \bullet Y} \text{ [sym]}$$

*Many more structural rules
are possible.*

Cut Elimination: The \Box Case

A cut on a principal $\Box A$ may be simplified into a cut on A .

$$\frac{\frac{X \vdash \bullet A}{X \vdash \Box A} [\Box R] \quad \frac{A \vdash Y}{\Box A \vdash \bullet Y} [\Box L]}{X \vdash \bullet Y} [Cut]$$

Cut Elimination: The \Box Case

A cut on a principal $\Box A$ may be simplified into a cut on A .

$$\frac{\frac{X \vdash \bullet A}{X \vdash \Box A} [\Box R] \quad \frac{A \vdash Y}{\Box A \vdash \bullet Y} [\Box L]}{X \vdash \bullet Y} [Cut]$$

$$\frac{\frac{X \vdash \bullet A}{\bullet X \vdash A} [display] \quad A \vdash Y}{\bullet X \vdash Y} [Cut]$$

$$\frac{\bullet X \vdash Y}{X \vdash \bullet Y} [display]$$

Virtues and Vices of Display Logic

DISPLAY	
<i>Cut-free</i>	+
<i>Explicit</i>	+
<i>Systematic</i>	+
<i>Separation</i>	+
<i>Subformula</i>	+
<i>Nonredundant</i>	—
<i>Gentzen-plus</i>	—

Virtues and Vices of Display Logic

DISPLAY	
<i>Cut-free</i>	+
<i>Explicit</i>	+
<i>Systematic</i>	+
<i>Separation</i>	+
<i>Subformula</i>	+
<i>Nonredundant</i>	—
<i>Gentzen-plus</i>	—

LABELLED SEQUENTS

Recall this derivation...

$$\begin{array}{c}
 \frac{A \vdash A}{\square A \vdash \bullet A} [\square L] \\
 \frac{\square A \vdash \bullet A}{\square A \circ \square B \vdash \bullet A} [K] \\
 \frac{\square A \circ \square B \vdash \bullet A}{\bullet(\square A \circ \square B) \vdash A} [display] \\
 \frac{\bullet(\square A \circ \square B) \vdash A}{\bullet(\square A \circ \square B) \vdash B} [\wedge R] \\
 \frac{\bullet(\square A \circ \square B) \vdash A \wedge B}{\bullet(\square A \circ \square B) \vdash A \wedge B} [W] \\
 \frac{\bullet(\square A \circ \square B) \vdash A \wedge B}{\square A \circ \square B \vdash \bullet A \wedge B} [display] \\
 \frac{\square A \circ \square B \vdash \bullet A \wedge B}{\square A \wedge \square B \vdash \bullet A \wedge B} [\wedge L] \\
 \frac{\square A \wedge \square B \vdash \bullet A \wedge B}{\square A \wedge \square B \vdash \square(A \wedge B)} [\square R]
 \end{array}$$

Here is another way to represent it

$$\begin{array}{c}
 \frac{\frac{\frac{}{v : A \vdash v : A} [\Box L]}{wRv, w : \Box A \vdash v : A} [K]}{wRv, w : \Box A, w : \Box B \vdash v : A} \\
 \frac{\frac{\frac{}{v : B \vdash v : B} [\Box L]}{wRv, w : \Box B \vdash v : B} [K]}{wRv, w : \Box A, w : \Box B \vdash v : B} [\wedge R] \\
 \frac{wRv, w : \Box A, w : \Box B, wRv, w : \Box A, w : \Box B \vdash v : A \wedge B} [\wedge L] \\
 \frac{wRv, w : \Box A, w : \Box B \vdash A \wedge B} [\Box R] \\
 \frac{wRv, w : \Box A \wedge \Box B \vdash v : A \wedge B} [\Box R] \\
 w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)
 \end{array}$$

Labelled Sequent Rules: Boolean Connectives

$$x : A \vdash x : A$$

(Plus weakening and contraction.)

$$\frac{x : A, x : B, X \vdash Y}{x : A \wedge B, X \vdash Y} [\wedge L]$$

$$\frac{X \vdash x : A, Y \quad X \vdash x : B, Y}{X \vdash x : A \wedge B, Y} [\wedge R]$$

$$\frac{x : A, X \vdash Y \quad x : B, X \vdash Y}{x : A \vee B, X \vdash Y} [\vee L]$$

$$\frac{X \vdash x : A, x : B, Y}{X \vdash x : A \vee B, Y} [\vee R]$$

$$\frac{X \vdash x : A, Y}{x : \neg A, X \vdash Y} [\neg L]$$

$$\frac{x : A, X \vdash Y}{X \vdash x : \neg A, Y} [\neg R]$$

Labelled Sequent Rules: Modal Operators

$$\frac{x : A, X \vdash Y}{y R x, y : \Box A, X \vdash Y} [\Box L]$$

$$\frac{x R y, X \vdash y : A, Y}{X \vdash x : \Box A, Y} [\Box R]$$

$$\frac{x R y, y : A, X \vdash Y}{x : \Diamond A, X \vdash Y} [\Diamond L]$$

$$\frac{X \vdash x : A, Y}{y R x, X \vdash y : \Diamond A, Y} [\Diamond R]$$

In $\Box R$ and $\Diamond L$, the label y must not be present in X, Y or be identical to x .

Labelled Sequents

In these rules (except for weakenings) relational statements (xRy) are introduced only on the left of the sequent.

We may without loss of deductive power, restrict our attention to sequents in $X \vdash Y$ which relational statements appear only in X and not in Y .

The ‘cash value’ of a labelled sequent $X \vdash Y$ on a Kripke model is found by replacing $x : A$ by $v_x(A) = 1$; X by its conjunction; Y by its disjunction; the \vdash by a conditional; and universally quantifying over all world labels.

Frame conditions

The ‘cash value’ of a labelled sequent $X \vdash Y$ on a Kripke model is found by replacing $x : A$ by $v_x(A) = 1$; X by its conjunction; Y by its disjunction; the \vdash by a conditional; and universally quantifying over all world labels.

$xRy, x : A \vdash y : B, x : C$ is valid on a model if and only if

$$(\forall x, y)((xRy \wedge v_x(A) = 1) \supset ((v_y(B) = 1) \vee v_x(C) = 1))$$

Translation

A systematic translation maps modal display derivations into labelled modal derivations.

The translation simplifies the proof structure, erasing display equivalences, which are mapped to identical labelled sequents (*modulo* relabelling).

For details, see Poggiolesi and Restall “Interpreting and Applying Proof Theory for Modal Logic” (2012).

Virtues and Vices

	DISPLAY	LABELLED
<i>Cut-free</i>	+	+
<i>Explicit</i>	+	+
<i>Systematic</i>	+	+
<i>Separation</i>	+	+
<i>Subformula</i>	+	+—
<i>Nonredundant</i>	—	+—
<i>Gentzen-plus</i>	—	+—

Virtues and Vices

	DISPLAY	LABELLED
<i>Cut-free</i>	+	+
<i>Explicit</i>	+	+
<i>Systematic</i>	+	+
<i>Separation</i>	+	+
<i>Subformula</i>	+	+—
<i>Nonredundant</i>	—	+—
<i>Gentzen-plus</i>	—	+—

TREE HYPERSEQUENTS

Inspecting the translation

Display equivalent sequents correspond to *nearly* identical labelled sequents.

$$A \vdash \bullet B$$

$$\bullet A \vdash B$$

Inspecting the translation

Display equivalent sequents correspond to *nearly* identical labelled sequents.

$$A \vdash \bullet B \quad \Rightarrow \quad vRw, v : A \vdash w : B$$

$$\bullet A \vdash B$$

Inspecting the translation

Display equivalent sequents correspond to *nearly* identical labelled sequents.

$$A \vdash \bullet B \quad \Rightarrow \quad vRw, v : A \vdash w : B$$

$$\bullet A \vdash B \quad \Rightarrow \quad wRv, w : A \vdash v : B$$

Inspecting the translation

Display equivalent sequents correspond to *nearly* identical labelled sequents.

$$A \vdash \bullet B \quad \Rightarrow \quad vRw, v : A \vdash w : B$$

$$\bullet A \vdash B \quad \Rightarrow \quad wRv, w : A \vdash v : B$$

All we *care* about is that one world accesses the other. We have

$$A \vdash \longrightarrow \vdash B$$

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

- There is one node for every label.

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

- ▶ There is one node for every label.
- ▶ Every node is a sequent.

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

- ▶ There is one node for every label.
- ▶ Every node is a sequent.
- ▶ For every instance of $x : A$ in antecedent position, put A in the antecedent of the sequent at the node corresponding to x .

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

- ▶ There is one node for every label.
- ▶ Every node is a sequent.
- ▶ For every instance of $x : A$ in antecedent position, put A in the antecedent of the sequent at the node corresponding to x .
- ▶ For every instance of $x : A$ in consequent position, put A in the consequent of the sequent at the node corresponding to x .

The Recipe

Replace the labelled sequent $\mathcal{R}, X \vdash Y$ by a *directed graph* of sequents:

- ▶ There is one node for every label.
- ▶ Every node is a sequent.
- ▶ For every instance of $x : A$ in antecedent position, put A in the antecedent of the sequent at the node corresponding to x .
- ▶ For every instance of $x : A$ in consequent position, put A in the consequent of the sequent at the node corresponding to x .
- ▶ If \mathcal{R} contains Rxy , then place an arc from x to y .

Three ways of presenting the one fact

- Display Sequent: $\bullet * (A \circ * \bullet B) \vdash * (D \circ E)$

Three ways of presenting the one fact

- ▶ Display Sequent: $\bullet * (A \circ * \bullet B) \vdash * (D \circ E)$
- ▶ Labelled Sequent: $vRw, uRv, u : B, w : D, w : E \vdash v : A$

Three ways of presenting the one fact

- ▶ Display Sequent: $\bullet * (A \circ * \bullet B) \vdash * (D \circ E)$
- ▶ Labelled Sequent: $vRw, uRv, u : B, w : D, w : E \vdash v : A$
- ▶ Delabelled Sequent: $B \vdash \longrightarrow D, E \vdash \longrightarrow \vdash A$

An example delabelling

$$\begin{array}{c}
 \frac{v : A \vdash v : A}{wRv, w : \Box A \vdash v : A} [\Box L] \quad \frac{v : B \vdash v : B}{wRv, w : \Box A \vdash v : A} [\Box L] \\
 \frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \quad \frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \\
 \hline
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \hline
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{wRv, w : \Box A \vdash v : A} [\Box L] \quad \frac{v : B \vdash v : B}{wRv, w : \Box A \vdash v : A} [\Box L] \\
 \frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \quad \frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \\
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \quad wRv, w : \Box A, w : \Box B \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B} [\wedge R] \\
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\square A \vdash \longrightarrow \vdash A} [\Box L] \\
 \hline
 wRv, w : \Box A, w : \Box B \vdash v : A \quad \frac{v : B \vdash v : B}{wRv, w : \Box A \vdash v : A} [\Box L] \\
 \hline
 wRv, w : \Box A, w : \Box B \vdash v : A \quad wRv, w : \Box A, w : \Box B \vdash v : A \quad \frac{[K]}{[K]} \\
 \hline
 wRv, w : \Box A, w : \Box B \vdash v : A \wedge B \quad \frac{[K]}{[\wedge R]} \\
 \hline
 w : \Box A, w : \Box B \vdash w : \Box(A \wedge B) \quad \frac{[\Box R]}{[\wedge R]} \\
 \hline
 w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \\
 \frac{v : B \vdash v : B}{wRv, w : \Box A \vdash v : A} [\Box L] \\
 \frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \quad wRv, w : \Box A, w : \Box B \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B} [\wedge R] \\
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \\
 \hline
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \hline
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

$\frac{B \vdash B}{wRv, w : \Box A \vdash v : A} [\Box L]$
 $\frac{wRv, w : \Box A \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [K]$
 $\frac{wRv, w : \Box A, w : \Box B \vdash v : A}{wRv, w : \Box A, w : \Box B \vdash v : A} [\wedge R]$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \\
 \hline
 \Box A, \Box B \vdash \longrightarrow \vdash A
 \end{array}
 \qquad
 \begin{array}{c}
 \frac{B \vdash B}{\Box B \vdash \longrightarrow \vdash B} [\Box L] \\
 \frac{\Box B \vdash \longrightarrow \vdash B}{wRv, w : \Box A, w : \Box B \vdash v : A} [K] \\
 \hline
 wRv, w : \Box A, w : \Box B \vdash v : A
 \end{array}
 \begin{array}{c}
 \hline
 wRv, w : \Box A, w : \Box B \vdash v : A \wedge B \\
 \hline
 w : \Box A, w : \Box B \vdash w : \Box(A \wedge B) \\
 \hline
 w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)
 \end{array}
 \begin{array}{c}
 [\Box R] \\
 [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \quad \frac{B \vdash B}{\Box B \vdash \longrightarrow \vdash B} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \quad \frac{\Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash B} [K] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \quad \Box A, \Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B} [\wedge R] \\
 \frac{wRv, w : \Box A, w : \Box B \vdash v : A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \quad \frac{B \vdash B}{\Box B \vdash \longrightarrow \vdash B} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \quad \frac{\Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash B} [K] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \quad \Box A, \Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B} [\wedge R] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B}{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)} [\Box R] \\
 \frac{w : \Box A, w : \Box B \vdash w : \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\begin{array}{c}
 \frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L] \quad \frac{B \vdash B}{\Box B \vdash \longrightarrow \vdash B} [\Box L] \\
 \frac{\Box A \vdash \longrightarrow \vdash A}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \quad \frac{\Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash B} [K] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \quad \Box A, \Box B \vdash \longrightarrow \vdash B}{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B} [\wedge R] \\
 \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B}{\Box A, \Box B \vdash \Box(A \wedge B)} [\Box R] \\
 \frac{\Box A, \Box B \vdash \Box(A \wedge B)}{w : \Box A \wedge \Box B \vdash w : \Box(A \wedge B)} [\wedge R]
 \end{array}$$

An example delabelling

$$\frac{\frac{\frac{A \vdash A}{\Box A \vdash \longrightarrow \vdash A} [\Box L]}{\Box A, \Box B \vdash \longrightarrow \vdash A} [K] \quad \frac{\frac{\frac{B \vdash B}{\Box B \vdash \longrightarrow \vdash B} [\Box L]}{\Box A, \Box B \vdash \longrightarrow \vdash B} [K]}{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B} [\wedge R] \\ \frac{\Box A, \Box B \vdash \longrightarrow \vdash A \wedge B}{\Box A, \Box B \vdash \Box(A \wedge B)} [\Box R] \\ \frac{\Box A, \Box B \vdash \Box(A \wedge B)}{\Box A \wedge \Box B \vdash \Box(A \wedge B)} [\wedge R]$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Another example delabelling

$$\begin{array}{c}
 \frac{x : A \vdash x : A}{x : \neg A, x : A \vdash} [\neg L] \\
 \frac{x : \neg A, x : A \vdash}{Ryx, y : \Box \neg A, x : A \vdash} [\Box L] \\
 \frac{Ryx, y : \Box \neg A, x : A \vdash}{Ryx, x : A \vdash y : \neg \Box \neg A} [\neg R] \\
 \frac{Ryx, x : A \vdash y : \neg \Box \neg A}{Rxy, x : A \vdash y : \neg \Box \neg A} [sym] \\
 \frac{Rxy, x : A \vdash y : \neg \Box \neg A}{x : A \vdash x : \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

$$\begin{array}{c}
 \frac{A \vdash A}{\neg A, A \vdash} [\neg L] \\
 \frac{\neg A, A \vdash}{\Box \neg A \vdash \longrightarrow A \vdash} [\Box L] \\
 \frac{\Box \neg A \vdash \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longrightarrow A \vdash} [\neg R] \\
 \frac{\vdash \neg \Box \neg A \longrightarrow A \vdash}{\vdash \neg \Box \neg A \longleftarrow A \vdash} [sym] \\
 \frac{\vdash \neg \Box \neg A \longleftarrow A \vdash}{A \vdash \Box \neg \Box \neg A} [\Box R]
 \end{array}$$

Tree Hypersequent Rules: Modal Operators

$$\frac{\mathcal{H}[X \vdash Y \multimap X', A \vdash Y']}{\mathcal{H}[X, \Box A \vdash Y \multimap X' \vdash Y']} [\Box L]$$

$$\frac{\mathcal{H}[X \vdash Y \multimap \vdash A]}{\mathcal{H}[X \vdash \Box A, Y]} [\Box R]$$

$$\frac{\mathcal{H}[X \vdash Y \multimap A \vdash]}{\mathcal{H}[\Diamond A, X \vdash Y]} [\Diamond L]$$

$$\frac{\mathcal{H}[X \vdash Y \multimap X' \vdash A, Y']}{\mathcal{H}[X \vdash \Diamond A, Y \multimap X' \vdash Y']} [\Diamond R]$$

Forms of Cut

$$\frac{\mathcal{H}[X \vdash A, Y] \quad \mathcal{H}[X, A \vdash Y]}{\mathcal{H}[X \vdash Y]} [Cut^a]$$

Forms of Cut

$$\frac{\mathcal{H}[X \vdash A, Y] \quad \mathcal{H}[X, A \vdash Y]}{\mathcal{H}[X \vdash Y]} [Cut^a]$$

$$\frac{\mathcal{H}[X \vdash A, Y] \quad \mathcal{H}'[X, A \vdash Y]}{(\mathcal{H} \oplus \mathcal{H}') [X \vdash Y]} [Cut^m]$$

Forms of Weakening

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iKL]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iKR]}$$

Forms of Weakening

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iKL]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iKR]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X' \vdash Y' \multimap X \vdash Y]} \text{ [eKL]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X \vdash Y \multimap X' \vdash Y']} \text{ [eKR]}$$

Forms of Weakening

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iKL]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iKR]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X' \vdash Y' \multimap X \vdash Y]} \text{ [eKL]}$$

$$\frac{\mathcal{H}[X \vdash Y]}{\mathcal{H}[X \vdash Y \multimap X' \vdash Y']} \text{ [eKR]}$$

$$\mathcal{H}[X, A \vdash A, Y] \quad \text{[axK]}$$

Forms of Contraction

$$\frac{\mathcal{H}[X, A, A \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iWL]}$$

$$\frac{\mathcal{H}[X \vdash A, A, Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iWR]}$$

Forms of Contraction

$$\frac{\mathcal{H}[X, A, A \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iWL]}$$

$$\frac{\mathcal{H}[X \vdash A, A, Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iWR]}$$

$$\frac{\mathcal{H}[X'' \vdash Y'' \curvearrowright X \vdash Y \curvearrowright X' \vdash Y']}{\mathcal{H}[X' \vdash Y' \curvearrowright X', X'' \vdash X', Y'']} \text{ [eWo]}$$

Forms of Contraction

$$\frac{\mathcal{H}[X, A, A \vdash Y]}{\mathcal{H}[X, A \vdash Y]} \text{ [iWL]}$$

$$\frac{\mathcal{H}[X \vdash A, A, Y]}{\mathcal{H}[X \vdash A, Y]} \text{ [iWR]}$$

$$\frac{\mathcal{H}[X'' \vdash Y'' \curvearrowright X \vdash Y \curvearrowleft X' \vdash Y']}{\mathcal{H}[X' \vdash Y' \curvearrowright X', X'' \vdash X', Y'']} \text{ [eWo]}$$

$$\frac{\mathcal{H}[X'' \vdash Y'' \curvearrowleft X \vdash Y \curvearrowright X' \vdash Y']}{\mathcal{H}[X \vdash Y \curvearrowleft X', X'' \vdash X', Y'']} \text{ [eWi]}$$

Cut Elimination

A cut elimination theorem for tree hypersequent systems is relatively straightforward.

One option is a contraction-free style argument (by Negri and von Plato), following the construction for Labelled Sequent systems.

Another is the Curry–Belnap argument.

Virtues and Vices

	DISPLAY	LABELLED	DELABELLED
<i>Cut-free</i>	+	+	+
<i>Explicit</i>	+	+	+
<i>Systematic</i>	+	+	+
<i>Separation</i>	+	+	+
<i>Subformula</i>	+	+—	+
<i>Nonredundant</i>	—	+—	+
<i>Gentzen-plus</i>	—	+—	+

Virtues and Vices

	DISPLAY	LABELLED	DELABELLED
<i>Cut-free</i>	+	+	+
<i>Explicit</i>	+	+	+
<i>Systematic</i>	+	+	+
<i>Separation</i>	+	+	+
<i>Subformula</i>	+	+—	+
<i>Nonredundant</i>	—	+—	+
<i>Gentzen-plus</i>	—	+—	+

Flat hypersequents (for $S5$),
and *structured* hypersequents
for two-dimensional modal logic.

Display Logic, Labelled Sequents and Hypersequents



NUEL D. BELNAP, JR.

“Display Logic.”

Journal of Philosophical Logic, 11:375–417, 1982.



HEINRICH WANSING

Displaying Modal Logic

Kluwer Academic Publishers, 1998.



SARA NEGRI

“Proof Analysis in Modal Logic.”

Journal of Philosophical Logic, 34:507–544, 2005.



ARNON AVRON

“Using Hypersequents in Proof Systems for Non-Classical Logics.”

Annals of Mathematics and Artificial intelligence, 4:225–248, 1991.

Delabelled Sequents



FRANCESCA POGGIOLESI

Gentzen Calculi for Modal Propositional Logic
Springer, 2011.



FRANCESCA POGGIOLESI AND GREG RESTALL

“Interpreting and Applying Proof Theory for Modal Logic.”
New Waves in Philosophical Logic, ed. Greg Restall and Gillian Russell,
Palgrave MacMillan, 2012.
<http://consequently.org/writing/interp-apply-ptml>

THANK YOU!

<https://consequently.org/class/2016/PTPLA-NASSLLI/>

@consequently / @standefer on Twitter