

Proofs, and what they're good for

Greg Restall



THE UNIVERSITY OF
MELBOURNE

AAP CONFERENCE · 2016

To explain the nature of *proof*,
from the perspective of a *normative
pragmatic account of meaning*, using
the formal tools of *proof theory*.

Outline

Motivation

Background

What Proofs Are

How Proofs Work

MOTIVATION

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx)).$

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade*

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$.

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why?

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an
arbitrary *drink*.

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an
arbitrary *drink*. It's either a *beer* or a *lemonade*.

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade* $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we have the conclusion that some *drink* is a *lemonade*.

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade*
 $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is
a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an
arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we
have the conclusion that some *drink* is a *lemonade*. If we don't have
that conclusion, then that arbitrary *drink* is a *beer*

Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade* $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we have the conclusion that some *drink* is a *lemonade*. If we don't have that conclusion, then that arbitrary *drink* is a *beer*, and so, all the *drinks* are *beers*


Example Proof 1

Every *drink* (in our fridge) is either a *beer* or a *lemonade* $(\forall x)(Dx \supset (Bx \vee Lx))$. So either every *drink* is a *beer* or some *drink* is a *lemonade* $(\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)$. Why? Take an arbitrary *drink*. It's either a *beer* or a *lemonade*. If it's a *lemonade*, we have the conclusion that some *drink* is a *lemonade*. If we don't have that conclusion, then that arbitrary *drink* is a *beer*, and so, all the *drinks* are *beers*, and so, we also have our conclusion.

Example Proof 1 (the formal structure)


$$\begin{array}{c}
 \frac{\frac{\frac{Ba \succ Ba \quad La \succ La}{Ba \vee La \succ Ba, La} \vee L}{Da \succ Da \quad Ba \vee La \succ Ba, La} \supset L}{Da \supset (Ba \vee La), Da \succ Ba, La} \supset L \\
 \frac{Da \supset (Ba \vee La), Da \succ Ba, La}{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, La} \forall L \\
 \frac{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, La \quad Da \succ Da}{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, Da \wedge La} \wedge R \\
 \frac{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, Da \wedge La}{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, (\exists x)(Dx \wedge Lx)} \exists R \\
 \frac{(\forall x)(Dx \supset (Bx \vee Lx)), Da \succ Ba, (\exists x)(Dx \wedge Lx)}{(\forall x)(Dx \supset (Bx \vee Lx)) \succ Da \supset Ba, (\exists x)(Dx \wedge Lx)} \supset R \\
 \frac{(\forall x)(Dx \supset (Bx \vee Lx)) \succ Da \supset Ba, (\exists x)(Dx \wedge Lx)}{(\forall x)(Dx \supset (Bx \vee Lx)) \succ (\forall x)(Dx \supset Bx), (\exists x)(Dx \wedge Lx)} \forall R \\
 \frac{(\forall x)(Dx \supset (Bx \vee Lx)) \succ (\forall x)(Dx \supset Bx), (\exists x)(Dx \wedge Lx)}{(\forall x)(Dx \supset (Bx \vee Lx)) \succ (\forall x)(Dx \supset Bx) \vee (\exists x)(Dx \wedge Lx)} \vee R
 \end{array}$$

Example Proof 2 (The Bridges of Königsberg)





It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*.

Example Proof 2 (The Bridges of Königsberg)





It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*. Why?

Example Proof 2 (The Bridges of Königsberg)





It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*. Why? Any bridge takes you from one landmass (A, B, C, D) to another.

Example Proof 2 (The Bridges of Königsberg)





It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*. Why? Any bridge takes you from one landmass (A, B, C, D) to another. In any *circuit*, you must *leave* a landmass as many times as you *arrive*.

Example Proof 2 (The Bridges of Königsberg)



It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*. Why? Any bridge takes you from one landmass (A, B, C, D) to another. In any *circuit*, you must *leave* a landmass as many times as you *arrive*. So, if you use every bridge exactly once, each landmass must have an even number of bridges entering and exiting it.

Example Proof 2 (The Bridges of Königsberg)



It's not possible to walk a circuit through Königsberg, *crossing each bridge exactly once*. Why? Any bridge takes you from one landmass (A, B, C, D) to another. In any *circuit*, you must *leave* a landmass as many times as you *arrive*. So, if you use every bridge exactly once, each landmass must have an even number of bridges entering and exiting it. Here, each landmass has an *odd* number of bridges, so a circuit is impossible.

Our Focus

Our focus is on *categorical* proofs from premises to a conclusion.

Our Focus

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic*

Our Focus

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic* ($\wedge, \vee, \neg, \supset, \forall, \exists, =$).

Our Focus

Our focus is on *categorical* proofs from premises to a conclusion. In particular, on categorical proofs in *first-order predicate logic* ($\wedge, \vee, \neg, \supset, \forall, \exists, =$).

But what I say here can be extended to proof relying on other concepts.

Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?

Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?

Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?
- ▶ What *grounds* the necessity in the connection between the premises and the conclusion?

Puzzles about proof

- ▶ How can proofs expand our knowledge, when the conclusion is (*in some sense*) already present in the premises?
- ▶ How can we be ignorant of a conclusion which logically follows from what we already know?
- ▶ What *grounds* the necessity in the connection between the premises and the conclusion?
- ▶ (Notice that these are important questions for proofs in first order predicate logic, as much as for proof more generally.)

BACKGROUND

Assertions and Denials

$[X : Y]$

... in a communicative practice

Assertions and denials are moves in a practice.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

I can ‘try on’ assertion or denial hypothetically.

Assertions and denials are moves in a practice.

I can *deny* what you *assert*.

I can *retract* an assertion or a denial.

I can ‘try on’ assertion or denial hypothetically.

They are connected to other speech acts, too, like imperatives, interrogatives, recognitives, observatives, *etc.*

Assertions and denials take a *stand*
(*pro* or *con*) on something.

DENIAL clashes with assertion.
ASSERTION clashes with denial.

Bounds for Positions

- ▶ These norms give rise to **BOUNDS** for positions.

Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY: $[A : A]$ is out of bounds.

Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY: $[A : A]$ is out of bounds.
- ▶ WEAKENING: If $[X : Y]$ is out of bounds, then $[X, A : Y]$ and $[X : A, Y]$ are also out of bounds.

Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY: $[A : A]$ is out of bounds.
- ▶ WEAKENING: If $[X : Y]$ is out of bounds, then $[X, A : Y]$ and $[X : A, Y]$ are also out of bounds.
- ▶ CUT: If $[X, A : Y]$ and $[X : A, Y]$ are out of bounds, then so is $[X : Y]$.

Bounds for Positions

- ▶ These norms give rise to BOUNDS for positions.
- ▶ IDENTITY: $[A : A]$ is out of bounds.
- ▶ WEAKENING: If $[X : Y]$ is out of bounds, then $[X, A : Y]$ and $[X : A, Y]$ are also out of bounds.
- ▶ CUT: If $[X, A : Y]$ and $[X : A, Y]$ are out of bounds, then so is $[X : Y]$.
- ▶ A position that is OUT OF BOUNDS doesn't succeed in taking a stand.

Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

x is a square $=_{df}$ x is a rectangle \wedge
all sides of x are equal in length.

Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

x is a square $=_{df}$ x is a rectangle \wedge
all sides of x are equal in length.

Concepts defined explicitly are *sharply delimited*
(contingent on the definition).

Explicit Definition

Define a concept by showing how you can compose that concept out of more primitive concepts.

x is a square $=_{df}$ x is a rectangle \wedge
all sides of x are equal in length.

Concepts defined explicitly are *sharply delimited*
(contingent on the definition).

Logical concepts are similarly sharply delimited,
but they cannot all be given explicit definitions.

Definition through a rule for use

$[X, A \wedge B : Y]$ is out of bounds

if and only if

$[X, A, B : Y]$ is out of bounds

Definition through a rule for use

$[X, A \wedge B : Y]$ is out of bounds

if and only if

$[X, A, B : Y]$ is out of bounds

$$\frac{X, A, B \vdash Y}{X, A \wedge B \vdash Y} \wedge Df$$

What about when to *deny* a conjunction?

$$\frac{\frac{\frac{}{A \wedge B \vdash A \wedge B} Id}{X \vdash B, Y \quad A, B \vdash A \wedge B} \wedge Df}{\frac{X \vdash A, Y \quad X, A \vdash A \wedge B, Y}{X \vdash A \wedge B, Y} Cut} Cut$$

What about when to *deny* a conjunction?

$$\frac{\frac{\frac{}{A \wedge B \vdash A \wedge B} Id}{A, B \vdash A \wedge B} \wedge Df}{\frac{X \vdash B, Y \quad A, B \vdash A \wedge B}{X, A \vdash A \wedge B, Y} Cut} \frac{X \vdash A, Y \quad X, A \vdash A \wedge B, Y}{X \vdash A \wedge B, Y} Cut$$

So, we have

$$\frac{X \vdash A, Y \quad X \vdash B, Y}{X \vdash A \wedge B, Y} \wedge R$$

Definitions for other logical concepts

$$\frac{X \vdash A, Y}{X, \neg A \vdash Y} \neg Df$$

$$\frac{X, A \vdash B, Y}{X \vdash A \supset B, Y} \supset Df$$

$$\frac{X \vdash A, B, Y}{X \vdash A \vee B, Y} \vee Df$$

Definitions for other logical concepts

$$\frac{X \vdash A, Y}{X, \neg A \vdash Y} \neg Df$$

$$\frac{X, A \vdash B, Y}{X \vdash A \supset B, Y} \supset Df$$

$$\frac{X \vdash A, B, Y}{X \vdash A \vee B, Y} \vee Df$$

$$\frac{X \vdash Fa, Y}{X \vdash (\forall x)Fx, Y} \forall Df$$

$$\frac{X, Fa \vdash Y}{X, (\exists x)Fx \vdash Y} \forall Df$$

$$\frac{X, Gb \vdash Gc, Y}{X \vdash b = c, Y} \forall Df$$

(Where a and G are not present in X and Y .)

Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)

Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)

Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)

Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ▶ Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)

Concepts defined in this way...

- ▶ Are *uniquely defined*. (If you and I use the same rule, we define the same concept.)
- ▶ Are *conservatively extending*. (Adding a logical concept to your vocabulary in this way doesn't constrain the bounds in the original language.)
- ▶ *Play useful dialogical roles*. (You can do things with these concepts that you cannot do without. Denying a conjunction does something different to denying the conjuncts.)
- ▶ Are *subject matter neutral*. (They work wherever you assert and deny—and have singular terms and predicates.)
- ▶ In Brandom's terms, they *make explicit* some of what was implicit in the practice of assertion and denial.

WHAT PROOFS ARE

A Tiny Proof

If it's Friday, I'm in Melbourne.

It's Friday.

Therefore, I'm in Melbourne.

A Tiny Proof

If it's Friday, I'm in Melbourne.

It's Friday.

Therefore, I'm in Melbourne.

$$\frac{\frac{}{A \supset B \vdash A \supset B} Id}{A \supset B, A \vdash B} \supset Df$$

A Tiny Proof

If it's Friday, I'm in Melbourne.

It's Friday.

Therefore, I'm in Melbourne.

$$\frac{\frac{}{A \supset B \vdash A \supset B} Id}{A \supset B, A \vdash B} \supset Df$$

[It's Friday \supset I'm in Melbourne, It's Friday : I'm in Melbourne]

(This is out of bounds.)

The Undeniable

Take a context in which I've asserted
it's Friday \supset I'm in Melbourne
and I've asserted *it's Friday*,
then *I'm in Melbourne* is undeniable.

The Undeniable

Take a context in which I've asserted
it's Friday \supset I'm in Melbourne
and I've asserted *it's Friday*,
then *I'm in Melbourne* is undeniable.

Adding the *assertion* makes explicit
what was *implicit* before that assertion.

The Undeniable

Take a context in which I've asserted
it's Friday \supset I'm in Melbourne
and I've asserted *it's Friday*,
then *I'm in Melbourne* is undeniable.

Adding the *assertion* makes explicit
what was *implicit* before that assertion.

The *stance* (*pro* or *con*)
on *I'm in Melbourne* was already made.

A *proof* of $X \vdash Y$ shows that the position $[X : Y]$ is out of bounds, by way of the defining rules for the concepts involved in the proof.

A *proof* of $X \vdash Y$ shows that the position $[X : Y]$ is out of bounds, by way of the defining rules for the concepts involved in the proof.

In this sense, proofs are *analytic*.

They apply, given the definitions, independently of the positions taken by those giving the proof.

What Proofs Prove

A proof of $A, B \vdash C, D$ can be seen
as a *proof* of C from $[A, B : D]$,
and a *refutation* of A from $[B : C, D]$,
and *more*.

HOW PROOFS WORK

Observation o: Proofs are *analytic*

These proofs are grounded in the *rules*
defining the concepts used in them.

Observation 1: *Specification* outstrips *Recognition*

Our ability to *specify* concepts and consequence
far outstrips our ability to *recognise* that consequence.

Peano Arithmetic and Goldbach's Conjecture

SUCCESSOR AXIOMS:

PA1: $\forall x \forall y (x' = y' \supset x = y)$;

PA2: $\forall x (0 \neq x')$.

ADDITION AXIOMS:

PA3: $\forall x (x + 0 = x)$;

PA4: $\forall x (x + y' = (x + y)')$.

MULTIPLICATION AXIOMS:

PA5: $\forall x (x \times 0 = 0)$;

PA6: $\forall x \forall y (x \times y' = (x \times y) + x)$.

INDUCTION SCHEME:

PA7: $(\phi(0) \wedge \forall x (\phi(x) \supset \phi(x')))) \supset \forall x \phi(x)$.

GOLDBACH'S CONJECTURE:

GC: $\forall x \exists y \exists z (\text{Prime } y \wedge \text{Prime } z \wedge 0'' \times x = y + z)$

Observation 1: *Specification* outstrips *Recognition*

Is $[PA : GC]$ out of bounds?

Observation 1: *Specification* outstrips *Recognition*

Is $[PA : GC]$ out of bounds?

We have no idea.

Observation 1: *Specification* outstrips *Recognition*

Is $[PA : GC]$ out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

Observation 1: *Specification* outstrips *Recognition*

Is $[PA : GC]$ out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

Our concepts are rich and expressive.
We can say things whose significance
we continue to work out.

Is $[PA : GC]$ out of bounds?

We have no idea.

This is not a *bug*. It's a *feature*.

Our concepts are rich and expressive.
We can say things whose significance
we continue to work out.

Verifying a putative proof is straightforward.
Checking that something *has* a proof is not so easy.

Are we logically omniscient?

Suppose that $PA \vdash GC$
(but we don't possess that proof)
and that we *know* PA .

Do we know GC ?

In a weak sense of 'know', *yes*, we do know CC

- ▶ It's a logical consequence of what we know.

In a weak sense of 'know', *yes*, we do know CC

- ▶ It's a logical consequence of what we know.
- ▶ It is implicitly present in what we already know.

In a weak sense of 'know', *yes*, we do know GC

- ▶ It's a logical consequence of what we know.
- ▶ It is implicitly present in what we already know.
- ▶ There is no epistemic possibility (no circumstance consistent with our knowledge) that leaves GC out.

In a not-so-weak sense, we don't know GC

- ▶ Do we *believe* GC?

In a not-so-weak sense, we don't know GC

- ▶ Do we *believe* GC?
- ▶ If we believed it, do we believe it *in the right way*?

In a not-so-weak sense, we don't know GC

- ▶ Do we *believe* GC?
- ▶ If we believed it, do we believe it *in the right way*?
- ▶ There is evidence for GC (its proof from PA, for example), but if that evidence plays no role in our belief...

Observation 2: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.

Observation 2: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.
- ▶ However, given plausible (minimal) assumptions concerning T , we can show that (*for example*) if $A, B \vdash C$ then $T\langle A \rangle, T\langle B \rangle \vdash T\langle C \rangle$.

Observation 2: Proofs Preserve Truth

- ▶ The account of consequence does not use the concept of *truth*.
- ▶ However, given plausible (minimal) assumptions concerning T , we can show that (*for example*) if $A, B \vdash C$ then $T\langle A \rangle, T\langle B \rangle \vdash T\langle C \rangle$.
- ▶ This *follows from* the concepts of consequence and truth.

Observation 3: Proofs Can Preserve Warrant

- ▶ The account of consequence does not use the concept of *warrant*.

Observation 3: Proofs Can Preserve Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (*for example*) if p is a proof for $A, B \vdash C$ then $x : A, y : B \vdash p(x, y) : C$.

Observation 3: Proofs Can Preserve Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (*for example*) if p is a proof for $A, B \vdash C$ then $x : A, y : B \vdash p(x, y) : C$.
- ▶ Here, p transforms warrants for the premises into warrant for the conclusion.

Observation 3: Proofs Can Preserve Warrant

- ▶ The account of consequence does not use the concept of *warrant*.
- ▶ However, given plausible (less minimal) assumptions concerning warrant, we can show that (*for example*) if p is a proof for $A, B \vdash C$ then $x : A, y : B \vdash p(x, y) : C$.
- ▶ Here, p transforms warrants for the premises into warrant for the conclusion.
- ▶ This works only for *categorical, conclusive* warrants (*grounds*), not for *defeasible* warrants.

A Caveat on Defeasible Warrants

Consider the “Lottery Paradox.”

A Caveat on Defeasible Warrants

Consider the “Lottery Paradox.”

$$\begin{aligned} & [(\exists x)(Tx \wedge Wx), \\ & (\forall x)(Tx \equiv (x = t_1 \vee x = t_2 \vee \dots \vee x = t_{1\,000\,000})) \\ & \quad : Wt_1, Wt_2, \dots, Wt_{1\,000\,000}] \end{aligned}$$

A Caveat on Defeasible Warrants

Consider the “Lottery Paradox.”

$$\begin{aligned} & [(\exists x)(Tx \wedge Wx), \\ & (\forall x)(Tx \equiv (x = t_1 \vee x = t_2 \vee \dots \vee x = t_{1\,000\,000})) \\ & \qquad : Wt_1, Wt_2, \dots, Wt_{1\,000\,000}] \end{aligned}$$

We have a very high degree of confidence in each part.

Each component is highly probable.

But the whole position is out of bounds.

Observation 4: Achilles and the Tortoise

“Well, now, let’s take a little bit of the argument in that First Proposition—just *two* steps, and the conclusion drawn from them. Kindly enter them in your note-book. And in order to refer to them conveniently, let’s call them *A*, *B*, and *Z* :—

(*A*) Things that are equal to the same are equal to each other.

(*B*) The two sides of this Triangle are things that are equal to the same.

(*Z*) The two sides of this Triangle are equal to each other.

Readers of Euclid will grant, I suppose, that *Z* follows logically from *A* and *B*, so that any one who accepts *A* and *B* as true, *must* accept *Z* as true?”

“Undoubtedly! The youngest child in a High School—as soon as High Schools are invented, which will not be till some two thousand years later—will grant *that*.”

“And if some reader had *not* yet accepted *A* and *B* as true, he might still accept the *sequence* as a *valid* one, I suppose?”

Observation 4: Achilles and the Tortoise

“No doubt such a reader might exist. He might say ‘I accept as true the Hypothetical Proposition that, *if* A and B be true, Z must be true; but, I *don't* accept A and B as true.’ Such a reader would do wisely in abandoning Euclid, and taking to football.”

“And might there not *also* be some reader who would say ‘I accept A and B as true, but I *don't* accept the Hypothetical’?”

“Certainly there might. *He*, also, had better take to football.”

“And *neither* of these readers,” the Tortoise continued, “is *as yet* under any logical necessity to accept Z as true?”

“Quite so,” Achilles assented.

“Well, now, I want you to consider *me* as a reader of the *second* kind, and to force me, logically, to accept Z as true.”

“A tortoise playing football would be—” Achilles was beginning

“—an anomaly, of course,” the Tortoise hastily interrupted. “Don’t wander from the point. Let’s have Z first, and football afterwards!”

“I’m to force you to accept Z , am I?” Achilles said musingly. “And your present position is that you accept A and B , but you *don't* accept the Hypothetical—”

“Let’s call it C ,” said the Tortoise.

“—but you *don't* accept

(C) If A and B are true, Z must be true.”

“That is my present position,” said the Tortoise.

“Then I must ask you to accept C .”

Our Analysis

$$A, B \vdash Z$$

Our Analysis

$A, B \vdash Z$

or

$A, A \supset Z \vdash Z$

Our Analysis

$$A, B \vdash Z$$

or

$$A, A \supset Z \vdash Z$$

This *doesn't* mean when I accept A
and I accept $A \supset Z$,
I ought to also accept Z .

Our Analysis

$$A, B \vdash Z$$

or

$$A, A \supset Z \vdash Z$$

This *doesn't* mean when I accept A
and I accept $A \supset Z$,
I ought to also accept Z .

However, if I assert A and $A \supset Z$ then Z is *undeniable*.

If I assert A and *if A then Z* and *deny Z* ,
then I am using ‘*if...then*’ in a way that
deviates from the defining rule for \supset ,
or I am explicitly contradicting myself.

If I assert A and *if* A *then* Z and *deny* Z ,
then I am using ‘*if...then*’ in a way that
deviates from the defining rule for \supset ,
or I am explicitly contradicting myself.

$$\frac{A \supset B \vdash A \supset B}{A \supset B, A \vdash B} \supset Df$$

An account of the logical concepts given in terms
of defining rules governing assertions and denials

An account of the logical concepts given in terms of defining rules governing assertions and denials helps explain how (*first order predicate logic*) proof works,

An account of the logical concepts given in terms of defining rules governing assertions and denials helps explain how (*first order predicate logic*) proof works, how possessing a proof can expand our knowledge, while proofs make explicit what is implicit in what we know.

THANK YOU!

[https://consequently.org/presentation/2016/
proofs-and-what-theyre-good-for-AAP/](https://consequently.org/presentation/2016/proofs-and-what-theyre-good-for-AAP/)

@consequently on Twitter