

CROP DATA SHEET · ROW CROP

Corn

Holds yield through heat and drought, then keeps building in normal seasons.

+34% YIELD UNDER EXTREME HEAT
119 DAYS ABOVE 86°F · P=0.015

+5-14%

ACROSS TRIALS

Yield gain range across independent replicated corn trials.

+12.4%

NO-IRRIGATION DROUGHT

Result under dryland conditions (p=0.06).


Cooler

CANOPY & STEM

Treated corn held lower canopy and stem temperatures under heat.



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Corn is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
2015	Arizona	Seed treatment	+34% (heat, $p=0.015$)
2016	Indiana	Seed treatment	+14.2% ($p=0.006$)
2020	Missouri	Seed treatment	+12.4% (drought, $p=0.06$)
2022	Iowa	Two-step program	+9.5% ($p=0.004$)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for corn at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • LEGUME • ROW CROP

Soybeans

More beans per acre, with stronger stands and fewer synthetic inputs.

+12.5% YIELD VS. UNTREATED
REPLICATED TRIAL • P=0.05

+8-13%

ACROSS TRIALS

Yield gain range across independent soybean trials.

30-50%


LESS NITROGEN

Fixes and frees more of its own nitrogen (Harman 2011).


Improved

ROOTS & MODULATION

Stronger root and nodule development for better nutrient reach.



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Soybeans is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
2023	Missouri	Seed treatment	+12.5% (p=0.05)
2020	Michigan	Seed treatment	+8.7% (p=0.06)
2021	Iowa	Seed treatment	+8.4% (p=0.08)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for soybeans at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · CEREAL · ROW CROP

Wheat

More grain and better protein, with less nitrogen.

+11.3%

YIELD, SEED-TREATED

REPLICATED TRIAL · P=0.0001 · +1.4% PROTEIN

+9-11%

ACROSS TRIALS

Yield gain range across independent wheat trials.

+1.4%

GRAIN PROTEIN


Higher protein alongside the yield gain.

+8.6%

STRAW YIELD

More whole-plant value from the same field.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Wheat is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
2016	Indiana	Seed treatment	+11.3% (p=0.0001)
2020	Indiana	Seed treatment	+9.8% (p=0.01)
2019	Wisconsin	Seed + foliar	+9.0% (p=0.004)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for wheat at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · CEREAL · ROW CROP

Barley

Stronger stands and steady yield through a dry, hot finish.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Barley has not been through a Symm-specific trial; the figures below are the general cross-crop results from independent peer-reviewed research, pointing to yield through a dry, hot finish and lower nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not barley-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for barley at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symmbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • OILSEED • ROW CROP

Canola

Stronger stands and steadier yield through heat and drought.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Canola has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in canola: carrying yield through heat and water stress, and improving nutrient-use efficiency.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not canola-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for canola at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · CEREAL

Rice

Holds yield through a hot, dry season, with less fertilizer.

+13–30% **YIELD, STRESS SEASON**
INDEPENDENT TRIALS, DRY YEAR

+13–30%

STRESS SEASON

Gain range in a hot, dry season vs. untreated.

30–50%


LESS NITROGEN

Matched yield at reduced N (Harman 2011).


Improved

TILLERING & VIGOR

More tillers and stronger early vigor.



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Rice is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
2023	Texas	Seed treatment	+29.7% (stress season)
2024	Texas	Foliar pass	+4.9%
Multi-yr	Sub-Saharan	On-farm	+28-75%, high win rate

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for rice at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · CEREAL

Millet

A hardy dryland grain made hardier, with less fertilizer.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Millet has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in millet: extending its natural drought and heat tolerance under low-input, dryland conditions.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not millet-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for millet at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symmbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · FORAGE

Alfalfa

Yield gains every cutting, with no drop in relative feed value.

+25% YIELD, LATE-SEASON FOLIAR
NO DECREASE IN RELATIVE FEED VALUE

+7-12%

SEED & IN-FURROW

Gain range for in-furrow and seed-treated cuttings.

+13-25%

FOLIAR, APR-AUG


Gain range for foliar application through the season.

No loss

FEED VALUE

Relative feed value held steady across cuttings.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Alfalfa is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Multi-yr	Temperate	Seed treatment	+9.9–10.8% / cutting (p<0.05)
Multi-yr	Arid	Foliar pass	+13.6–25.0% (p<0.05)
Multi-yr	Temperate / arid	In-furrow	+7.0–12.3% (p<0.05)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for alfalfa at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · FORAGE

Pasture Grass

More forage off every cutting, on less synthetic input.

+6%

FRESH-WEIGHT YIELD

ACROSS 20 CUTTINGS VS. UNTREATED

+6%

SEASON YIELD

Fresh-weight gain across a full cutting season.

30-50%

LESS NITROGEN


Matched yield at reduced N (Harman 2011).

Improved

ROOT ZONE

Deeper roots for better drought carry-through.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Pasture Grass is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Multi-yr	Temperate	Seed / foliar	+6% fresh weight (20 cuttings)
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for pasture grass at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · FIBER & GRAIN

Hemp

More grain and far more straw, with less synthetic input.


+35% **STRAW YIELD**
FOLIAR PASS VS. UNTREATED

+18.5%
GRAIN YIELD
Foliar application vs. untreated.

+35.2%
STRAW YIELD
More fiber tonnage from the same field.

30–50%
LESS NITROGEN
Matched yield at reduced N (Harman 2011).

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress
Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth
Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone
A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Hemp is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
2021	Temperate	Foliar pass	+18.5% grain, +35.2% straw
2021	Temperate	Seed treatment	+15.1% grain, +31.9% straw

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for hemp at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · FIBER · ROW CROP

Cotton

Stronger plants and better fiber quality, with less nitrogen.

Higher

LINT VALUE

FIBER QUALITY VS. UNTREATED

Higher

LINT VALUE

Better fiber quality than untreated controls.

30–50%

LESS NITROGEN


Matched yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger roots for heat and drought carry-through.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Cotton is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Multi-yr	Australia	Foliar / seed	Higher lint value vs. control
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for cotton at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • LEGUME • SPECIALTY

Guar

A tough, drought-hardy crop made tougher, with less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Guar has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in guar: extending its natural drought and heat tolerance, and fixing and freeing more of its own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not guar-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for guar at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@syμβiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • LEGUME

Dry Beans

Stronger stands and more pods, fixing more of their own nitrogen.

+8-10% TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Dry Beans has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in dry beans: carrying pod set through heat and drought and freeing more of their own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not dry beans-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for dry beans at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEGUME

Field Peas

Better emergence and steadier yield, with less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Field Peas has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in field peas: improving establishment through cool, wet or dry starts and fixing more of their own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not field peas-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for field peas at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • LEGUME

Lentils

Tougher stands through heat and drought, on less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Lentils has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in lentils: carrying yield through heat and water stress and fixing more of their own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not lentils-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for lentils at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • LEGUME

Mung Bean

A resilient pulse made more resilient under heat and drought.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Mung Bean has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in mung bean: carrying pod set through heat and drought and fixing more of its own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not mung bean-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for mung bean at 100 acres with typical yield and price.

Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEGUME

Peanut

Stronger stands and better fill through heat and drought.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Peanut has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in peanut: supporting root and peg development through heat and water stress, and fixing more of its own nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not peanut-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for peanut at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEGUME · VEGETABLE

Sugar Snap Peas

Faster emergence and a bigger harvest, with less nitrogen.

+34.9%

HARVEST

GERMINATED 46% FASTER VS. UNTREATED

+34.9%

HARVEST

More harvest vs. untreated.

46% faster

GERMINATION

Quicker, more even emergence.


30–50%

LESS NITROGEN

Fixes and frees more of its own N (Harman 2011).



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Sugar Snap Peas is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Seed treatment	+34.9% harvest, germinated 46% faster

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for sugar snap peas at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE

Tomatoes

More fruit and bigger fruit, with less nitrogen.

+8.4%

HARVEST WEIGHT

+5.4% AVERAGE WEIGHT PER FRUIT

+8.4%

HARVEST WEIGHT

More total harvest vs. untreated.

+5.4%


PER FRUIT

Heavier average fruit.


30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Tomatoes is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Container	Replicated	Two-step program	+8.4% weight, +5.4% per fruit

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for tomatoes at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symmbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE

Peppers

Many more peppers per plant, with less synthetic input.

+91% **TOTAL WEIGHT**
+64% FRUIT COUNT VS. UNTREATED

+91%

TOTAL WEIGHT

More pepper weight per plant vs. untreated.

+64%


FRUIT COUNT

More fruit set per plant.


30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Peppers is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	+91% weight, +64% count

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for peppers at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE

Eggplant

More resilient plants and better fruit set under heat.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Eggplant has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in eggplant: carrying fruit set through heat and drought, and improving nutrient-use efficiency.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not eggplant-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for eggplant at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE

Cucumbers

A far bigger season-long harvest, from healthier vines.

+148.8% SEASON HARVEST WEIGHT
REPLICATED TRIAL · $P < 0.006$

+148.8%

SEASON HARVEST

Cumulative harvest weight across the season ($p < 0.006$).

30-50%

LESS NITROGEN


Matched yield at reduced N (Harman 2011).

Improved

VIGOR & SET

Stronger vines and more fruit set under heat.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Cucumbers is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Container	Replicated	Two-step program	+148.8% season harvest (p<0.006)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for cucumbers at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · CUCURBIT

Squash

A bigger harvest across varieties, on less synthetic input.

+47.9% HARVEST, SUMMER SQUASH
BEST-VARIETY RESULT VS. UNTREATED

+12.8–46.8%

WINTER SQUASH

Gain range across six varieties.

+47.9%

SUMMER SQUASH


Best-variety harvest gain.

30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Squash is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	Winter +12.8–46.8% (6 varieties)
Field	Replicated	Two-step program	Summer squash +47.9%

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for squash at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE

Okra

A heat-loving crop made steadier through drought, on less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Okra has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in okra: carrying pod set through heat and drought and improving nutrient-use efficiency.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not okra-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for okra at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · ALLIUM

Onions

Reliable emergence even from heat-stressed, stored seed.

0→50%

GERMINATION RECOVERY

SEED HELD ONE YEAR AT 35°C

0→50%

GERMINATION

Emergence from seed held a year at 35°C vs. 0% untreated.

30-50%

LESS NITROGEN


Matched yield at reduced N (Harman 2011).

Improved

VIGOR

Stronger early stands and establishment.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Onions is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Trial	Heat-stressed seed	Seed treatment	Germination 0% → 50%

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for onions at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · ALLIUM

Garlic

Bigger bulbs and more of them, with less synthetic input.

+27% OVERALL YIELD

BULBS 8.2% LARGER VS. UNTREATED

+27%

YIELD

Overall yield gain vs. untreated.

+8.2%

BULB SIZE


Larger individual bulbs.

30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Garlic is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	+27% yield, +8.2% bulb size

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for garlic at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · ROOT

Carrot

Better stands and more uniform roots, with less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Carrot has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in carrot: improving establishment and root uniformity through heat and drought, on less nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not carrot-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for carrot at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · PERENNIAL

Asparagus

Stronger crowns and steadier spears, on less synthetic input.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Asparagus has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in asparagus: supporting crown vigor and establishment through heat and drought, and reducing nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not asparagus-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for asparagus at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · BRASSICA

Broccoli

Tighter heads and steadier yield through heat stress.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Broccoli has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in broccoli: carrying head quality through heat and water stress, and cutting nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not broccoli-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for broccoli at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · VEGETABLE · BRASSICA

Cabbage

Heavier heads and a bigger harvest, on less synthetic input.

+24.8%

HARVEST WEIGHT

FIELD TRIAL VS. UNTREATED

+24.8%

HARVEST WEIGHT

Heavier harvest vs. untreated.

30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).


Improved

VIGOR

Stronger, darker, more uniform heads.



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Cabbage is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	+24.8% harvest weight

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for cabbage at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEAFY VEGETABLE · BRASSICA

Kale

A much bigger season-long cut, from healthier plants.

+111%

SEASON HARVEST WEIGHT

CUMULATIVE ACROSS THE HARVEST WINDOW

+111%

SEASON HARVEST

Cumulative harvest weight across the season.

30-50%

LESS NITROGEN


Matched yield at reduced N (Harman 2011).

Improved

VIGOR & QUALITY

Darker, more robust leaves.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Kale is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	+111% season harvest weight

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for kale at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEAFY VEGETABLE

Lettuce

A bigger, more nutrient-dense harvest, with less nitrogen.

+14.9% **HARVEST WEIGHT**
FIELD TRIAL VS. UNTREATED

+14.9%

HARVEST WEIGHT

More harvest vs. untreated.

Higher

NUTRIENT DENSITY


More N, P, K and calcium in treated leaves ($p < 0.001$).

30–50%

LESS NITROGEN

Matched yield at reduced N (Harman 2011).

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Lettuce is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Field	Replicated	Two-step program	+14.9% harvest weight
Independent	Testing	Nutrient uptake	Higher N/P/K/Ca, protein, dry matter (p<0.001)

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for lettuce at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEAFY VEGETABLE

Leafy Greens

A bigger, more nutrient-dense harvest, with less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Leafy Greens has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in leafy greens: improving establishment and quality through heat and water stress, and raising nutrient density.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not leafy greens-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for leafy greens at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · LEAFY VEGETABLE

Spinach

Better germination and stronger seedlings, even after drought.

+27% GERMINATION
SEED-TREATED VS. UNTREATED

+27%

GERMINATION

More seed emerged vs. untreated.

85.7→95.6%

VIABILITY

Higher seedling viability after drought.


+53.7%

SEEDLING BIOMASS

More biomass after one week of drought.



WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Spinach is part of the verified trial record. Figures are from independent replicated trials with cooperator identities withheld, shown as ranges, never a guarantee for a given field.

YEAR	REGION	APPLICATION	RESULT
Container	Replicated	Seed treatment	+27% germination (36/50 vs 46/50)
Container	Drought	Seed treatment	Viability 85.7→95.6%, +53.7% biomass

Independent contract-research trials; cooperator names withheld. Figures are verified ranges, never a guarantee for a given field.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for spinach at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@symbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Apple

Stronger, more resilient trees and better fruit, on less synthetic nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Apple has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in apple: holding fruit set and quality through heat and drought, supporting young-tree establishment, and cutting nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not apple-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for apple at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@syμβiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Cherry

Healthier trees and better fruit set through heat and drought.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Cherry has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in cherry: carrying fruit set and quality through heat spikes and water stress, and reducing nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not cherry-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for cherry at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@syμβiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Peach

More resilient trees and better-sized fruit, with fewer inputs.

+8-10% TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Peach has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in peach: protecting fruit sizing through summer heat and drought, and easing nitrogen demand.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not peach-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for peach at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Avocado

Stronger roots and steadier trees under heat and water stress.

+8-10% TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Avocado has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in avocado: supporting root health and tree vigor through heat, drought and salinity, on less nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not avocado-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for avocado at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Citrus

Tree vigor and root health under sustained stress, with less nitrogen.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Citrus has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in citrus: supporting tree vigor and root health under heat, drought and salinity, and trimming nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not citrus-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for citrus at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD CROP

Olive

Resilient trees and steady set through hot, dry seasons.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Olive has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in olive: carrying trees through heat and drought and improving nutrient-use efficiency.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not olive-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for olive at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • ORCHARD • VINE

Kiwifruit

Stronger vines and better fruit through heat and water stress.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Kiwifruit has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in kiwifruit: supporting vine vigor and fruit quality through heat and drought, on less nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not kiwifruit-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for kiwifruit at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · ORCHARD · VINE

Grape

A stronger canopy and better fruit quality, with less synthetic input.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Grape has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in grape: supporting vine vigor and fruit quality through heat and drought, and cutting nitrogen cost.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not grape-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for grape at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET • BERRY

Blueberries

Stronger plants and better fruit set, on less synthetic input.

+8-10% TYPICAL YIELD GAIN

CROSS-CROP AVERAGE • UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Blueberries has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in blueberries: carrying fruit set through heat and drought and improving nutrient uptake in acidic soils.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not blueberries-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for blueberries at 100 acres with typical yield and price.

Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · BERRY

Blackberry

Healthier canes and better fruit set through heat and drought.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Blackberry has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in blackberry: supporting cane vigor and fruit set through heat and water stress, on less nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not blackberry-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for blackberry at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · BERRY

Raspberry

Stronger canes and steadier fruit set, with fewer inputs.

+8-10% TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Raspberry has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in raspberry: protecting fruit set through heat and drought and supporting root health.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not raspberry-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for raspberry at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · BERRY

Cranberry

Healthier vines and steadier set in demanding bog conditions.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Cranberry has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in cranberry: supporting vine vigor and fruit set through stress, and improving nutrient-use efficiency.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not cranberry-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for cranberry at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@syμβiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)

CROP DATA SHEET · SPECIALTY · VINE

Hops

Stronger bines and steadier cones through a hot season.

+8-10%

TYPICAL YIELD GAIN

CROSS-CROP AVERAGE · UP TO ~14% UNDER STRESS

+8-10%

YIELD

Typical gain vs. untreated; greater as heat or drought stress rises.

30-50%

LESS NITROGEN


Matched full-rate yield at reduced N (Harman 2011).

Improved

ROOTS & VIGOR

Stronger root development and establishment across trials.

WHEN TO APPLY



Apply by **seed treatment**, **in-furrow** at planting, or as a **foliar pass**. Earlier is better, but a foliar pass well into the **vegetative stage** still delivers a real benefit — it is never too late to start.

HOW THE PROGRAM WORKS

Holds up under stress

Living inputs help the plant manage heat, drought and salinity, so more of the season goes into the crop.

More growth

Symbiotic signalling lifts growth rate and nutrient-use efficiency across the canopy.

Healthier root zone

A richer root environment frees up soil nutrients, so the crop leans less on synthetic NPK.

FIELD-TRIAL EVIDENCE

Hops has not yet been through a Symm-specific field trial, so the figures below are the general cross-crop results from independent peer-reviewed research. They point to where the program helps most in hops: supporting bine vigor and cone development through heat and drought, on less nitrogen.

YEAR	REGION	APPLICATION	RESULT
—	Peer-reviewed (Cornell / Harman)	Trichoderma research	+8-10% yield, improved roots
2011	Peer-reviewed · Harman	Nitrogen study	30-50% N reduction, no yield loss

General cross-crop figures from independent peer-reviewed research, shown as ranges; not hops-specific trial results and never a guarantee for a given field. Yield gain is typically +8 to 10%, up to ~14% under abiotic stress.



See your numbers.

Scan or tap to open our benefit calculator, pre-filled for hops at 100 acres with typical yield and price. Adjust to your farm and see your projected net gain and return.

[ESTIMATE YOUR BENEFIT →](#)

✓ CHEMICAL-FREE

✓ PESTICIDE-FREE

✓ OMRI LISTED

✓ NON-GMO

Build your program for the season.

advisors@sympbiotic.com · (845) 393-9539 · Symm 2026

[CONTACT A REP →](#)