

Analysis of Categorical Data

Dr. Supaporn Erjongmanee

Department of Computer Engineering
Kasetsart University
fengspe@ku.ac.th

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 1



Department of Computer Engineering
Kasetsart University

1

Outline

- Analysis of Categorical Data

- Introduction

- Homogeneity test

- Independence test

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 2



Department of Computer Engineering
Kasetsart University

2

Introduction

- A study of data in categories
- Case: Population I of interest; Each population is separated into J categories
 - Example: 3 department stores vs. 5 payment methods (cash, check, store credit card, Visa, Mastercard)
- Homogeneity (Hypothesis) Test
 - Proportions of all categories in each population are the same

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 3



Department of Computer Engineering
Kasetsart University

3

Introduction (cont.)

- In general, data are put in the table
- Let n_{ij} = number of samples in (i,j) category
- Table contains $\{n_{ij}\}$'s is called two-way contingency table

	1	2	...	j	...	J
1	n_{11}	n_{12}	...	n_{1j}	...	n_{1J}
2	n_{21}					
...	...					
i	n_{i1}			n_{ij}		
...	...					
I	n_{I1}					n_{IJ}

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 4



Department of Computer Engineering
Kasetsart University

4

Outline

- Analysis of Categorical Data
 - Introduction
 - Homogeneity test
 - Independence test

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 5



Department of Computer Engineering
Kasetsart University

5

Homogeneity Test

- Population I of interest; Each population is separated into J categories

- Let
 - n_{ij} = number of samples in (i,j) category
 - n_j = number of samples in j category = $\sum_i n_{ij}$
 - n_i = number of samples in i population = $\sum_j n_{ij}$
 - n = number of all samples = $\sum_i \sum_j n_{ij}$
 - p_{ij} = proportions of samples in (i,j) category
- Hypothesis test
 - Null hypothesis (H_0): $p_{1j} = p_{2j} = \dots = p_{Ij}$
 - Proportion of samples in j category for each population is the same
 - Alternative hypothesis (H_a): H_0 is not true

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 6



Department of Computer Engineering
Kasetsart University

6

Homogeneity Test (cont.)

P(samples in category j)


- Let \hat{e}_{ij} = expected number of samples = $n_i p_j = n_i \frac{n_j}{n}$

- Test statistic

$$\chi^2 = \sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$$

- Rejection region

$$\chi^2 \geq \chi_{\alpha, (I-1)(J-1)}^2$$



- In each row i , there are J cells but $n_i = \sum_j n_{ij}$ is fixed. Hence, d.f. per row = $J-1$. There are I rows. Thus, sum of d.f. from all rows = $I(J-1)$
- In addition, we estimate p_1, p_2, \dots, p_J with $\sum_i p_i = 1$. There are $J-1$ parameters to estimate.
- At the end, resulting d.f. = $I(J-1) - (J-1) = (I-1)(J-1)$

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 7



Department of Computer Engineering
Kasetsart University

7

Example

- A can food company have three product sizes; each size is produced at different production lines
- Test in nonconformity of cans at significance level 0.5
 - Blemish, Crack, Improper pull tab location, Missing pull tab, Others

		Nonconformity					Sample size
		Blemish	Crack	Location	Missing	Others	
Production line	1	34	65	17	21	13	150
	2	23	52	25	19	6	125
	3	32	28	16	14	10	100
Total		89	145	58	54	29	375

n_i

n_j

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 8



Department of Computer Engineering
Kasetsart University

8

Example (cont.)

- Hypothesis
 - H_0 : All production lines are homogeneous in term of nonconformity categories (Blemish, Crack, Improper pull tab location, Missing pull tab, Others)
 - I = number of production lines = 3
 - J = types of nonconformity = 5
 - That is we test whether $p_{1j} = p_{2j} = p_{3j}$ for $j = 1, 2, \dots, 5$
 - H_a : Production lines are not homogeneous

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 9



Department of Computer Engineering
Kasetsart University

9

Example (cont.)

- Find \hat{e}_{ij} = expected number of samples = $n_i \frac{n_j}{n}$

		\hat{e}_{ij}					
		Blemish	Crack	Location	Missing	Others	Sample size
Production line	1	$\frac{150(89)}{375} = 35.60$	$\frac{150(145)}{375} = 58.00$	$\frac{150(58)}{375} = 23.20$	$\frac{150(54)}{375} = 21.60$	$\frac{150(29)}{375} = 11.60$	150
	2	$\frac{125(89)}{375} = 29.67$	48.33	19.33	18.00	9.67	125
	3	$\frac{100(89)}{375} = 23.73$	38.7	15.47	14.40	7.73	100
Total		89	145	58	54	29	375

n_i

n_j

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 10



Department of Computer Engineering
Kasetsart University

10

Example (cont.)

- Find test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$

		$\frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$				
		Blemish	Crack	Location	Missing	Others
Production line	1	$\frac{(34-35.60)^2}{35.60}$ = 0.072	$\frac{(65-58.00)^2}{58.00}$ = 0.845	$\frac{(17-23.20)^2}{23.20}$ = 1.657	$\frac{(21-21.60)^2}{21.60}$ = 0.017	$\frac{(13-11.60)^2}{11.60}$ = 0.169
	2	$\frac{(23-29.67)^2}{29.67}$ = 1.498	0.278	1.661	0.056	1.391
	3	$\frac{(32-23.73)^2}{23.73}$ = 2.879	2.943	0.018	0.011	0.664

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 14.159$

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 11



Department of Computer Engineering
Kasetsart University

11

Example (cont.)

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 14.159$
- Find rejection region:
 - Degree of freedom = $(I-1)(J-1) = (3-1)(5-1) = (2)(4) = 8$
 - $\chi^2_{0.05,8} = 15.507$
- Thus, we do not reject hypothesis at $\alpha = 0.05$
- At significance level = 0.05, all production lines are homogeneous in term of nonconformity categories

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 12



Department of Computer Engineering
Kasetsart University

12

Example (cont.)

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 14.159$
- Find p-value
 - Degree of freedom = $(I-1)(J-1) = (3-1)(5-1) = (2)(4) = 8$
 - P-Value = 0.077
- Thus, we do not reject hypothesis since p-value > $\alpha = 0.05$
- At significance level = 0.05, all production lines are homogeneous in term of nonconformity categories

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 13




Department of Computer Engineering
Kasetsart University

13

Example 2

- Compare two books whether they were written by the same author or not



- How to compare these two books?

Image Source: <http://www.clipartpanda.com/categories/school-book-clipart>

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 14



Department of Computer Engineering
Kasetsart University

14

Example 2

- Compare whether the frequencies of words in three of Austen's works are the same

Word	Sense and Sensibility	Emma	Sandition
a	147	186	101
an	25	26	11
this	32	39	15
that	94	105	37
with	59	74	28
without	18	10	10

- Test homogeneity



Outline

- Analysis of Categorical Data
 - Introduction
 - Homogeneity test
 - Independence test



Introduction

- A study of data in categories
- Case: Single population with two factors; One factor with I categories, and the other factor with J categories
 - Example: One department store, 6 departments (male clothes, female clothes, children, cosmetics, shoes, grocery) vs. 5 payment methods (cash, check, store credit card, Visa, Mastercard)
- Independence Test
 - Two factors occur independently

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 17



Department of Computer Engineering
Kasetsart University

17

Introduction (cont.)

- In general, data are put in the table
- Let n_{ij} = number of samples in (i,j) category
- Table contains $\{n_{ij}\}$'s is called two-way contingency table

	1	2	...	j	...	J
1	n_{11}	n_{12}	...	n_{1j}	...	n_{1J}
2	n_{21}					
...	...					
i	n_{i1}			n_{ij}		
...	...					
I	n_{I1}					n_{IJ}

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 18



Department of Computer Engineering
Kasetsart University

18

Independence Test

- Single population with two factors; One factor with I categories, and the other factor with J categories

- Let
 - n_{ij} = number of samples in (i,j) category
 - n_j = number of samples in j category = $\sum_i n_{ij}$
 - n_i = number of samples in i category = $\sum_j n_{ij}$
 - n = number of all samples = $\sum_i \sum_j n_{ij}$
 - p_{ij} = proportions of samples in (i,j) category
- Hypothesis test
 - Null hypothesis (H_0): $p_{ij} = p_i p_j$
 - Proportion of samples in categories i and j are independent
 - Alternative hypothesis (H_a): H_0 is not true

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 19



Department of Computer Engineering
Kasetsart University

19

Independence Test (cont.)

If two factors are independent, $p_{ij} = p_i p_j$

- Let \hat{e}_{ij} = expected number of samples = $np_{ij} = np_i p_j = n \frac{n_i}{n} \frac{n_j}{n} = \frac{n_i n_j}{n}$
- Test statistic
 - $\chi^2 = \sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$
- Rejection region
 - $\chi^2 \geq \chi_{\alpha, (I-1)(J-1)}^2$

Derivation of \hat{e}_{ij} is different from Homogeneity test

Same \hat{e}_{ij} as Homogeneity Test

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 20



Department of Computer Engineering
Kasetsart University

20

Example

- Study of gasoline station condition and aggressiveness in gasoline pricing
- Two factors: gasoline station condition (modern, standard, sub-standard) vs. aggressiveness in pricing (aggressive, neutral, nonaggressive)
- Test whether two factors are independent of each other at significance level = 0.01

		Aggressiveness in pricing			Sample Size
		Aggressive	Neutral	Non Aggressive	
Condition	Substandard	24	15	17	56
	Standard	52	73	80	205
	Modern	58	86	36	180
	Total	134	174	133	441

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 21



Department of Computer Engineering
Kasetsart University

21

Example (cont.)

- Hypothesis
 - H_0 : Gasoline station condition and aggressiveness in pricing are independent
 - I = number of conditions = 3
 - J = levels of pricing aggressiveness = 3
 - We test on $p_{ij} = p_i p_j$
 - H_a : Gasoline station condition and aggressiveness in pricing are not independent

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 22



Department of Computer Engineering
Kasetsart University

22

Example (cont.)

- Find \hat{e}_{ij} = expected number of samples = $\frac{n_i n_j}{n}$

		\hat{e}_{ij}			Sample Size
		Aggressive	Neutral	Non Aggressive	
Condition	Substandard	$\frac{56(134)}{441}$ =17.02	$\frac{56(174)}{441}$ =22.10	$\frac{56(133)}{441}$ =16.89	56
	Standard	$\frac{205(134)}{441}$ =62.29	80.88	61.83	205
	Modern	$\frac{180(134)}{441}$ =54.69	71.02	54.29	180
Total		134	174	133	441

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 23



Department of Computer Engineering
Kasetsart University

23

Example (cont.)

- Find test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$

		$\frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}}$		
		Aggressive	Neutral	Non Aggressive
Condition	Substandard	$\frac{(24-17.02)^2}{17.02}$ = 2.867	$\frac{(15-22.10)^2}{22.10}$ = 2.278	$\frac{(17-16.89)^2}{16.89}$ = 0.001
	Standard	$\frac{(52-62.29)^2}{62.29}$ = 1.700	0.769	5.343
	Modern	$\frac{(58-54.69)^2}{54.69}$ = 0.200	3.160	6.160

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 22.476$

Supaporn Erjongmanee
fengspe@ku.ac.th

Analysis of Categorical Data
Slide 24



Department of Computer Engineering
Kasetsart University

24

Example (cont.)

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 22.476$
- Given $\alpha = 0.01$, find p-value
 - Degree of freedom = $(I-1)(J-1) = (3-1)(3-1) = 4$
 - P-value 0.00016
- $P\text{-value} < \alpha = 0.01 \Rightarrow$ Null hypothesis is rejected
- Gasoline station condition and aggressiveness in pricing are dependent

```
from scipy.stats import chi2  
  
1-chi2.cdf(22.476,4)  
  
0.0001611050155756466
```



Example (cont.)

- Test statistic = $\sum_i \sum_j \frac{(n_{ij} - \hat{e}_{ij})^2}{\hat{e}_{ij}} = 22.476$
- Given $\alpha = 0.01$, find rejection region
 - Degree of freedom = $(I-1)(J-1) = (3-1)(3-1) = 4$
 - Thus, $\chi^2_{0.01,4} = 13.277$
- Null hypothesis is rejected
- Gasoline station condition and aggressiveness in pricing are dependent



Example 2

- Is there a relationship between marital status and educational level?

Education	Married once	Married more than once
College degree	550	61
No college degree	681	144

- Test independency



References

1. J.L. Devore and K.N.Berk, Modern Mathematical Statistics with Applications, Springer, 2012.
2. J.A. Rice, Mathematical Statistics and Data Analysis, Duxbury Press, 1995.

