

# Analysis of Variance Two Factors

**Dr. Supaporn Erjongmanee**

Department of Computer Engineering  
Kasetsart University  
fengspe@ku.ac.th

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 1



Department of Computer Engineering  
Kasetsart University

1

## Outline

- Two-Factor ANOVA

- Additive Factors

- Interaction Factors

- Interaction Plot

- Additions

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 2



Department of Computer Engineering  
Kasetsart University

2

## Two-Factor ANOVA : Additive Factors

- Compare two or more populations on two factors
  - Factor A has **I** level of treatments
  - Factor B has **J** level of treatments
  - Example: test washing detergents on pens
    - A = Brand of pens, **I** = 3
    - B = Washing detergent, **J** = 4

		Washing detergent			
		1	2	3	4
Brand of pens	1	0.97	0.48	0.48	0.46
	2	0.77	0.14	0.22	0.25
	3	0.67	0.39	0.57	0.19

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 3



Department of Computer Engineering  
Kasetsart University

3

## Additive Two-Factor ANOVA : Effects Model

$$X_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

- $X_{ij}$  = random sample j of treatment i
- $\mu$  = overall mean of treatment
- $\alpha_i$  = effect due to factor A at level i
- $\beta_j$  = effect due to factor B at level j
- $\varepsilon_{ij}$  = random error from sample j of treatment i
  - Assumed to be independent and normally distributed with mean = 0, variance =  $\sigma^2$
- Factor A is independent of factor B

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 4



Department of Computer Engineering  
Kasetsart University

4

## Additive Two-Factor ANOVA : Effects Model (cont.)

$$E(X_{ij}) = \mu + \alpha_i + \beta_j$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

- If  $\alpha_i = 0$  and  $\beta_j = 0$ , then all treatments have the same response

$$E(X_{ij}) = \mu$$

- Thus, null hypotheses for two additive factor ANOVA
  - Hypothesis on A: factor A at any level i has no effect on overall mean.
    - $H_{0A}: \alpha_1 = \alpha_2 = \dots = \alpha_I = 0$
  - Hypothesis on B: factor B at any level j has no effect on overall mean.
    - $H_{0B}: \beta_1 = \beta_2 = \dots = \beta_J = 0$



## Two-Factor ANOVA : Additive Factors (cont.)

- Let
  - $\mu_{ij}$  = mean of treatment i of factor A and treatment j of factor B
  - **I** = number of treatments from factor A
  - **J** = number of treatments from factor B
- Hypothesis on A: factor A at any level i has no effect on true mean.
  - $H_{0A}: \alpha_1 = \alpha_2 = \dots = \alpha_I = 0$
  - $H_{aA}$ : Not all  $\alpha_i$ 's are equal (Factor A has effect.)
- Hypothesis on B: factor B at any level j has no effect on true mean.
  - $H_{0B}: \beta_1 = \beta_2 = \dots = \beta_J = 0$
  - $H_{aB}$ : Not all  $\beta_j$ 's are equal (Factor B has effect.)



## Two-Factor ANOVA : Additive Factors (cont.)

- Test statistic (cont.):

- **I** = Number of treatments from factor A

- **J** = Number of treatments from factor B

$$SST = \sum_{i=1}^I \sum_{j=1}^J (X_{ij} - \bar{X})^2 \quad df = IJ - 1$$

$$SSA = \sum_{i=1}^I \sum_{j=1}^J (\bar{X}_i - \bar{X})^2 \quad df = I - 1$$

$$SSB = \sum_{i=1}^I \sum_{j=1}^J (\bar{X}_j - \bar{X})^2 \quad df = J - 1$$

$$SSE = \sum_{i=1}^I \sum_{j=1}^J (X_{ij} - \bar{X}_i - \bar{X}_j + \bar{X})^2 \quad df = (I - 1)(J - 1)$$

$$SST = SSA + SSB + SSE$$



## Two-Factor ANOVA : Additive Factors (cont.)

- Test statistic (cont.):

- **I** = Number of treatments from factor A

- **J** = Number of treatments from factor B

Another option:  
Sample-based  
computation

$$SST = \sum_{i=1}^I \sum_{j=1}^J X_{ij}^2 - \frac{1}{IJ} (\sum_{i=1}^I \sum_{j=1}^J X_{ij})^2 \quad df = IJ - 1$$

$$SSA = \frac{1}{J} \sum_{i=1}^I (\sum_{j=1}^J X_{ij})^2 - \frac{1}{IJ} (\sum_{i=1}^I \sum_{j=1}^J X_{ij})^2 \quad df = I - 1$$

$$SSB = \frac{1}{I} \sum_{j=1}^J (\sum_{i=1}^I X_{ij})^2 - \frac{1}{IJ} (\sum_{i=1}^I \sum_{j=1}^J X_{ij})^2 \quad df = J - 1$$

$$SSE = SST - SSA - SSB \quad df = (I - 1)(J - 1)$$



## Two-Factor ANOVA : Additive Factors (cont.)

- Test statistic (cont.):

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$SSA/(I-1)$	$f_A = MSA / MSE$	$f_A > F_{\alpha, I-1, (I-1)(J-1)}$
$H_{0B}$ vs. $H_{aB}$	$SSB/(J-1)$	$f_B = MSB / MSE$	$f_B > F_{\alpha, J-1, (I-1)(J-1)}$

## Example

- Test 4 washing detergents on 3 brands of pens at significance level = 0.05

		Washing detergent				
		1	2	3	4	
Brand of pens	1	0.97	0.48	0.48	0.46	0.598
	2	0.77	0.14	0.22	0.25	0.345
	3	0.67	0.39	0.57	0.19	0.455
		0.803	0.337	0.423	0.300	0.466

- A = Brand of pens, I = 3
- B = Washing detergents, J = 4
- $SST^* = 0.6947$ ,  $df = 11$
- $SSA^* = 0.1282$ ,  $df = 2$
- $SSB^* = 0.4797$ ,  $df = 3$
- $SSE = 0.6947 - 0.1282 - 0.4797 = 0.0868$ ,  $df = 11 - 2 - 3 = 6$

## Example (cont.)

- Test 4 washing detergents on 3 brands of pens (cont.):
  - $SST^* = 0.6947$ ,  $df = 11$
  - $SSA^* = 0.1282$ ,  $df = 2$
  - $SSB^* = 0.4797$ ,  $df = 3$
  - $SSE = 0.0868$ ,  $df = 6$

Hypothesis	Mean Square (MS = SS/df)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$0.1282/2$ $= 0.0641$	$f_A = 4.43$	$F_{0.05, 2, 6} = 5.14$
$H_{0B}$ vs. $H_{aB}$	$0.4797 / 3$ $= 0.1599$	$f_B = 11.05$	$F_{0.05, 3, 6} = 4.76$
Error	$0.0868/6$ $= 0.0144$		

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 11



Department of Computer Engineering  
Kasetsart University

11

## Example (cont.)

- Test 4 washing detergents on 3 brands of pens (cont.):

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$0.1282/2$ $= 0.0641$	$f_A = 4.43$	$F_{0.05, 2, 6} = 5.14$
$H_{0B}$ vs. $H_{aB}$	$0.4797 / 3$ $= 0.1599$	$f_B = 11.05$	$F_{0.05, 3, 6} = 4.76$
Error	$0.0868/6$ $= 0.0144$		

- $H_{0A}$  is not rejected. Factor A (brand of pens) has no effect on mark removal
- $H_{0B}$  is rejected. Factor B (washing detergents) has effect on mark removal

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 12



Department of Computer Engineering  
Kasetsart University


12

## Example

- Factor A (brand of pens) has no effect on mark removal
- Factor B (washing detergents) has effect on mark removal


		Washing detergent				
		1	2	3	4	
Brand of pens	1	0.97	0.48	0.48	0.46	0.598
	2	0.77	0.14	0.22	0.25	0.345
	3	0.67	0.39	0.57	0.19	0.455
		0.803	0.337	0.423	0.300	0.466

Brand of pens



Supaporn Erj  
fengspe@ku.ac.th

Washing detergent



Department of Computer Engineering  
Kasetsart University

13

## Multiple Comparisons

- To specify which mean is different from others
  - Fix on factor A or B
- Find Tukey's Honestly Significant Difference (HSD) using the following formula

$$\text{Factor A: } w_A = q_{\alpha, I, (I-1)(J-1)} \sqrt{\frac{MSE}{J}}$$

$$\text{Factor B: } w_B = q_{\alpha, J, (I-1)(J-1)} \sqrt{\frac{MSE}{I}}$$

- $q_{\alpha, m, n}$  = q-value from studentized range distribution with 2 degrees of freedom  $m, n$
- Follow the same steps as finding other HSD

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 14



Department of Computer Engineering  
Kasetsart University

14

## Example

$$\text{Factor A: } w_A = q_{\alpha, I, (I-1)(J-1)} \sqrt{\frac{MSE}{J}}$$

- Test 4 washing detergents on 3 brands of pens at significance level = 0.05


		Washing detergents				
		1	2	3	4	
Brand of pens	1	0.97	0.48	0.48	0.46	0.598
	2	0.77	0.14	0.22	0.25	0.345
	3	0.67	0.39	0.57	0.19	0.455
		0.803	0.337	0.423	0.300	0.466

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A} \text{ vs. } H_{aA}$	$\frac{0.1282}{2} = 0.0641$	$f_A = 4.43$	$F_{0.05, 2, 6} = 5.14$
$H_{0B} \text{ vs. } H_{aB}$	$\frac{0.4797}{3} = 0.1599$	$f_B = 11.05$	$F_{0.05, 3, 6} = 4.76$
Error	$\frac{0.0868}{6} = 0.0144$		

- Fixed at brands of pens:

$$w_A = q_{0.05, 3, 6} \sqrt{\frac{MSE}{J}} = 4.34 \sqrt{\frac{0.0144}{4}} = 0.261$$

- Sort factor-A sample means:  $\bar{x}_2, \bar{x}_3, \bar{x}_1$  0.345, 0.455, 0.598
- 1 group of means:  $\{\bar{x}_1, \bar{x}_2, \bar{x}_3\}$



Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 15

15

## Example (cont.)

$$\text{Factor B: } w_B = q_{\alpha, J, (I-1)(J-1)} \sqrt{\frac{MSE}{I}}$$

- Test 4 washing detergents on 3 brands of pens at significance level = 0.05


		Washing detergents				
		1	2	3	4	
Brand of pens	1	0.97	0.48	0.48	0.46	0.598
	2	0.77	0.14	0.22	0.25	0.345
	3	0.67	0.39	0.57	0.19	0.455
		0.803	0.337	0.423	0.300	0.466

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A} \text{ vs. } H_{aA}$	$\frac{0.1282}{2} = 0.0641$	$f_A = 4.43$	$F_{0.05, 2, 6} = 5.14$
$H_{0B} \text{ vs. } H_{aB}$	$\frac{0.4797}{3} = 0.1599$	$f_B = 11.05$	$F_{0.05, 3, 6} = 4.76$
Error	$\frac{0.0868}{6} = 0.0144$		

- Fixed at detergent factor:

$$w_B = q_{0.05, 4, 6} \sqrt{\frac{MSE}{I}} = 4.90 \sqrt{\frac{0.0144}{3}} = 0.340$$

- Sort factor-B sample means:  $\bar{x}_4, \bar{x}_2, \bar{x}_3, \bar{x}_1$  0.300, 0.337, 0.423, 0.803
- 2 groups of means:  $\{\bar{x}_4, \bar{x}_2, \bar{x}_3\}$  and  $\bar{x}_1$



Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 16



Dep  
Kas

16



## Example 2

- Test 4 coatings on 3 soil type for corrosion at significance level = 0.05

		Soil Type (B)			$\bar{x}_j$
		1	2	3	
Coating (A)	1	64	49	50	54.33
	2	53	51	48	50.67
	3	47	45	50	47.33
	4	51	43	52	48.67
	$\bar{x}_i$	53.75	47.00	50.00	50.25

- A = Coatings, I = 4, B = Soil Types, J = 3
- SST = 242.063, df = 11  $SST = \sum_{i=1}^I \sum_{j=1}^J (X_{ij} - \bar{X})^2$
- SSA = 83.583, df = 3  $SSA = \sum_{i=1}^I \sum_{j=1}^J (\bar{X}_i - \bar{X})^2$
- SSB = 91.500, df = 2  $SSB = \sum_{i=1}^I \sum_{j=1}^J (\bar{X}_j - \bar{X})^2$
- SSE = SST - SSA - SSB = 66.979, df = 11 - 3 - 2 = 6

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 17



Department of Computer Engineering  
Kasetsart University

17

## Example 2 (cont.)

- Test 4 coatings on 3 soil types for corrosion (cont.):

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$83.583/3 = 27.861$	$f_A = 2.495$	$F_{0.05, 3, 6} = 4.7571$
$H_{0B}$ vs. $H_{aB}$	$91.500 / 2 = 45.750$	$f_B = 4.098$	$F_{0.05, 2, 6} = 5.1433$
Error	$66.97/6 = 11.163$		

- $H_{0A}$  is not rejected. Factor A (coatings) has no effect on corrosion
- $H_{0B}$  is not rejected. Factor B (soil types) has no effect on corrosion

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 18



Department of Computer Engineering  
Kasetsart University

18

# Outline

- Two-Factor ANOVA
  - Additive Factors
  - Interaction Factors
  - Interaction Plot
- Additions



## Two-Factor ANOVA : Interaction Factors

- Compare two or more populations on two interaction factors
  - Factor A has **I** level of treatments
  - Factor B has **J** level of treatments
  - **K** Samples from treatment of factors and B are collected
  - Example: 3 varieties of tomatoes on 4 planting density

	Planting Density											
Variety	10,000			20,000			30,000			40,000		
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5
I	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2



## Interaction Two-Factor ANOVA : Effects Model

$$X_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

- $X_{ij}$  = random sample  $j$  of treatment  $i$
- $\mu$  = overall mean of treatment  $i$  on factor  $j$
- $\alpha_i$  = effect due to factor A at level  $i$
- $\beta_j$  = effect due to factor B at level  $j$
- $\gamma_{ij}$  = interaction parameter between factors A and B
- $\varepsilon_{ij}$  = random error from sample  $j$  of treatment  $i$ 
  - Assumed to be independent and normally distributed with mean = 0, variance =  $\sigma^2$
- Factor A is not independent of factor B
  - If  $\gamma_{ij}$  's are all zeros, factors A and B are independent.

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 21



Department of Computer Engineering  
Kasetsart University

21

## Interaction Two-Factor ANOVA : Effects Model (cont.)


$$X_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

- If  $\alpha_i = 0$ ,  $\beta_j = 0$ , and  $\gamma_{ij} = 0$ , then all treatments have the same response

$$E(X_{ijk}) = \mu$$

- Thus, null hypotheses for interaction factor ANOVA
  - Hypothesis on A and B: factors A and B at any level  $i$  has no effect on overall mean.
    - $H_{0AB}: \gamma_{ij} = 0$  for all  $i, j$   Test first If reject, no need to test  $H_{0A}$ ,  $H_{0B}$
  - Hypothesis on A: factor A at any level  $i$  has no effect on overall mean.
    - $H_{0A}: \alpha_1 = \alpha_2 = \dots = \alpha_I = 0$
  - Hypothesis on B: factor B at any level  $j$  has no effect on overall mean.
    - $H_{0B}: \beta_1 = \beta_2 = \dots = \beta_J = 0$

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 22



Department of Computer Engineering  
Kasetsart University

22

## Two-Factor ANOVA : Interaction Factors (cont.)

- Let
  - $\mu_{ijk}$  = mean of treatment i of factor A and treatment j of factor B
  - **I** = number of treatments from factor A
  - **J** = number of treatments from factor B
  - **K** = number of samples per treatments from factors A and B
- Hypothesis :
  - $H_{0AB}$ :  $\gamma_{ij} = 0$  for all i,j
  - $H_{aAB}$ : At least  $\gamma_{ij} \neq 0$
  - $H_{0A}$ :  $\alpha_1 = \alpha_2 = \dots = \alpha_I = 0$
  - $H_{aA}$ : At least  $\alpha_i \neq 0$
  - $H_{0B}$ :  $\beta_1 = \beta_2 = \dots = \beta_J = 0$
  - $H_{aB}$ : At least  $\beta_j \neq 0$



## Two-Factor ANOVA : Interaction Factors

- Test statistic (cont.):

$$SST = \sum_i \sum_j \sum_k (X_{ijk} - \bar{X})^2 \quad df = IJK - 1$$

$$SSA = \sum_i \sum_j \sum_k (\bar{X}_i - \bar{X})^2 \quad df = I - 1$$

$$SSB = \sum_i \sum_j \sum_k (\bar{X}_j - \bar{X})^2 \quad df = J - 1$$

$$SSAB = \sum_i \sum_j \sum_k (\bar{X}_{ij} - \bar{X}_i - \bar{X}_j + \bar{X})^2 \quad df = (I - 1)(J - 1)$$

$$SSE = \sum_i \sum_j \sum_k (X_{ijk} - \bar{X}_{ij})^2 \quad df = IJ(K - 1)$$

$$SST = SSA + SSB + SSAB + SSE$$



## Two-Factor ANOVA : Additive Factors (cont.)

### • Test statistic (cont.):

Another option:  
Sample-based computation

$$SST = \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K X_{ijk}^2 - \frac{1}{IJK} \left( \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K X_{ijk} \right)^2 \quad df = IJK - 1$$

$$SSA = \frac{1}{JK} \sum_{i=1}^I \left( \sum_{j=1}^J \sum_{k=1}^K X_{ijk} \right)^2 - \frac{1}{IJK} \left( \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K X_{ijk} \right)^2 \quad df = I - 1$$

$$SSB = \frac{1}{IK} \sum_{j=1}^J \left( \sum_{i=1}^I \sum_{k=1}^K X_{ijk} \right)^2 - \frac{1}{IJK} \left( \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K X_{ijk} \right)^2 \quad df = J - 1$$

$$SSE = \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K X_{ijk}^2 - \frac{1}{K} \sum_{i=1}^I \sum_{j=1}^J \left( \sum_{k=1}^K X_{ijk} \right)^2 \quad df = (I - 1)(J - 1)$$

$$SSAB = SST - SSA - SSB - SSE$$

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 25



Department of Computer Engineering  
Kasetsart University

25

## Two-Factor ANOVA : Interaction Factors (cont.)

### • Test statistic (cont.):

- I = Number of treatments from factor A
- J = Number of treatments from factor B
- K = number of samples per treatments from factors A and B

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$SSA/(I-1)$	$f_A = MSA / MSE$	$f_A > F_{\alpha, I-1, IJ(K-1)}$
$H_{0B}$ vs. $H_{aB}$	$SSB/(J - 1)$	$f_B = MSB / MSE$	$f_B > F_{\alpha, J-1, IJ(K-1)}$
$H_{0AB}$ vs. $H_{aAB}$	$\frac{SSAB}{(I - 1)(J - 1)}$	$f_{AB} = MSAB / MSE$	$f_{AB} > F_{\alpha, (I-1)(J-1), IJ(K-1)}$

Supaporn  
fengspe@ku.ac.th

Department of Computer Engineering  
Kasetsart University

26

## Example

- Test 3 varieties of tomatoes on 4 planting density at significance level = 0.01

	Planting Density											
Variety	10,000			20,000			30,000			40,000		
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5
Ife	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2

- A = Varieties of tomatoes, I = 3
- B = Planting densities, J = 4
- K = Number of samples per factors A and B = 3
- IJK = 36

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 27



Department of Computer Engineering  
Kasetsart University

27

## Example (cont.)

- Test 3 varieties of tomatoes on 4 planting density at significance level = 0.01

	Planting Density											
Variety	10,000			20,000			30,000			40,000		
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5
Ife	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2

- SST = 460.36, df = 35 (IJK - 1)
- SSA = 327.60, df = 2 (I-1)
- SSB = 86.69, df = 3 (J-1)
- SSE = 38.04, df = 24 (IJ(K-1)) -> MSE = 38.04 / 24 = 1.59
- SSAB = 460.36 - 327.60 - 86.69 - 38.04 = 8.03, df = 35 - 2 - 3 - 24 = 6

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 28



Department of Computer Engineering  
Kasetsart University

28

## Example (cont.)

- Test statistic :

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	163.8	$f_A = 103.02$	$F_{0.01, 2, 24} = 5.61$
$H_{0B}$ vs. $H_{aB}$	28.9	$f_B = 18.18$	$F_{0.01, 3, 24} = 4.72$
$H_{0AB}$ vs. $H_{aAB}$	1.34	$f_{AB} = 0.84$	$F_{0.01, 6, 24} = 3.67$

- $H_{0AB}$  is not rejected. Interaction has no effect.
- $H_{0A}$  is rejected. Factor A (varieties of tomatoes) has effect on average product
- $H_{0B}$  is rejected. Factor B (planting densities) has effect on average product

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 29



Department of Computer Engineering  
Kasetsart University

29

## Multiple Comparisons

- When interaction is rejected and one or both factors has the effect, we can perform multiple comparisons.
- To specify which mean is different from others
  - Fix on factor A or B
  - Find Tukey's Honestly Significant Difference (HSD) using the following formula
 
$$\text{Factor A: } w_A = q_{\alpha, I, J(K-1)} \sqrt{\frac{MSE}{JK}}$$

$$\text{Factor B: } w_B = q_{\alpha, J, IJ(K-1)} \sqrt{\frac{MSE}{IK}}$$
  - $q_{\alpha, m, n}$  = q-value from studentized range distribution with 2 degrees of freedom  $m, n$
- Follow the same steps as finding other HSD

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 30



Department of Computer Engineering  
Kasetsart University

30

## Example (cont.)

$$I = 3, J = 4, K = 3$$

$$IJ(K-1) = 24$$

- Test 3 varieties of tomatoes on 4 planting density at  $\alpha = 0.01$

	Planting Density												
Variety	10,000			20,000			30,000			40,000			$\bar{x}_i$
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5	11.33
Ife	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5	12.21
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2	18.13

**Factor A:**  $w_A = q_{\alpha, I, IJ(K-1)} \sqrt{\frac{MSE}{JK}}$   $w_A = q_{0.01, 3, 24} \sqrt{\frac{MSE}{JK}} = 4.55 \sqrt{\frac{1.59}{12}} = 1.66$

- Sort sample means ( $\bar{x}_i$ ):  $\bar{x}_1$ ,  $\bar{x}_2$ ,  $\bar{x}_3$  : 11.33, 12.21, 18.13
- 2 groups of means:  $\{\bar{x}_1, \bar{x}_2\}$  and  $\bar{x}_3$

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 31



Department of Computer Engineering  
Kasetsart University

31

## Example (cont.)

$$I = 3, J = 4, K = 3$$

$$IJ(K-1) = 24$$

- Test 3 varieties of tomatoes on 4 planting density at  $\alpha = 0.01$

	Planting Density											
Variety	10,000			20,000			30,000			40,000		
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5
Ife	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2
$\bar{x}_j$	11.48			14.39			15.78			13.91		

**Factor B:**  $w_B = q_{\alpha, J, IJ(K-1)} \sqrt{\frac{MSE}{IK}}$   $w_B = q_{0.01, 4, 24} \sqrt{\frac{MSE}{IK}} = 4.91 \sqrt{\frac{1.59}{9}} = 2.06$

- Sort sample means ( $\bar{x}_i$ ):  $\bar{x}_1$ ,  $\bar{x}_4$ ,  $\bar{x}_2$ ,  $\bar{x}_3$  : 11.48, 13.91, 14.39, 15.78
- 2 groups of mean:  $\bar{x}_1$  and  $\{\bar{x}_4, \bar{x}_2, \bar{x}_3\}$

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 32



Department of Computer Engineering  
Kasetsart University

32



## Example 2

- Test 2 varieties of Iron (Fe) on 3 concentration doses
- 18 samples per category

Fe <sup>2+</sup>			Fe <sup>3+</sup>		
10.2	1.2	0.3	10.2	1.2	0.3
0.71	2.20	2.25	2.20	4.04	2.71
1.66	2.93	3.93	2.69	4.16	5.43
2.01	3.08	5.08	3.54	4.42	6.38
2.16	3.49	5.82	3.75	4.93	6.38
2.42	4.11	5.84	3.83	5.49	8.32
2.42	4.95	6.89	4.08	5.77	9.04
2.56	5.16	8.50	4.27	5.86	9.56
2.60	5.54	8.56	4.53	6.28	10.01
3.31	5.68	9.44	5.32	6.97	10.08
3.64	6.25	10.52	6.18	7.06	10.62
3.74	7.25	13.46	6.22	7.78	13.80
3.74	7.90	13.57	6.33	9.23	15.99
4.39	8.85	14.76	6.97	9.34	17.90
4.50	11.96	16.41	6.97	9.91	18.25
5.07	15.54	16.96	7.52	13.46	19.32
5.26	15.89	17.56	8.36	18.40	19.87
8.15	18.30	22.82	11.65	23.89	21.60
8.24	18.59	29.13	12.45	26.39	22.25

- A = Forms of Irons, I = 2, B = Concentration of doses, J = 3
- K = Number of samples per factors A and B = 18, IJK = 108

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 33



Department of Computer Engineering  
Kasetsart University

33

## Example 2 (cont.)

- Test 2 varieties of Iron (Fe) on 3 concentration doses

$$\bar{x} = 8.64$$

$$\bar{x}_{Fe^{2+}} = 7.88$$

$$\bar{x}_{Fe^{3+}} = 9.40$$

$$\bar{x}_{10.2} = 4.82$$

$$\bar{x}_{1.2} = 8.92$$

$$\bar{x}_{0.3} = 12.19$$

- SST = 3992.37, df = 107 (IJK - 1)
- SSA = 62.26, df = 1 (I - 1)
- SSB = 983.62, df = 2 (J - 1)
- SSE = 2938.20, df = 102 (IJ(K - 1))
- SSAB = SST - SSA - SSB - SSE = 8.29, df = 2

Fe <sup>2+</sup>			Fe <sup>3+</sup>		
10.2	1.2	0.3	10.2	1.2	0.3
0.71	2.20	2.25	2.20	4.04	2.71
1.66	2.93	3.93	2.69	4.16	5.43
2.01	3.08	5.08	3.54	4.42	6.38
2.16	3.49	5.82	3.75	4.93	6.38
2.42	4.11	5.84	3.83	5.49	8.32
2.42	4.95	6.89	4.08	5.77	9.04
2.56	5.16	8.50	4.27	5.86	9.56
2.60	5.54	8.56	4.53	6.28	10.01
3.31	5.68	9.44	5.32	6.97	10.08
3.64	6.25	10.52	6.18	7.06	10.62
3.74	7.25	13.46	6.22	7.78	13.80
3.74	7.90	13.57	6.33	9.23	15.99
4.39	8.85	14.76	6.97	9.34	17.90
4.50	11.96	16.41	6.97	9.91	18.25
5.07	15.54	16.96	7.52	13.46	19.32
5.26	15.89	17.56	8.36	18.40	19.87
8.15	18.30	22.82	11.65	23.89	21.60
8.24	18.59	29.13	12.45	26.39	22.25

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 34



Department of Computer Engineering  
Kasetsart University

34

## Example 2(cont.)

### • Test statistic :

- SST = 3992.37, df = 107
- SSA = 62.26, df = 1
- SSB = 983.62, df = 2
- SSE = 2938.20, df = 102
- SSAB = 8.29, df = 2

$$MSE = 2938.20 / 102 = 28.81$$

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection region
$H_{0A}$ vs. $H_{aA}$	$62.26/1 = 62.26$	$f_A = 62.26/28.81 = 2.16$	$F_{0.01, 1, 102} = 6.89$ $F_{0.05, 1, 102} = 3.93$ $F_{0.1, 1, 102} = 2.76$
$H_{0B}$ vs. $H_{aB}$	$983.62/2 = 491.81$	$f_B = 491.81/28.81 = 17.07$	$F_{0.01, 2, 102} = 4.82$ $F_{0.05, 2, 102} = 3.09$ $F_{0.1, 2, 102} = 2.36$
$H_{0AB}$ vs. $H_{aAB}$	$8.29/2 = 4.15$	$f_{AB} = 4.15/28.81 = 0.14$	$F_{0.01, 2, 102} = 4.82$ $F_{0.05, 2, 102} = 3.09$ $F_{0.1, 2, 102} = 2.36$

- $H_{0AB}$  is not rejected. Interaction has no effect.
- $H_{0A}$  is not rejected. Factor A (forms of Iron) has no effect on % of retained iron
- $H_{0B}$  is rejected. Factor B (Concentration) has effect on % of retained iron

Supaporn  
fengspei

35

## Example 2(cont.)

### • Test statistic :

- SST = 3992.37, df = 107
- SSA = 62.26, df = 1
- SSB = 983.62, df = 2
- SSE = 2938.20, df = 102
- SSAB = 8.29, df = 2

$$MSE = 2938.20 / 102 = 28.81$$

Hypothesis	Mean Square (MS)	Test statistic (f)	P-value
$H_{0A}$ vs. $H_{aA}$	$62.26/1 = 62.26$	$f_A = 62.26/28.81 = 2.16$	0.14 Accept at 0.1
$H_{0B}$ vs. $H_{aB}$	$983.62/2 = 491.81$	$f_B = 491.81/28.81 = 17.07$	$4.02 \times 10^{-7}$ Reject at 0.01
$H_{0AB}$ vs. $H_{aAB}$	$8.29/2 = 4.15$	$f_{AB} = 4.15/28.81 = 0.14$	0.87 Accept at 0.1

- $H_{0AB}$  is not rejected. Interaction has no effect.
- $H_{0A}$  is not rejected. Factor A (forms of Iron) has no effect on % of retained iron
- $H_{0B}$  is rejected. Factor B (Concentration) has effect on % of retained iron

Si  
fe

Adelphi University

36

## Outline

- Two-Factor ANOVA
  - Additive Factors
  - Interaction Factors
  - Interaction Plot
- Addition



## Additive Two-Factor ANOVA : Effects Model

$$X_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

Adding effect of factors A and B

- $X_{ij}$  = random sample j of treatment i
- $\mu$  = overall mean of treatment
- $\alpha_i$  = effect due to factor A at level i
- $\beta_j$  = effect due to factor B at level j
- $\varepsilon_{ij}$  = random error from sample j of treatment i
  - Assumed to be independent and normally distributed with mean = 0, variance =  $\sigma^2$
- Factor A is independent of factor B



## Interaction Plot : Example 1

$$X_{ij} = \mu + \alpha_i + \beta_j + \varepsilon_{ij}$$

$$\sum \alpha_i = 0$$


$$\sum \beta_j = 0$$

Adding effect of factors A and B

2 factors affect job applicants

A. Weight

B. Relationship type



Both lines (means) are (almost) parallel

Two lines are not crossing -> Change in one factor does not affect the other factor

When value of one factor changes, mean changes

Image source: [http://onlinestatbook.com/2/analysis\\_of\\_variance/multiway.html](http://onlinestatbook.com/2/analysis_of_variance/multiway.html)

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 39



Department of Computer Engineering  
Kasetsart University

39

## Interaction Plot : Example 3

2 factors affect on students' scores

A. Typing experience: none, some, lots (x-axis)

B. Exam method: book, computer (lines)

Check whether  $\left\{ \begin{array}{l} \text{each factor affects on students' scores or not} \\ \text{both factors interact with each other or not} \end{array} \right.$

2 factors: A and B



8 possibilities of factor effects

How many possibilities of factor affects?

Source: <http://www.psychstat.missouristate.edu/multibook/mlt09.htm>

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 40




Department of Computer Engineering  
Kasetsart University

40

## Interaction Plot : Example 3 (cont.)

2 factors affect on students' scores


- A. Typing experience: none, some, lots (x-axis)
- B. Exam method: book, computer (lines)




No effect from either A or B



Effect from A only



Effect from B only



Effect on A and B, but not AxB

Source: <http://www.psychstat.missouristate.edu/multibook/mlt09.htm>

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 41




Department of Computer Engineering  
Kasetsart University

41


## Interaction Plot : Example 3 (cont.)

2 factors affect on students' scores


- A. Typing experience: none, some, lots (x-axis)
- B. Exam method: book, computer (lines)




Effect on AxB only



Effect from A and AxB



Effect from B and AxB



Effect on A, B, and AxB

Source: <http://www.psychstat.missouristate.edu/multibook/mlt09.htm>

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 42



Department of Computer Engineering  
Kasetsart University

42

## Interaction Plot : Example 5

- Test 3 varieties of tomatoes on 4 planting density at significance level = 0.01

Variety	Planting Density											
	10,000			20,000			30,000			40,000		
H	10.5	9.2	7.9	12.8	11.2	13.3	12.1	12.6	14.0	10.8	9.1	12.5
Ife	8.1	8.6	10.1	12.7	13.7	11.5	14.4	15.4	13.7	11.3	12.5	14.5
P	16.1	15.3	17.5	16.6	19.2	18.5	20.8	18.0	21.0	18.4	18.9	17.2

- A = Varieties of tomatoes, I = 3
- B = Planting densities, J = 4
- K = Number of samples per factors A and B = 3
- IJK = 36

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 43




Department of Computer Engineering  
Kasetsart University

43

## Interaction Plot : Example 5 (cont.)

- Test statistic :

Hypothesis	Mean Square (MS)	Test statistic (f)	Reject
$H_{0A}$ vs. $H_{aA}$	163.8	$f_A = 103.02$	$F_{0.01}$
$H_{0B}$ vs. $H_{aB}$	28.9	$f_B = 18.18$	$F_{0.01}$
$H_{0AB}$ vs. $H_{aAB}$	1.34	$f_{AB} = 0.84$	$F_{0.01}$



- $H_{0AB}$  is not rejected. Interaction has no effect.
- $H_{0A}$  is rejected. Factor A (varieties of tomatoes) has effect on average product
- $H_{0B}$  is rejected. Factor B (planting densities) has effect on average product

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 44



Department of Computer Engineering  
Kasetsart University

44

## Interaction Plot : Example 6

- Test 2 varieties of Iron (Fe) on 3 concentration doses
- 18 samples on % retainment per category

Fe <sup>2+</sup>			Fe <sup>3+</sup>		
10.2	1.2	0.3	10.2	1.2	0.3
0.71	2.20	2.25	2.20	4.04	2.71
1.66	2.93	3.93	2.69	4.16	5.43
2.01	3.08	5.08	3.54	4.42	6.38
2.16	3.49	5.82	3.75	4.93	6.38
2.42	4.11	5.84	3.83	5.49	8.32
2.42	4.95	6.89	4.08	5.77	9.04
2.56	5.16	8.50	4.27	5.86	9.56
2.60	5.54	8.56	4.53	6.28	10.01
3.31	5.68	9.44	5.32	6.97	10.08
3.64	6.25	10.52	6.18	7.06	10.62
3.74	7.25	13.46	6.22	7.78	13.80
3.74	7.90	13.57	6.33	9.23	15.99
4.39	8.85	14.76	6.97	9.34	17.90
4.50	11.96	16.41	6.97	9.91	18.25
5.07	15.54	16.96	7.52	13.46	19.32
5.26	15.89	17.56	8.36	18.40	19.87
8.15	18.30	22.82	11.65	23.89	21.60
8.24	18.59	29.13	12.45	26.39	22.25

- A = Forms of Irons, I = 2, B = Concentration of doses, J = 3
- K = Number of samples per factors A and B = 18, IJK = 108

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 45



Department of Computer Engineering  
Kasetsart University


45

## Interaction Plot : Example 6 (cont.)

Use interaction plot as GUIDELINE,  
not actual decision on rejection in ANOVA

- Test statistic :

Hypothesis	Mean Square (MS)	Test statistic (f)	Rejection r
H <sub>0A</sub> vs. H <sub>aA</sub>	62.26/1 = 62.26	f <sub>A</sub> = 62.26/28.81 = 2.16	F <sub>0.01, 1, 102</sub> = 6.90 F <sub>0.05, 1, 102</sub> = 3.84 F <sub>0.1, 1, 102</sub> = 2.77
H <sub>0B</sub> vs. H <sub>aB</sub>	983.62/2 = 491.81	f <sub>B</sub> = 491.81/28.81 = 17.07	F <sub>0.01, 2, 102</sub> = 4.61 F <sub>0.05, 2, 102</sub> = 3.10 F <sub>0.1, 2, 102</sub> = 2.35
H <sub>0AB</sub> vs. H <sub>aAB</sub>	8.29/2 = 4.15	f <sub>AB</sub> = 4.15/28.81 = 0.14	F <sub>0.01, 2, 102</sub> = 4.61 F <sub>0.05, 2, 102</sub> = 3.10 F <sub>0.1, 2, 102</sub> = 2.35



- H<sub>0AB</sub> is not rejected. Interaction has no effect.
- H<sub>0A</sub> is not rejected. Factor A (forms of Iron) has no effect on % of retained iron
- H<sub>0B</sub> is rejected. Factor B (Concentration) has effect on % of retained iron

46

## Outline

- Two-Factor ANOVA
  - Additive Factors
  - Interaction Factors
  - Interaction Plots
- Additions



## More on Multiple Comparison

- Is it possible that  $H_{0A}$  (or  $H_{0B}$ ) is not rejected but result to multiple group of means?
- Is it also possible that  $H_{0A}$  (or  $H_{0B}$ ) is rejected but have one group of means?

ANOVA tests on ALL means whether they are identical

Multiple comparison tests on PAIRWISE means

Measured by MSE

Not measured by MSE

ANOVA detects variability among all means

ANOVA test is more sensitive than Multiple comparison





## More on Multiple Comparison (cont.)

ANOVA tests on ALL means whether they are identical

Multiple comparison tests on PAIRWISE means

ANOVA detects lower variability among all means

ANOVA test is more sensitive than Multiple comparison




Image source: <http://blog.minitab.com/blog/adventures-in-statistics-2/understanding-analysis-of-variance-anova-and-the-f-test>

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 49



Department of Computer Engineering  
Kasetsart University

49

## Interaction Two-Factor ANOVA : Effects Mode

$$X_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk}$$

$$\sum \alpha_i = 0$$

$$\sum \beta_j = 0$$

- If  $\alpha_i = 0$ ,  $\beta_j = 0$ , and  $\gamma_{ij} = 0$ , then all treatments have the same response

$$E(X_{ijk}) = \mu$$

- Thus, null hypotheses for interaction factor ANOVA

- Hypothesis on A and B:** factors A and B at any level i has no effect on overall mean.

- $H_{0AB}: \gamma_{ij} = 0$  for all i,j



Test first

If reject, no need to test  $H_{0A}$ ,  $H_{0B}$

- Hypothesis on A:**

factor A at any level i has no effect on overall mean.

- $H_{0A}: \alpha_1 = \alpha_2 = \dots = \alpha_i = 0$

A interacts with B

Effect of A depends on B  
Thus, A and B both affect.

- Hypothesis on B:** factor B at any level j has no effect on overall mean.

- $H_{0B}: \beta_1 = \beta_2 = \dots = \beta_j = 0$

Supaporn Erjongmanee  
fengspe@ku.ac.th

Statistics in Computer Engineering  
Slide 50



Department of Computer Engineering  
Kasetsart University

50

## More on ANOVA

- When  $H_{0AB}$  is rejected and  $H_{0A}$  (or  $H_{0B}$ ) is not rejected, what does it mean?

A and B interact.

A and B both affect.

That's why all means are equal  
and  $H_{0A}$  is not rejected.

A has effect but its effect (that depends on B) is the same for all responses



## References

1. J.L. Devore and K.N. Berk, Modern Mathematical Statistics with Applications, Springer, 2012.
2. J.A. Rice, Mathematical Statistics and Data Analysis, Duxbury Press, 1995.

