

Multiple Linear Regression

Dr. Supaporn Erjongmanee

Department of Computer Engineering
Kasetsart University
fengspe@ku.ac.th

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 1



Department of Computer Engineering
Kasetsart University

1

Outline

- Introduction
- Estimating Parameters
- Residual, SSE, SST, R^2
- Example

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 2



Department of Computer Engineering
Kasetsart University

2

Multiple Regression Model

Multiple Regression



- General model

$$Y = b_0 + b_1x_1 + b_2x_2 + \dots + b_kx_k + \varepsilon$$

- where

- X = independent/predictor variables
- Y = dependent variable
- k = number of predictors
- ε is normally distributed with $\mu = 0$, $\text{var} = \sigma^2$
 - When $\sigma^2 \rightarrow 0$, ε is close to zero, Y is closer to true regression line
 - When σ^2 is large, ε is close to zero, Y is closer to true regression line
- b_j = rate how y increases according to x_j increases

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 3



Department of Computer Engineering
Kasetsart University

3

Observation Set

- Let x_{ij} = observed j^{th} predictor in i^{th} data set
 - $i = 1, 2, \dots, n$, $j = 1, 2, \dots, k$
- The data set is composed of n sets:

- $(x_{11}, x_{12}, x_{13}, \dots, x_{1k}, y_1)$ $y_1 = b_0 + b_1x_{11} + b_2x_{12} + \dots + b_kx_{1k} + \varepsilon$

- $(x_{21}, x_{22}, x_{23}, \dots, x_{2k}, y_2)$ $y_2 = b_0 + b_1x_{21} + b_2x_{22} + \dots + b_kx_{2k} + \varepsilon$

- ...

- $(x_{n1}, x_{n2}, x_{n3}, \dots, x_{nk}, y_n)$ $y_n = b_0 + b_1x_{n1} + b_2x_{n2} + \dots + b_kx_{nk} + \varepsilon$

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 4



Department of Computer Engineering
Kasetsart University

4

Outline

- Introduction
- Estimating Parameters
- Residual, SSE, SST, R^2
- Example



Estimating Parameters

$$Y = b_0 + b_1x_1 + b_2x_2 + \dots + b_kx_k + \varepsilon$$

- Parameters to estimate are
 - b_1, b_2, \dots, b_k
- Also, apply principle of least squares to minimize sum of errors $[g(\cdot)]$ to find estimated parameters

$$g(b_0, b_1, b_2, b_k) = \sum_{i=1}^n [y_i - (b_0 + b_1 x_{i1} + \dots + b_k x_{ik})]^2$$



Estimating Parameters (cont.)

$$g(b_0, b_1, b_2, b_k) = \sum_{i=1}^n [y_i - (b_0 + b_1 x_{i1} + \dots + b_k x_{ik})]^2$$

- Take partial derivatives of $g(\cdot)$ with respect to each b_j and set to zero.

$$\begin{aligned} b_0 \sum_{i=1}^n 1 + b_1 \sum_{i=1}^n x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} &= \sum_{i=1}^n y_i \\ b_0 \sum_{i=1}^n x_{i1} + b_1 \sum_{i=1}^n x_{i1} x_{i1} + \dots + b_k \sum_{i=1}^n x_{i1} x_{ik} &= \sum_{i=1}^n x_{i1} y_i \\ \dots \\ b_0 \sum_{i=1}^n x_{ik} + b_1 \sum_{i=1}^n x_{ik} x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} x_{ik} &= \sum_{i=1}^n x_{ik} y_i \end{aligned}$$

- Help for solving b_j 's
 - Use software for help finding values of b_j 's
 - Use matrix to help out calculation



Regression with Matrices

$$y = b_0 + b_1 x_1 + b_2 x_2 + \dots + b_k x_k + \varepsilon$$

$$\mathbf{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \quad \mathbf{X} = \begin{bmatrix} 1 & x_{11} & \dots & x_{1k} \\ & \vdots & & \\ 1 & x_{n1} & \dots & x_{nk} \end{bmatrix} \quad \boldsymbol{\beta} = \begin{bmatrix} b_0 \\ \vdots \\ b_k \end{bmatrix} \quad \boldsymbol{\varepsilon} = \begin{bmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

$$\begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} 1 & x_{11} & \dots & x_{1k} \\ & \vdots & & \\ 1 & x_{n1} & \dots & x_{nk} \end{bmatrix} \begin{bmatrix} b_0 \\ \vdots \\ b_k \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_n \end{bmatrix}$$



Regression with Matrices (cont.)

$$\begin{aligned}
 g(b_0, b_1, b_2, b_k) \\
 &= \sum_{i=1}^n [y_i - (b_0 + b_1 x_{i1} + \dots + b_k x_{ik})]^2 \\
 &= (\mathbf{y} - \mathbf{X}\boldsymbol{\beta})'(\mathbf{y} - \mathbf{X}\boldsymbol{\beta}) \\
 &= \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|^2
 \end{aligned}$$

- Apply principle of least squares to minimize sum of errors (\cdot) to find $\boldsymbol{\beta}$
- Take partial derivatives of $g(\cdot)$ with respect to each b_j and set to zero.

$$\begin{aligned}
 b_0 \sum_{i=1}^n 1 + b_1 \sum_{i=1}^n x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} &= \sum_{i=1}^n y_i \\
 b_0 \sum_{i=1}^n x_{i1} + b_1 \sum_{i=1}^n x_{i1} x_{i1} + \dots + b_k \sum_{i=1}^n x_{i1} x_{ik} &= \sum_{i=1}^n x_{i1} y_i \\
 \dots \\
 b_0 \sum_{i=1}^n x_{ik} + b_1 \sum_{i=1}^n x_{ik} x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} x_{ik} &= \sum_{i=1}^n x_{ik} y_i
 \end{aligned}$$



Regression with Matrices (cont.)

- Take partial derivatives of $g(\cdot)$ with respect to each b_j and set to zero.

$$\begin{aligned}
 b_0 \sum_{i=1}^n 1 + b_1 \sum_{i=1}^n x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} &= \sum_{i=1}^n y_i \\
 b_0 \sum_{i=1}^n x_{i1} + b_1 \sum_{i=1}^n x_{i1} x_{i1} + \dots + b_k \sum_{i=1}^n x_{i1} x_{ik} &= \sum_{i=1}^n x_{i1} y_i \\
 \dots \\
 b_0 \sum_{i=1}^n x_{ik} + b_1 \sum_{i=1}^n x_{ik} x_{i1} + \dots + b_k \sum_{i=1}^n x_{ik} x_{ik} &= \sum_{i=1}^n x_{ik} y_i
 \end{aligned}$$

- In a matrix, we can write:

$$\begin{bmatrix} \sum_{i=1}^n 1 & \sum_{i=1}^n x_{i1} & \dots & \sum_{i=1}^n x_{ik} \\ \sum_{i=1}^n x_{i1} & \sum_{i=1}^n x_{i1} x_{i1} & \dots & \sum_{i=1}^n x_{i1} x_{ik} \\ \dots & \dots & \dots & \dots \\ \sum_{i=1}^n x_{ik} & \sum_{i=1}^n x_{ik} x_{i1} & \dots & \sum_{i=1}^n x_{ik} x_{ik} \end{bmatrix} \begin{bmatrix} b_0 \\ \vdots \\ b_k \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^n y_i \\ \sum_{i=1}^n x_{i1} y_i \\ \vdots \\ \sum_{i=1}^n x_{ik} y_i \end{bmatrix}$$



Regression with Matrices (cont.)

- In a matrix, we can write:

$$\begin{bmatrix} \sum_{i=1}^n 1 & \sum_{i=1}^n x_{i1} & \dots & \sum_{i=1}^n x_{ik} \\ \sum_{i=1}^n x_{i1} & \sum_{i=1}^n x_{i1} x_{i1} & \dots & \sum_{i=1}^n x_{i1} x_{ik} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^n x_{ik} & \sum_{i=1}^n x_{ik} x_{i1} & \dots & \sum_{i=1}^n x_{ik} x_{ik} \end{bmatrix} \begin{bmatrix} b_0 \\ \vdots \\ b_k \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^n y_i \\ \sum_{i=1}^n x_{i1} y_i \\ \vdots \\ \sum_{i=1}^n x_{ik} y_i \end{bmatrix}$$

$$X'X\beta = X'y$$

Note: X' = transpose of X

$$\hat{\beta} = (X'X)^{-1}X'y$$

$$\hat{y} = X\hat{\beta}$$

After finding $b_0, b_1, b_2, \dots, b_k$, we can compute SSE, SST, R^2 ,

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 11



Department of Computer Engineering
Kasetsart University

11

Example

- Use data of 6 cars, build a model to predict horsepower (hp) using two inputs: engine size (liters) and fuel type

Make	hp	Engine size	Fuel
Ford	132	2.0	Regular
Mazda	167	2.0	Premium
Subaru	170	2.5	Regular
Lexus	204	2.5	Premium
Mitsubishi	230	3.0	Regular
BMW	260	3.0	Premium

- Let
 - x_1 = Engine size
 - x_2 = Fuel (replace 0 for regular, 1 for premium)
 - Y = hp

Source: [1]

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 12



Department of Computer Engineering
Kasetsart University

12

Example (cont.)

- $n = 6$
- $k = 2$

Make	hp	Engine size	Fuel
Ford	132	2.0	Regular
Mazda	167	2.0	Premium
Subaru	170	2.5	Regular
Lexus	204	2.5	Premium
Mitsubishi	230	3.0	Regular
BMW	260	3.0	Premium

$$\hat{\beta} = (X'X)^{-1}X'y$$

$$X = \begin{bmatrix} 1 & 2.0 & 0 \\ 1 & 2.0 & 1 \\ 1 & 2.5 & 0 \\ 1 & 2.5 & 1 \\ 1 & 3.0 & 0 \\ 1 & 3.0 & 1 \end{bmatrix} \quad y = \begin{bmatrix} 132 \\ 167 \\ 170 \\ 204 \\ 230 \\ 260 \end{bmatrix} \quad X'X = \begin{bmatrix} 6 & 15 & 3 \\ 15 & 38.5 & 7.5 \\ 3 & 7.5 & 3 \end{bmatrix}$$

$$X'y = \begin{bmatrix} 1163 \\ 3003 \\ 631 \end{bmatrix}$$

Source: [1]

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 13



Department of Computer Engineering
Kasetsart University

13

Example (cont.)

$$\hat{\beta} = (X'X)^{-1}X'y$$

$$X = \begin{bmatrix} 1 & 2.0 & 0 \\ 1 & 2.0 & 1 \\ 1 & 2.5 & 0 \\ 1 & 2.5 & 1 \\ 1 & 3.0 & 0 \\ 1 & 3.0 & 1 \end{bmatrix} \quad y = \begin{bmatrix} 132 \\ 167 \\ 170 \\ 204 \\ 230 \\ 260 \end{bmatrix} \quad X'X = \begin{bmatrix} 6 & 15 & 3 \\ 15 & 38.5 & 7.5 \\ 3 & 7.5 & 3 \end{bmatrix}$$

$$X'y = \begin{bmatrix} 1163 \\ 3003 \\ 631 \end{bmatrix}$$

$$\hat{\beta} = (X'X)^{-1}X'y = \begin{bmatrix} 79/12 & -5/2 & -1/3 \\ -5/2 & 1 & 0 \\ -1/3 & 0 & 2/3 \end{bmatrix} \begin{bmatrix} 1163 \\ 3003 \\ 631 \end{bmatrix}$$

$$\hat{\beta} = \begin{bmatrix} \hat{\beta}_0 \\ \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} = \begin{bmatrix} -61.42 \\ 95.50 \\ 33.00 \end{bmatrix}$$

$$y = -61.42 + 95.5x_1 + 33.0x_2$$

b_1, b_2 are not zero.
Both factors (engine size, fuel) affect horsepower (hp).

Source: [1]

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 14



Department of Computer Engineering
Kasetsart University

14

Example (cont.)

$$y = -61.42 + 95.5x_1 + 33.0x_2$$

What is SSE?
SST?
R²?

- Let
 - x_1 = Engine size
 - x_2 = Fuel (replace 0 for regular, 1 for premium)
 - Y = hp
- If fuel has no effect, we increase engine size by one, horsepower is increased by 95.5
- If engine size has no effect, we increase fuel by one (change from regular to premium), horsepower is increased by 33.

Source: [1]

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 15



Department of Computer Engineering
Kasetsart University

15

Outline

- Introduction
- Estimating Parameters
- Residual, SSE, SST, R²
- Example

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 16



Department of Computer Engineering
Kasetsart University

16

Residuals, Sum of Errors (SSE), σ^2

$$SSE = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

$$\sigma^2 = \frac{SSE}{n-2}$$

$$\hat{\beta} = (X'X)^{-1}X'y$$

$$\hat{y} = X\hat{\beta}$$

How to compute residuals ?

$$\text{Residual} = y - \hat{y} = y - X\hat{\beta}$$

How to compute SSE ?

$$SSE = (y - \hat{y})(y - \hat{y}) = \|y - \hat{y}\|^2$$

$$\sigma^2 = \frac{SSE}{[n-(k+1)]} = MSE$$

How to compute σ^2 ?

What is σ^2 ?

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 17

Department of Computer Engineering
Kasetsart University

17

Example (cont.)

$$SSE = \|y - \hat{y}\|^2$$

$$\hat{y} = X\hat{\beta} = \begin{bmatrix} 1 & 2.0 & 0 \\ 1 & 2.0 & 1 \\ 1 & 2.5 & 0 \\ 1 & 2.5 & 1 \\ 1 & 3.0 & 0 \\ 1 & 3.0 & 1 \end{bmatrix} \begin{bmatrix} -61.42 \\ 95.50 \\ 33.00 \end{bmatrix} = \begin{bmatrix} 129.58 \\ 162.58 \\ 177.33 \\ 210.33 \\ 225.08 \\ 258.08 \end{bmatrix}$$

$$\text{Residual} = y - \hat{y} = \begin{bmatrix} 132 \\ 167 \\ 170 \\ 204 \\ 230 \\ 260 \end{bmatrix} - \begin{bmatrix} 129.58 \\ 162.58 \\ 177.33 \\ 210.33 \\ 225.08 \\ 258.08 \end{bmatrix} = \begin{bmatrix} 2.42 \\ 4.42 \\ -7.33 \\ -6.33 \\ 4.92 \\ 1.92 \end{bmatrix}$$

$$SSE = \|y - \hat{y}\|^2 = 2.42^2 + \dots + 1.92^2 = 147.08$$

$$X = \begin{bmatrix} 1 & 2.0 & 0 \\ 1 & 2.0 & 1 \\ 1 & 2.5 & 0 \\ 1 & 2.5 & 1 \\ 1 & 3.0 & 0 \\ 1 & 3.0 & 1 \end{bmatrix}$$

$$\hat{\beta} = \begin{bmatrix} \hat{\beta}_0 \\ \hat{\beta}_1 \\ \hat{\beta}_2 \end{bmatrix} = \begin{bmatrix} -61.42 \\ 95.50 \\ 33.00 \end{bmatrix}$$

$$y = \begin{bmatrix} 132 \\ 167 \\ 170 \\ 204 \\ 230 \\ 260 \end{bmatrix}$$

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 18

Department of Computer Engineering
Kasetsart University

18

Example (cont.)

$$\sigma^2 = \frac{SSE}{[n-(k+1)]} = MSE$$

$$SSE = ||\mathbf{y} - \hat{\mathbf{y}}||^2 = 2.42^2 + \dots + 1.92^2 = 147.08$$

$$\sigma^2 = \frac{SSE}{[n-(k+1)]} = \frac{147.08}{[6-(2+1)]} = 49.03 \rightarrow \sigma = \sqrt{49.03} = 7.002$$

data = 6

2 factors: x_1, x_2

Make	hp	Engine size	Fuel
Ford	132	2.0	Regular
Mazda	167	2.0	Premium
Subaru	170	2.5	Regular
Lexus	204	2.5	Premium
Mitsubishi	230	3.0	Regular
BMW	260	3.0	Premium

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 19



Department of Computer Engineering
Kasetsart University

19

Total Sum of Error (SST) and Regression Sum of Squares (SSR)

$$SST = (\mathbf{y} - \bar{\mathbf{y}})(\mathbf{y} - \bar{\mathbf{y}}) = ||\mathbf{y} - \bar{\mathbf{y}}||^2$$

$$SST = ||\mathbf{y} - \bar{\mathbf{y}}||^2$$

$$= [(\mathbf{y} - \hat{\mathbf{y}}) + (\hat{\mathbf{y}} - \bar{\mathbf{y}})]'[(\mathbf{y} - \hat{\mathbf{y}}) + (\hat{\mathbf{y}} - \bar{\mathbf{y}})]$$

$$= ||\mathbf{y} - \hat{\mathbf{y}}||^2 + ||\hat{\mathbf{y}} - \bar{\mathbf{y}}||^2$$

Other terms are zeros

Note that $SST = SSE + SSR$

$$SSR = ||\hat{\mathbf{y}} - \bar{\mathbf{y}}||^2 = \text{Regression sum of squares} \\ = \text{Sum of squared fitted values from } \bar{\mathbf{y}}$$

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 20



Department of Computer Engineering
Kasetsart University

20

Coefficient of Determination (R^2)

$$R^2 = 1 - \frac{SSE}{SST}$$

Proportion of fitted values that can be explained by multiple linear model



Example (cont.)

$$SST = ||\mathbf{y} - \bar{\mathbf{y}}||^2$$

$$R^2 = 1 - \frac{SSE}{SST}$$

$$SST = ||\mathbf{y} - \bar{\mathbf{y}}||^2 = \sum_{i=1}^n (y_i - 193.83)^2 = 10,900.83$$

$$SSE = ||\mathbf{y} - \hat{\mathbf{y}}||^2 = 2.42^2 + \dots + 1.92^2 = 147.08$$

$$SSR = SST - SSE = 10,573.75$$

$$R^2 = 1 - \frac{SSE}{SST} = 1 - \frac{147.08}{10900.83} = 0.9865$$

98.65% of fitted values that can be explained by multiple linear model

Make	hp
Ford	132
Mazda	167
Subaru	170
Lexus	204
Mitsubishi	230
BMW	260

$$\bar{y} = 193.83$$



Outline

- Introduction
- Estimating Parameters
- Residual, SSE, SST, R^2
- Example



Example 2

- What affects human wingspan? Height? Foot length?
- Use the given data to find the followings:
 - σ , σ^2
 - SSE
 - R^2
 - SST, SSR
 - MSR, MSE

$$\sigma^2 = \frac{SSE}{[n - (k + 1)]}$$

$$R^2 = 1 - \frac{SSE}{SST}$$

$$SSR = SST - SSE$$

$$n = 16, k = 2$$



Example 2 (cont.)

- What affects human wingspan? Height? Foot length?
- Use the given data to find the followings:
 - At $\alpha = 0.05$, does height affect wingspan?
 - At $\alpha = 0.05$, does foot length affect wingspan?



Example 2 (cont.)

- Use the given data to find the followings:
 - At $\alpha = 0.05$, does height affect wingspan (not consider foot length)?



Example 2 (cont.)

- Compute correlation between
 - Wingspan and height (r_1)
 - Wingspan and foot length (r_2)
 - Height and foot length (r_3)

$$r = \frac{S_{xy}}{\sqrt{S_{xx}}\sqrt{S_{yy}}} \\ = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}$$

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 27



Department of Computer Engineering
Kasetsart University

27

References

1. J.L. Devore and K.N. Berk, Modern Mathematical Statistics with Applications, Springer, 2012.

Supaporn Erjongmanee
fengspe@ku.ac.th

Statistics in Computer Engineering
Slide 28



Department of Computer Engineering
Kasetsart University

28